

**This item is the archived peer-reviewed author-version of:**

Changes in trauma-related emotions following treatment with dialectical behavior therapy for posttraumatic stress disorder after childhood abuse

**Reference:**

Görg Nora, Böhnke Jan R., Priebe Kathlen, Rausch Sophie, Wekenmann Stefanie, Ludäscher Petra, Bohus Martin, Kleindienst Nikolaus.- Changes in trauma-related emotions following treatment with dialectical behavior therapy for posttraumatic stress disorder after childhood abuse  
Journal of traumatic stress - ISSN 1573-6598 - 32:5(2019), p. 764-773  
Full text (Publisher's DOI): <https://doi.org/10.1002/JTS.22440>  
To cite this reference: <https://hdl.handle.net/10067/1812250151162165141>

## **Changes in Trauma-related Emotions Following Treatment with Dialectical Behavior Therapy for Posttraumatic Stress Disorder after Childhood Abuse**

Nora Görg<sup>a,b</sup>, Jan R. Böhnke<sup>c</sup>, Kathlen Priebe<sup>a</sup>, Sophie Rausch<sup>a</sup>, Stefanie Wekenmann<sup>d</sup>, Petra Ludäscher<sup>a</sup>, Martin Bohus<sup>a,e</sup>, & Nikolaus Kleindienst<sup>a</sup>

<sup>a</sup>Institute of Psychiatric and Psychosomatic Psychotherapy, Central Institute of Mental Health Mannheim, J5, D-68159 Mannheim, Germany / Medical Faculty Mannheim, Heidelberg University, Germany

<sup>b</sup>Department of Clinical Psychology and Psychotherapy, Institute of Psychology, Goethe University, Frankfurt am Main, Germany

<sup>c</sup>Dundee Centre for Health and Related Research, School of Nursing and Health Sciences, University of Dundee, Dundee DD1 4HJ, UK

<sup>d</sup>Department of Psychiatry and Psychotherapy, University of Tübingen, Calwerstr. 14, 72076 Tübingen, Germany

<sup>e</sup>Collaborative Antwerp Psychiatric Research Institute, University of Antwerp, Universiteitsplein 1, 2610, Wilrijk, Antwerp, Belgium

**Author accepted manuscript to be published in the Journal of Traumatic Stress.**

**Corresponding author:**

Nora Görg  
ZI Mannheim  
J5  
D-68159 Mannheim  
Germany  
Phone: 0049 621 1703 4412  
Fax: 00621-1703 4005  
Email: [nora.goerg@zi-mannheim.de](mailto:nora.goerg@zi-mannheim.de)

### Abstract

Dialectical Behavior Therapy for Posttraumatic Stress Disorder (DBT-PTSD) is a trauma-focused therapy effectively reducing core PTSD symptoms such as intrusions, hyperarousal, and avoidance. Preliminary data indicate effects on elevated trauma-related emotions (e.g., guilt and shame) and possibly radical acceptance of the traumatic event. However, it is unclear if improvements in these variables are significant after controlling for changes in core PTSD symptoms and to which extent non-clinical levels are obtained. Forty-two individuals meeting the criteria for PTSD after childhood abuse participating in a three-month residential DBT-PTSD program were evaluated at the start of the exposure phase and at the end of the treatment. A non-clinical sample with a history of childhood abuse served as a reference for the investigated outcomes. Multivariate analyses of variance and multivariate analyses of covariance controlling for change in core PTSD symptoms were used to evaluate changes in a broad range of elevated trauma-related emotions (fear, anger, guilt, shame, disgust, sadness, and helplessness) and in radical acceptance. Both elevated trauma-related emotions and radical acceptance significantly improved during DBT-PTSD ( $\Lambda = 0.34$ ,  $p < 0.001$ ,  $\eta^2 = 0.56$ ;  $t_{40} = -5.66$ ,  $p < 0.001$ ,  $SMD = 0.88$ ) even after controlling for changes in PTSD symptoms ( $\Lambda = 0.35$ ,  $p < 0.001$ ,  $\eta^2 = 0.65$ ;  $\Lambda = 0.86$ ,  $p = 0.018$ ,  $\eta^2 = 0.14$ ). At the end of DBT-PTSD between 76.2% (guilt) and 31.0% (acceptance) of patients showed non-clinical levels of the investigated outcomes. To conclude, both trauma-related emotions and radical acceptance changed after the three-month residential DBT-PTSD program.

When Posttraumatic Stress Disorder (PTSD) was introduced in the Diagnostic and Statistical Manual of Mental Disorders (DSM) (American Psychiatric Association, 1980), it was classified as an anxiety disorder. In recent years, a broader range of emotions that are elevated in PTSD have been considered relevant. Consequently, PTSD was reclassified in the DSM-5, moving it from the anxiety disorders to a new category of trauma- and stressor-related disorders. The core symptom clusters are re-experiencing, avoidance, and hyperarousal. These constitute previous diagnoses of PTSD and were complemented by a diagnostic criterion comprising persistent negative affect, inability to experience positive affect, and distorted cognitions like self-blame (American Psychiatric Association, 2013). The ICD-11 proposes two distinct sibling conditions, PTSD and Complex PTSD (CPTSD), under the category ‘disorders specifically associated with stress’ (WHO, 2018). For a diagnosis of CPTSD the core symptoms of PTSD are required in addition to the following: 1) severe and pervasive problems in affect regulation; 2) persistent beliefs about oneself as diminished, defeated or worthless, accompanied by deep and pervasive feelings of shame, guilt or failure related to the traumatic event; and 3) persistent difficulties in sustaining relationships and in feeling close to others (WHO, 2018). In a recent study the highest conditional prevalence for PTSD was associated with kidnapping or rape, and the highest CPTSD rates were associated with childhood sexual abuse (CSA) (Maercker, Hecker, Augsburger, & Kliem, 2018).

The relevance of non-fear emotions in modern conceptualizations of PTSD is well supported: high levels of shame, guilt, anger, and disgust correlate with increased re-experiencing, avoidance, and hyperarousal (Badour, Feldner, Blumenthal, & Knapp, 2013; Orth & Wieland, 2006; Pugh, Taylor, & Berry, 2015; Rupp, Doebler, Ehring, & Vossbeck-Elsebusch, 2017; Saraiya & Lopez-Castro, 2016; Street, Gibson, & Holohan, 2005); elevated levels of shame and guilt are also associated with suicidal ideation (Bryan, Morrow, Etienne, & Ray-Sannerud, 2013; You, Talbot, He, & Conner, 2012).

Core symptoms of PTSD can be effectively treated by trauma-focused treatments: Prolonged Exposure (PE), Narrative Exposure Therapy (NET), Cognitive Processing Therapy (CPT), and Eye Movement and Desensitization and Reprocessing (EMDR) (Cusack et al., 2016; Watts et al., 2013). However, while meta-analyses on trauma-focused treatment showed large reductions in PTSD symptoms across different subgroups (Cusack et al., 2016; Watts et al., 2013), meta-analyses specifically studying treatment effects in patients with PTSD related to childhood abuse (CA) yielded only moderate effect sizes (Ehring et al., 2014) and low recovery rates (Dorrepaal et al., 2014). Furthermore, studies on CA-related PTSD often excluded patients with substance abuse, dissociative disorders, borderline personality disorder (BPD), or suicidality. Thus, specialized treatments for CA-related PTSD are clearly needed—especially for patients with complex symptoms including emotion dysregulation and comorbidities such as BPD (Dorrepaal et al., 2014). Treatments for abuse-related PTSD (Cloitre, Petkova, Wang, & Lu, 2012; Harned, Korslund, & Linehan, 2014; Steil, Dyer, Priebe, Kleindienst, & Bohus, 2011) emphasize the use of emotion regulation strategies. In particular, Dialectical Behavior Therapy for Complex Posttraumatic Stress Disorder (DBT-PTSD) was tailored for patients with CPTSD related to CA, i.e. for patients with severe emotion dysregulation, high levels of guilt, shame, self-contempt, self-harm, suicidal ideation, dissociation, and interpersonal problems (Bohus et al., *in press*). The treatment principles further described by Bohus and Priebe (2018) are as follows: DBT-PTSD is a multi-component treatment following algorithms derived from DBT (Linehan, 1993). It includes clear structures and a dynamic hierarchization of treatment focuses. A major element of DBT are so-called “skills”, i.e. short mental self-instructions aiming at interrupting and modifying automatic appraisals and behavioral patterns. Skills can be further used to manage extreme levels of distress that are often associated with dissociation. In patients with emotion regulation difficulties, imagination of stressful life events often triggers dissociation that hampers

emotional learning (Krause-Utz et al., 2018). Accordingly, imaginal exposure was supplemented by anti-dissociative skills (skills-based exposure). Skills are further used to modulate over-activated emotions. Since standard DBT does not include elaborated trauma-specific interventions in sufficient detail, we supplemented DBT with trauma-focused cognitive and exposure-based interventions (Ehlers, Clark, Hackmann, McManus, & Fennell, 2005).

As outlined above, patients with CPTSD after CA suffer from severe negative self-concepts accompanied by deep and pervasive trauma-related emotions like guilt or shame. In accordance with Brewin (1996), we conceptualize these emotions as secondary emotions that are consequences of post-traumatic appraisals of the event. Assumptions like “it was all my fault”, “I deserve what happened”, or “something is wrong with me” are pseudo-causal explanations that may help to explain the inexplicable in the short-term. Guilt and shame might also help to decrease primary emotions that were present during the traumatic event such as helplessness, disgust, or fear. Accordingly, we assume that imaginal exposure of trauma-related primary lead to a better acceptance of these emotions. The emotional experience that the trauma - and its consequences - were not “my fault” should also lead to a significant decrease of secondary emotions like guilt or shame.

Furthermore, patients learn to process automatic thoughts and maladaptive trauma-related emotions from a more distant perspective. This leads to more control over the emotion-driven action tendencies and their replacement with functional behavior. Specific techniques used in DBT-PTSD are derived from both Acceptance and Commitment Therapy (ACT) (Hayes, Strosahl, & Wilson, 2011) and Compassion-Focused Therapy (CFT) (Gilbert, 2014). ACT provides helpful interventions for the recognition and implementation of values that improve the quality of life. Because the self-concept is often characterized by trauma-related emotions such as shame, guilt, self-hate, or disgust, many patients have difficulties dealing with themselves in a sympathetic and self-valuing manner. This is also frequently

reflected in interpersonal problems. In CFT, these difficulties are addressed through training of a compassionate perspective towards oneself and other people.

Both a pilot study (Steil et al., 2011) and an RCT (Bohus et al., 2013) showed the general safety and efficacy of DBT-PTSD. In the RCT, between-group effect-sizes with respect to the CAPS (i.e. PTSD core symptoms) were large ( $SMD = 1.35$  vs a treatment-as-usual wait list condition) and statistically significant (Bohus et al., 2013). Pre- to post-treatment changes in radical acceptance—an important aspect of DBT-PTSD—have only been investigated in one study (Görg et al., 2017) yielding a moderate increase in radical acceptance from pre- to post-DBT-PTSD ( $SMD = 0.42$ ). Empirical studies on DBT-PTSD and other trauma-focused treatments confirmed the change in non-fear-emotions including guilt and shame. With respect to trauma-related *guilt*, pre-to-post treatment decreases were reported in DBT-PTSD after CSA (Görg et al., 2017) in trauma- or present-focused group therapy for CSA (Ginzburg et al., 2009), in cognitive behavior therapy for women with intimate partner violence (Allard, Norman, Thorp, Browne, & Stein, 2016), in CPT and PE for female rape victims (Nishith, Nixon, & Resick, 2005), and in CPT for veterans (Held, Owens, Monroe, & Chard, 2017). These findings are complemented by other studies that reported decreases in trauma-related guilt in mixed PTSD samples receiving PE (Langkaas et al., 2017; Stapleton, Taylor, & Asmundson, 2006), Imagery Rescripting (IR) (Langkaas et al., 2017), EMDR (Stapleton et al., 2006), relaxation training (Stapleton et al., 2006), or imaginal exposure enhanced by IR (Arntz, Tiesema, & Kindt, 2007). Consistent with that, trauma-related self-blame changed significantly from pre- to post-IR for posttraumatic nightmares in a mixed trauma sample (Long et al., 2011). While non-significant, it tended to decrease in a sample with mixed trauma types receiving PE (Kumpula et al., 2017). Similarly, a study by Harned, Ruork, Liu, and Tkachuck (2015) reported significant decreases in both guilt and shame during exposure sessions from the start to the end of the exposure phase. The treat-

ment in this study combined DBT and PE for women with PTSD and BPD. This finding of decreased *shame* coincided with results reported for DBT-PTSD for CSA (Görg et al., 2017) as well as for PE and IR in a sample of PTSD patients after mixed traumatic events (Langkaas et al., 2017). Through these treatments, trauma-related shame decreased from pre- to post-treatment. Further studies investigated changes in different aspects of *anger*: Stapleton et al. (2006) found significant decreases in trauma-related anger in all treatment conditions (PE; EMDR; relaxation training) in a mixed trauma sample. Langkaas et al. (2017) also reported that internalized anger decreased significantly after both PE and IR. Consistent with these findings, in-session anger decreased significantly from pre- to post-exposure therapy in a military sample (Pitman, Orr, Altman, & Longpre, 1996). In patients with abuse-related PTSD, state anger decreased significantly after CPT (Galovski, Elwood, Blain, & Resick, 2014), Stress Inoculation Training, and PE (Cahill, Rauch, Hembree, & Foa, 2003), there were no significant reductions after PE in another study (Foa, Riggs, Massie, & Yarczower, 1995). Significant reductions in anger expression were observed after exposure therapy for PTSD related to CA (Cloitre et al., 2010). Finally, reductions in *sadness* were observed after exposure therapy in the military sample of Pitman et al. (1996) as were reductions in trauma-related *disgust* after DBT-PTSD for CSA (Görg et al., 2017).

However, despite the evidence indicating that trauma-related emotions decreased from pre- to post-treatment, the evidence for a substantial impact of treatments for CA-related PTSD on these emotions is limited. The few treatments that focus on PTSD related to CA remain inconclusive. While anger was investigated by Cloitre et al. (2010), shame and guilt were not. On the other hand, the evidence from the pilot study by Harned et al. (2014) is still very limited as for “the small sample size, which was further complicated by the higher than usual drop-out rate”. While studies on DBT-PTSD suggested strong evidence for its general efficacy (Bohus et al., 2013; Steil et al., 2011), the potential impact on trauma-related emo-



tions and radical acceptance was only investigated in one small study (Görg et al., 2017), and this requires replication from an independent sample. Furthermore, it is unclear to what extent the trauma-related emotions post-treatment correspond to a non-clinical level.

Approaches proposed for comparing patients' scores with non-clinical values are either exclusively based on values observed in the non-clinical reference group or rely on relative approaches based on values in the patient and non-clinical reference groups (for a review see Bauer, Lambert, & Nielsen, 2004; Nasiakos, Cribbie, & Arpin-Cribbie, 2010). In the first type, an equivalence range (e.g., a multiple of the standard deviation around the mean in the non-clinical population) is defined. Particular scores falling within this range of equivalence are considered to be non-clinical, and scores outside this range are considered clinical (Bauer et al., 2004; Kendall, Marrs-Garcia, Nath, & Sheldrick, 1999; Nasiakos et al., 2010). An alternative approach is based on the relative probabilities of belonging to the patient group vs. the non-clinical group (Bauer et al., 2004; Jacobson & Truax, 1991) such as "criterion C" proposed by Jacobson and Truax (1991). This has been recommended for its robustness and empirically-supported validity.

Previous research on trauma-related emotions has not addressed the role of core PTSD symptoms. The literature summarized above does not exclude the possibility that therapies for PTSD primarily impact core symptoms while changes in trauma-related emotions may be an epiphenomenon related to modification of PTSD core symptoms. Clinically, it would be important to know whether trauma-related emotions can be addressed by treating core PTSD symptoms or—as recommended by Cloitre et al., 2012—additionally require specific interventions aiming at directly addressing dysfunctional emotions.

In summary, previous studies of trauma-focused treatments including DBT-PTSD reported decreases in several trauma-related emotions. However, important points have not been investigated or remain unclear. These points include the potentially confounding role of core

PTSD symptoms, a comparison of trauma-related emotions post-treatment with non-clinical populations, and further data on change in radical acceptance.

### **Aims of this study**

This study assessed the levels of trauma-related emotions in PTSD (fear, anger, guilt, shame, disgust, sadness, and helplessness), and radical acceptance of the traumatic event at the start of the exposure phase and at the end of residential DBT-PTSD. Our research had three aims. First, to replicate and extend the respective results of Görg et al. (2017) in a larger sample: We investigated whether trauma-related emotions and radical acceptance change after DBT-PTSD. Second, we tested whether the hypothesized changes remain significant after controlling for a change in core PTSD symptoms measured with the Davidson Trauma Scale (DTS) (Davidson et al., 1997). Finally, we aimed to test how many patients show a non-clinical level after DBT-PTSD with respect to acceptance and trauma-related emotions.

## **Methods**

### **Participants**

The patients were recruited from a three-month residential DBT-PTSD treatment program at the Central Institute of Mental Health, Mannheim, Germany. Patients had to meet the diagnosis of PTSD after CA. Exclusion criteria were acute alcohol or drug dependence, schizophrenia, bipolar disorder, and medical contraindications to exposure (e.g., body mass index  $\leq 16$ ). All participants gave written informed consent after three weeks in treatment. At this time-point, patients committed to an individualized treatment plan including exposure. By then, patients had also received sufficient psychoeducation on the trauma-related emotions that were believed to be beneficial for the data quality due to frequent alexithymia in PTSD (Becirovic, Avdibegovic, Softic, Mirkovic-Hajdukovic, & Becirovic, 2017). The study was conducted according to the Declaration of Helsinki and was approved by the Ethics

Committee of the Medical Faculty Mannheim, Heidelberg University. The trial was registered at the WHO Clinical Trials Registry Platform (DRKS00006226).

There were 110 patients treated during the study period, but 28 patients were not recruited because they met exclusion criteria ( $n = 1$ ), received other treatments ( $n = 20$ ), had insufficient German literacy ( $n = 1$ ), or early drop-out ( $n = 6$ ). Of the 82 patients who were informed about the study, 60 patients agreed to participate. Of these, 18 were excluded (missing data,  $n = 4$ , drop out,  $n = 1$ , exclusion criteria fulfilled,  $n = 2$ , inclusion criteria not fulfilled,  $n = 3$ , not receiving full DBT-PTSD,  $n = 8$ ). The final sample included 42 patients (90.5% female) with a mean age of 42.17 ( $SD = 9.64$ ) years and 10.10 ( $SD = 1.14$ ) years of education. No significant differences between the patients from the final sample and those who did not participate were detected. On average, patients provided the first questionnaire after 34.19 days of the usual 90 days in treatment ( $SD = 9.12$ , median = 32.50). This corresponds with the exposure-based interventions, which comprise the treatment phase of interest as in this study.

On average, patients had a total of 2.00 ( $SD = 1.01$ ) current co-occurring disorders in addition to PTSD. The most frequent comorbidity was major depressive disorder affecting 73.8% ( $n = 31$ ) of the sample followed by BPD (35.7%,  $n = 15$ ). Two patients (4.8%) were each diagnosed with dysthymia or somatoform disorder. Altogether, 90.5% ( $n = 38$ ) of patients had been sexually abused—78.6% of the total sample ( $n = 33$ ) before the age of 18. Physical abuse was reported by 81.0% ( $n = 34$ ) of patients with 78.6% of the total sample ( $n = 33$ ) before the age of 18. Overall, 71.4% ( $n = 30$ ) of patients had been exposed to both physical and sexual abuse at some point in their lives. A total of 90.5% received psychotropic medication—mainly antidepressants (85.7%, mostly SSRIs). Other sample characteristics are presented in Table 1.

## **Procedure**

Patients were treated between 2014 and 2016 at the specialized trauma-unit of the Central Institute for Mental Health, Mannheim. All participants received residential DBT-PTSD, which has been described in detail elsewhere (Bohus et al., 2013; Bohus & Priebe, 2018). In short, DBT-PTSD consisted of several phases spread over 12 weeks. These phases include the following mandatory modules: psychoeducation and skills acquisition, skills-based exposure, radical acceptance, and psychosocial integration. These modules were supplemented by optional modules (e.g., to address trauma-related nightmares) that facilitates an adaptation to the diverse symptom constellations in CPTSD according to predefined rules. Skills were modified for the specific needs of traumatized clients and include: 1) mindfulness, 2) emotional regulation, 3) and distress tolerance skills. The final phase worked on personal goals that served to regain a life worth living. Supervision and consultation team meetings of therapists aimed to maintain treatment fidelity. In addition to specific trauma-focused therapy, patients could participate in general art therapy or could receive advice from social services. Treatment and assessments were provided by clinical psychologists with DBT-PTSD training who were supervised by one of the treatment developers.

## **Measures**

Inclusion and exclusion criteria as well as co-occurring diagnoses were determined with the German version of the Structured Clinical Interview for DSM-IV (Wittchen, Wunderlich, Gruschwitz, & Zaudig, 1997). In addition, diagnosis of BPD was assessed with the International Personality Disorder Examination for ICD-10 (Mombour et al., 1996). Trauma-related emotions were rated on seven visual analogue scales each ranging from 0-100. These scales assessed trauma-related shame, guilt, disgust, anger, fear, helplessness, sadness, and radical acceptance. The emotion items were: "When I think of the traumatic event now, I feel [emotion]". For acceptance, the question was: "When I think of the traumat-

ic experience now, I can accept it as part of my life story." The severity of core PTSD symptoms was assessed by the Davidson Trauma Scale (DTS; Davidson et al., 1997). It consists of 17 items rating the frequency and severity PTSD symptoms and showed good test-retest reliability (.86), good convergent validity with other PTSD symptom scales, and sensitivity to treatment response. The DTA authors defined the clinical cut-off as 40 (Davidson et al., 1997). Cronbach's  $\alpha$  of the DTS in this study was .93. Trauma history was assessed via a questionnaire that required patients to indicate if they had experienced one of the following traumatic events: sexual abuse, physical abuse, suicide, or homicide of a close person, war, imprisonment, torture, accidents, or natural disasters.

### **Data analysis**

Significance testing of changes in trauma-related emotions and radical acceptance was carried out from repeated measures multivariate analyses of variance (MANOVA). To test our second hypothesis, a repeated-measure multivariate analysis of covariance (MANCOVA) was computed, controlling for changes in the DTS. The third hypothesis explored to which degree the patients' trauma-related emotions and acceptance post treatment would show levels comparable to a non-clinical population. Since the trauma-related emotions in our study explicitly refer back to a traumatic event, reference values were sought from a sample representing a population with the same past trauma experience but not suffering from PTSD. We used the data from a subsample published by Rausch et al. (2016) who tested healthy traumatized women with a history of CA before the age of 18 without having developed PTSD, any other lifetime axis I disorder, or BPD as a reference for our sample. Although the data from the non-clinical population were not part of our study, they were sufficient to define cut-offs between clinical and non-clinical levels of emotions (and radical acceptance) via the criteria of Kendall et al. (1999) and Jacobson and Truax (1991). According to Kendall the cut-off is defined as  $mean_{reference\ group} \pm 1SD_{reference\ group}$  for each of the trauma-

related emotions and for radical acceptance, respectively, where the non-clinical data serve as the reference group. According to Jacobson & Truax (1991), the cut-off is defined as the point that is equally likely for a score to be drawn from the clinical (our patient sample at the start of the exposure phase) or the non-clinical data (the sample from Rausch et al., 2016). Effect sizes were computed as  $\eta^2$  and as standardized mean differences (*SMD*). The  $\eta^2$  can be interpreted as the proportion of explained variance; by convention, *SMDs* of 0.2, 0.5, and 0.8 correspond to small, medium, and large effects (Cohen, 1988). Individual changes for each trauma-related emotion were illustrated with Bertin plots. SPSS v. 21 was used for descriptive and inferential statistics; Bertin plots were realized in R (v. 3.1.3).

### Results

According to the MANOVA (included all trauma-related negative emotions investigated in this sample), these emotions clearly changed from the start of the exposure phase to the end of DBT-PTSD (Wilks'  $\Lambda = 0.34$ ,  $p < .001$ ,  $\eta^2 = 0.56$ ) even after controlling for the change in the total DTS score (Wilks'  $\Lambda = 0.65$ ,  $p < .001$ ,  $\eta^2 = 0.35$ ).

The decline encompassed a wide range of trauma-related emotions illustrated in Figure 1, Table 2, and Supplement 2: The change was statistically significant for all emotions except anger. The effect-sizes were large for all emotions except anger and sadness (i.e., for shame, guilt, helplessness, fear, and disgust). Average DTS scores at the start and end of DBT-PTSD were 88.30 ( $SD = 22.96$ ) and 61.97 ( $SD = 26.85$ ). Changes in all emotions but anger were positively and significantly correlated with changes in the DTS. Inter-correlations between changes in emotions were positive except for a non-significant negative association between anger and guilt (see Supplement 1). When controlling for changes in the DTS total score, changes in trauma-related emotions were significant for all variables except anger and sadness (see Table 2). Radical acceptance increased significantly over time ( $t_{40} = -5.66$ ,  $p < .001$ ,  $SMD = 0.88$ ); this remained statistically significant after controlling for changes in the

DTS total score (Wilks'  $\Lambda = 0.86$ ,  $p = .018$ ,  $\eta^2 = 0.14$ ). A post-hoc power analysis (setting  $\alpha$  at 0.05,  $1-\beta$  at 0.80, and assuming a correlation between pre- and post-measures of  $r = .50$ ) indicated that the study was adequately powered to detect effect sizes of  $\eta^2 \geq .06$ .

Supplement 2 indicates that the average levels of trauma-related emotions at the start of the exposure phase differed from respective levels in the non-clinical reference group and approached non-clinical reference values by the end of DBT-PTSD. Table 2 shows the percentage of participants' emotions falling into the non-clinical range (according to Kendall et al. (1999)) by the end of treatment ranged from 47.6% (for sadness) to 76.2% (for guilt). The respective percentages on the basis of the criterion of Jacobson and Truax (1991) ranged from 35.7% (for anger) to 81.0% (for guilt). This indicates that by the end of treatment, 81.0% of patients had achieved a level of trauma-related guilt that resembled the non-clinical control group rather than the baseline values of the patient group. With respect to radical acceptance, the criterion by Kendall et al. (1999) was fulfilled by 31.0% of the patients, and the criterion by Jacobson and Truax (1991) was fulfilled by 38.1% of the patients. The percentages of patients who met the Kendall criterion at the start of the exposure phase ranged from 7.3% to 54.8% (see Table 2).

## Discussion

### Main findings and comparison to prior research

We investigated levels of trauma-related emotions and of radical acceptance in patients with PTSD related to CA pre- and post-DBT-PTSD. Overall, patients reported statistically significant decreases in trauma-related emotions and increases in radical acceptance. Anger was the only outcome measure that did not improve in a statistically significant way. Changes in guilt, shame, disgust, fear, helplessness, and radical acceptance were significant when controlling for changes in core PTSD symptoms. This indicates that a relevant proportion of change in trauma-related emotions and radical acceptance was not driven by a change

in core PTSD symptoms. Depending on the outcome variable, 31.0 to 76.2% showed a non-clinical level after treatment (Kendall et al., 1999) i.e., post-treatment scores of these participants were similar to the traumatized but non-clinical reference sample.

The decrease in trauma-related emotions is in line with prior research. Changes were most pronounced for trauma-related shame and guilt ( $SMD = 1.5$  for both emotions). The results complement earlier studies regarding similar changes (Allard et al., 2016; Arntz, Tiesema, & Kindt, 2007; Cloitre et al., 2010; Ginzburg et al., 2009; Görg et al., 2017; Harned et al., 2015; Held et al., 2017; Langkaas et al., 2017; Long et al., 2011; Nishith et al., 2005). The large effects with respect to shame and guilt in our study may reflect the attention to these emotions in DBT-PTSD. Guilt is addressed as a secondary emotion, which helps to reduce the experience of helplessness (Bohus & Priebe, 2018). In particular, DBT-PTSD applies both cognitive interventions according to Ehlers et al. (2005) and skills according to Linehan (1993) to address shame and guilt. During skill-assisted exposure, patients were repeatedly asked to test their hypotheses about their own responsibility, their faults, and their contributions to the abuse. This procedure might explain the change in secondary guilt cognitions (e.g., “this happened since I am bad and seductive”) and related emotions (guilt and shame).

The fact that changes in trauma-related anger were least obvious may relate to functional aspects of trauma-related anger (Langkaas et al., 2017); this anger might result from a decrease in inappropriate self-blame (patients realizing that the perpetrator is responsible for the traumatic event, not themselves). The notion that a change in dysfunctional cognitions often results in appropriate anger directed to the perpetrator is substantiated in a study by Kleim, Graham, Bryant, and Ehlers (2013) in non-clinical traumatized controls with prior exposure to assault or motor vehicle accidents: These participants also reported feelings of anger in response to the intrusive experience although the level of self-reported anger was significantly lower than levels reported in patients with PTSD.



It is also not surprising that some patients experience high levels of sadness at the end of the three-month treatment. Radical acceptance of the traumatic event is frequently accompanied by appropriate sadness. Future studies should test whether sadness is reduced in follow-up studies. Significant increases in radical acceptance with a large effect size ( $SMD = 0.88$ ) extends the findings of an earlier study that reported moderate but non-significant increases in radical acceptance from the start to the end of DBT-PTSD (Görg et al., 2017).

The study further extended previous findings because it showed that a relevant proportion of change in trauma-related emotions and radical acceptance is not merely an epiphenomenal effect following changes in core PTSD symptoms. Finally, this is the first study to show that a substantial proportion of patients showed a non-clinical level of trauma-related emotions by the end of the treatment—especially for guilt and shame (76.2% and 69.0%, respectively).

### **Limitations and strengths of the study**

Although treatments for PTSD have shown general efficacy (Cusack et al., 2016; Watts et al., 2013), there is a need to better characterize efficacy profiles of available treatments. Efficacy profiles of PTSD treatments are relevant because modern conceptualizations of PTSD include a wide spectrum of conditions. Notably, the DSM-5 (American Psychiatric Association, 2013) and the ICD-11 (WHO, 2018) included non-fear emotions among diagnostic criteria. The ICD-11 conceptualization of CPTSD lists “deep and pervasive feelings of shame, guilt, or failure related to the traumatic event,” among characteristics separating CPTSD from PTSD. Shame and guilt are the trauma-related emotions that decreased the most after treatment with DBT-PTSD. Future research is clearly warranted to investigate the impact of DBT-PTSD and of other treatments for different trauma-related emotions. Patients reported different profiles of trauma-related emotions that might require specifically tailored

interventions depending on which emotions are most pronounced. Empirically-based decision rules should guide differential indications (Deisenhofer et al., 2018).

### **Implications for future research and clinical practice**

Although treatments for PTSD have shown general efficacy (Cusack et al., 2016; Watts et al., 2013), there is a need to better characterize efficacy profiles of available treatments. Efficacy profiles of PTSD treatments are relevant, since modern conceptualizations of PTSD account for a wide spectrum of conditions. Notably, the DSM-5 (American Psychiatric Association, 2013) and the ICD-11 (WHO, 2018) included non-fear emotions among diagnostic criteria. The ICD-11 conceptualization of CPTSD lists “deep and pervasive feelings of shame, guilt, or failure related to the traumatic event,” among characteristics separating CPTSD from PTSD. Shame and guilt are the trauma-related emotions which decreased most after treatment with DBT-PTSD. Future research is clearly warranted to investigate the impact of DBT-PTSD and of other treatments for different trauma-related emotions. Patients reported different profiles of trauma-related emotions which might require specifically tailored interventions depending on which emotions are most pronounced. Empirically-based decision rules should guide differential indication (Deisenhofer et al., 2018).

### **Conclusion**

Trauma-related emotions and radical acceptance changed from the start of the exposure phase to the end of the three-month residential DBT-PTSD. A high proportion of change in these variables cannot be explained by changes in core PTSD symptoms alone. The most encouraging results were observed with respect to trauma-related guilt and shame given that most patients reached a non-clinical level with respect to these emotions.

### References

- Allard, C. B., Norman, S. B., Thorp, S. R., Browne, K. C., & Stein, M. B. (2016). Mid-Treatment Reduction in Trauma-Related Guilt Predicts PTSD and Functioning Following Cognitive Trauma Therapy for Survivors of Intimate Partner Violence *Journal of Interpersonal Violence*. doi: 10.1177/0886260516636068
- American Psychiatric Association. (1980). *Diagnostic and statistical manual of mental disorders (3rd ed.)*. Washington, DC: American Psychiatric Association.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th ed.)*. Washington, DC: American Psychiatric Association.
- Arntz, A., Tiesema, M., & Kindt, M. (2007). Treatment of PTSD: A comparison of imaginal exposure with and without imagery rescripting. *Journal of Behavior Therapy and Experimental Psychiatry*, 38(4), 345-370. doi:10.1016/j.jbtep.2007.10.006
- Badour, C. L., Feldner, M. T., Blumenthal, H., & Knapp, A. (2013). Preliminary Evidence for a Unique Role of Disgust-Based Conditioning in Posttraumatic Stress. *Journal of Traumatic Stress*, 26(2), 280-287. doi:10.1002/jts.21796
- Bauer, S., Lambert, M. J., & Nielsen, S. L. (2004). Clinical significance methods: a comparison of statistical techniques. *Journal of Personality Assessment*, 82(1), 60-70.
- Becirovic, E., Avdibegovic, E., Softic, R., Mirkovic-Hajdukovic, M., & Becirovic, A. (2017). Alexithymia in war veterans with post-traumatic stress disorder. *European Psychiatry*, 41, S720. doi:https://doi.org/10.1016/j.eurpsy.2017.01.1301
- Bohus, M., Dyer, A. S., Priebe, K., Krüger, A., Kleindienst, N., . . . Steil, R. (2013). Dialectical Behaviour Therapy for Post-traumatic Stress Disorder after Childhood Sexual Abuse in Patients with and without Borderline Personality Disorder: A Randomised Controlled Trial. *Psychotherapy and Psychosomatics*, 82(4), 221-233. doi:10.1159/000348451

- Bohus, M., & Priebe, K. (2018). DBT–PTSD: A Treatment Programme for Complex PTSD After Childhood Abuse. In M. A. Swales (Ed.), *The Oxford Handbook of Dialectical Behaviour Therapy*. Oxford: Oxford University Press.
- Bohus, M., Schmahl, C., Fydrich, T., Steil, R., Müller-Engelmann, M., Herzog, J., Ludäscher, P. Kleindienst, N. Priebe, K. (in press). A Research Programme to Evaluate DBT-PTSD, a Modular Treatment Approach for Complex PTSD after Childhood Abuse. *Borderline Personality Disorder and Emotion Dysregulation*.
- Brewin, C. R., Dalgleish, T., & Joseph, S. (1996). A Dual Representation Theory of Post traumatic Stress Disorder. *Psychological Review*, 103(4), 670-686.
- Bryan, C. J., Morrow, C. E., Etienne, N., & Ray-Sannerud, B. (2013). Guilt, shame, and suicidal ideation in a military outpatient clinical sample. *Depression and Anxiety*, 30(1), 55-60.
- Cahill, S. P., Rauch, S. A., Hembree, E. A., & Foa, E. B. (2003). Effect of Cognitive-Behavioral Treatments for PTSD on Anger. *Journal of Cognitive Psychotherapy*, 17(2), 113-131. doi:10.1891/jcop.17.2.113.57434
- Cloitre, M., Petkova, E., Wang, J., & Lu, F. (2012). An examination of the influence of a sequential treatment on the course and impact of dissociation among women with PTSD related to childhood abuse. *Depression and Anxiety*, 29, 709-717.
- Cloitre, M., Stovall-McClough, K. C., Nooner, K., Zorbas, P., Cherry, S., & Jackson, C. L. (2010). Treatment for PTSD related to childhood abuse: a randomized controlled trial. *American Journal of Psychiatry*, 167(8), 915-924.
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences (2nd ed.)*. Hillsdale: Lawrence Erlbaum Associates.

Cusack, K., Jonas, D. E., Forneris, C. A., Wines, C., Sonis, J., . . . Gaynes, B. N. (2016).

Psychological treatments for adults with posttraumatic stress disorder: A systematic review and meta-analysis. *Clinical Psychology Review, 43*(Supplement C), 128-141.

Davidson, J. R. T., Book, S. W., Colket, J. T., Tupler, L. A., Roth, S., David, D., . . .

Feldman, M. (1997). Assessment of a new self-rating scale for post-traumatic stress disorder. *Psychological Medicine, 27*, 153-160.

Deisenhofer, A., Delgadillo, J., Rubel, J. A., Böhnke, J. R., Zimmermann, D., Schwartz, B.,

& Lutz, W. (2018). Individual treatment selection for patients with posttraumatic stress disorder. *Depression and Anxiety, 35*(6), 541-550. doi:10.1002/da.22755

Dorrepaal, E., Thomaes, K., Hoogendoorn, A. W., Veltman, D. J., Draijer, N., & van

Balkom, A. J. L. M. (2014). Evidence-based treatment for adult women with child abuse-related Complex PTSD: a quantitative review. *European Journal of Psychotraumatology, 5*. doi:10.3402/ejpt.v5.23613

Ehlers, A., Clark, D. M., Hackmann, A., McManus, F., & Fennell, M. (2005). Cognitive therapy for post-traumatic stress disorder: development and evaluation. *Behaviour Research and Therapy, 43*, 413-431. doi:10.1016/j.brat.2004.03.006

Ehring, T., Welboren, R., Morina, N., Wicherts, J. M., Freitag, J., & Emmelkamp, P. M. G.

(2014). Meta-analysis of psychological treatments for posttraumatic stress disorder in adult survivors of childhood abuse. *Clinical Psychology Review, 34*(8), 645-657.

Foa, E. B., Riggs, D. S., Massie, E. D., & Yarczower, M. (1995). The impact of fear

activation and anger on the efficacy of exposure treatment for posttraumatic stress disorder. *Behavior Therapy, 26*(3), 487-499. doi:10.1016/s0005-7894(05)80096-6

Galovski, T. E., Elwood, L. S., Blain, L. M., & Resick, P. A. (2014). Changes in Anger in

Relationship to Responsivity to PTSD Treatment. *Psychological trauma : theory, research, practice and policy, 6*(1), 56-64. doi:10.1037/a0031364

- Gilbert, P. (2014). The origins and nature of compassion focused therapy. *British Journal of Clinical Psychology, 53*(1), 6-41. doi:10.1111/bjc.12043
- Ginzburg, K., Butler, L. D., Giese-Davis, J., Cavanaugh, C. E., Neri, E., Koopman, C., . . . Spiegel, D. (2009). Shame, guilt, and posttraumatic stress disorder in adult survivors of childhood sexual abuse at risk for human immunodeficiency virus: outcomes of a randomized clinical trial of group psychotherapy treatment. *Journal of Nervous and Mental Disease, 197*(7), 536-542. doi:10.1097/NMD.0b013e3181ab2ebd
- Görg, N., Priebe, K., Böhnke, J. R., Steil, R., Dyer, A. S., & Kleindienst, N. (2017). Trauma-related emotions and radical acceptance in dialectical behavior therapy for posttraumatic stress disorder after childhood sexual abuse. *Borderline Personality Disorder and Emotion Dysregulation, 4*(1), 15. doi:10.1186/s40479-017-0065-5
- Harned, M. S., Korslund, K. E., & Linehan, M. M. (2014). A pilot randomized controlled trial of Dialectical Behavior Therapy with and without the Dialectical Behavior Therapy Prolonged Exposure protocol for suicidal and self-injuring women with borderline personality disorder and PTSD. *Behaviour Research and Therapy, 55*, 7-17. doi:10.1016/j.brat.2014.01.008
- Harned, M. S., Ruork, A. K., Liu, J., & Tkachuck, M. A. (2015). Emotional activation and habituation during Imaginal exposure for PTSD among women with borderline personality disorder. *Journal of Traumatic Stress, 28*(3), 253-257.
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (2011). *Acceptance and Commitment Therapy. The Process and Practice of Mindful Change* (2 ed.). New York: Guilford Press.
- Held, P., Owens, G. P., Monroe, J. R., & Chard, K. M. (2017). Increased Mindfulness Skills as Predictors of Reduced Trauma-Related Guilt in Treatment-Seeking Veterans. *Journal of Traumatic Stress, 30*(4), 425-431. doi:10.1002/jts.22209

- Jacobson, N. S., & Truax, P. (1991). Clinical Significance: A Statistical Approach to Defining Meaningful Change in Psychotherapy Research. *Journal of Consulting and Clinical Psychology, 59*(1), 12-19.
- Kendall, P. C., Marrs-Garcia, A., Nath, S. R., & Sheldrick, R. C. (1999). Normative comparisons for the evaluation of clinical significance. *Journal of Consulting and Clinical Psychology, 67*(3), 285-299.
- Kleim, B., Graham, B., Bryant, R. A., & Ehlers, A. (2013). Capturing Intrusive Re-experiencing in Trauma Survivors' Daily Lives Using Ecological Momentary Assessment. *Journal of Abnormal Psychology, 122*(4), 998-1009.
- Krause-Utz, A. & Elzinga, B. (2018). Current Understanding of the Neural Mechanisms of Dissociation in Borderline Personality Disorder. *Current Behavioral Neuroscience Reports, 5*(1), 113–123. doi: 10.1007/s40473-018-0146-9
- Kumpula, M. J., Pentel, K. Z., Foa, E. B., LeBlanc, N. J., Bui, E., McSweeney, L. B., . . . Rauch, S. A. M. (2017). Temporal Sequencing of Change in Posttraumatic Cognitions and PTSD Symptom Reduction During Prolonged Exposure Therapy. *Behavior Therapy, 48*(2), 156-165. doi:10.1016/j.beth.2016.02.008
- Langkaas, T. F., Hoffart, A., Øktedalen, T., Ulvenes, P. G., Hembree, E. A., & Smucker, M. (2017). Exposure and non-fear emotions: A randomized controlled study of exposure-based and rescripting-based imagery in PTSD treatment. *Behaviour Research and Therapy, 97*, 33-42. doi:10.1016/j.brat.2017.06.007
- Linehan, M. M. (1993). *Cognitive-behavioral therapy of borderline personality disorder*. New York: Guilford Press.
- Long, M. E., Davis, J. L., Springer, J. R., Elhai, J. D., Rhudy, J. L., Teng, E. J., & Frueh, B. C. (2011). The role of cognitions in imagery rescripting for posttraumatic nightmares. *Journal of Clinical Psychology, 67*(10), 1008-1016. doi:10.1002/jclp.20804

- Maercker, A., Hecker, T., Augsburger, M., & Kliem, S. (2018). ICD-11 Prevalence Rates of Posttraumatic Stress Disorder and Complex Posttraumatic Stress Disorder in a German Nationwide Sample. *The Journal of Nervous and Mental Disease, 206*(4), 270-276. doi:10.1097/nmd.0000000000000790
- Maikovich-Fong, A. K., & Jaffee, S. R. (2010). Sex differences in childhood sexual abuse characteristics and victims' emotional and behavioral problems: Findings from a national sample of youth. *Child Abuse & Neglect, 34*(6), 429-437.
- Mombour, W., Zaudig, M., Berger, P., Gutierrez, K., Berner, W., Berger, K., . . . von Bose, M. (1996). *IPDE, International Personality Disorder Examination von A.W. Loranger. Deutschsprachige Ausgabe*. Bern: Huber.
- Nasiakos, G., Cribbie, R. A., & Arpin-Cribbie, C. A. (2010). Equivalence-based measures of clinical significance: assessing treatments for depression. *Psychotherapy Research, 20*(6), 647-656. doi:0.1080/10503307.2010.501039
- Nishith, P., Nixon, R. D. V., & Resick, P. A. (2005). Resolution of trauma-related guilt following treatment of PTSD in female rape victims: A result of cognitive processing therapy targeting comorbid depression? *Journal of Affective Disorders, 86*(2-3), 259-265. doi:10.1016/j.jad.2005.02.013
- Orth, U., & Wieland, E. (2006). Anger, hostility, and posttraumatic stress disorder in trauma-exposed adults: a meta-analysis. *Journal of Consulting and Clinical Psychology, 74*(4), 698-706. doi:10.1037/0022-006x.74.4.698
- Pitman, R. K., Orr, S. P., Altman, B., & Longpre, R. E. (1996). Emotional processing and outcome of imaginal flooding therapy in Vietnam veterans with chronic post-traumatic stress disorder. *Comprehensive Psychiatry, 37*(6), 409-418.



- Pugh, L. R., Taylor, P. J., & Berry, K. (2015). The role of guilt in the development of post-traumatic stress disorder: A systematic review. *Journal of Affective Disorders, 182*, 138-150. doi:10.1016/j.jad.2015.04.026
- Rausch, S., Herzog, J., Thome, J., Ludäscher, P., Müller-Engelmann, M., Steil, R., . . . Kleindienst, N. (2016). Women with exposure to childhood interpersonal violence without psychiatric diagnoses show no signs of impairment in general functioning, quality of life and sexuality. *Borderline Personality Disorder and Emotion Dysregulation, 3*(1), 13. doi:10.1186/s40479-016-0048-y
- Rupp, C., Doebler, P., Ehring, T., & Vossbeck-Elsebusch, A. N. (2017). Emotional Processing Theory Put to Test: A Meta-Analysis on the Association Between Process and Outcome Measures in Exposure Therapy. *Clinical Psychology and Psychotherapy, 24*(3), 697-711. doi:10.1002/cpp.2039
- Saraiya, T., & Lopez-Castro, T. (2016). Ashamed and Afraid: A Scoping Review of the Role of Shame in Post-Traumatic Stress Disorder (PTSD). *Journal of Clinical Medicine, 5*(11), 94. doi:10.3390/jcm5110094
- Stapleton, J. A., Taylor, S., & Asmundson, G. J. (2006). Effects of three PTSD treatments on anger and guilt: exposure therapy, eye movement desensitization and reprocessing, and relaxation training. *Journal of Traumatic Stress, 19*(1), 19-28.
- Steil, R., Dyer, A., Priebe, K., Kleindienst, N., & Bohus, M. (2011). Dialectical Behavior Therapy for Posttraumatic Stress Disorder Related to Childhood Sexual Abuse: A Pilot Study of an Intensive Residential Treatment Program. *Journal of Traumatic Stress, 24*(1), 102-106. doi:10.1002/jts.20617
- Street, A. E., Gibson, L. E., & Holohan, D. R. (2005). Impact of childhood traumatic events, trauma-related guilt, and avoidant coping strategies on PTSD symptoms in female survivors of domestic violence. *Journal of Traumatic Stress, 18*(3), 245-252.

- Wade, D., Varker, T., Kartal, D., Hetrick, S., O'Donnell, M., & Forbes, D. (2016). Gender difference in outcomes following trauma-focused interventions for posttraumatic stress disorder: Systematic review and meta-analysis. *Psychological Trauma: Theory, Research, Practice, and Policy*, 8(3), 356-364. doi:10.1037/tra0000110
- Watts, B. V., Schnurr, P. P., Mayo, L., Young-Xu, Y., Weeks, W. B., & Friedman, M. J. (2013). Meta-analysis of the efficacy of treatments for posttraumatic stress disorder. *Journal of Clinical Psychiatry*, 74(6). doi:10.4088/JCP.12r08225
- WHO. (2018). *ICD-11. International Classification of Diseases 11th Revision*. Retrieved from <https://icd.who.int/>
- Wittchen, H., Wunderlich, U., Gruschwitz, S., & Zaudig, M. (1997). *Strukturiertes Klinisches Interview für DSM-IV. Achse 1: Psychische Störungen*. Göttingen: Hogrefe.
- You, S., Talbot, N. L., He, H., & Conner, K. R. (2012). Emotions and suicidal ideation among depressed women with childhood sexual abuse histories. *Suicide and Life-Threatening Behavior*, 42(3), 244-254. doi:10.1111/j.1943-278X.2012.00086.x

*Table 1. Patient Characteristics (N=42).*

	Mean	SD	Range
Age in years	42.17	9.64	23-60
Years in school	10.10	1.14	8-12
DTS at intake	88.39	23.24	29-128
Weeks in therapy	11.95	1.48	8-14
	n	% Total sample	
Sexual abuse	38	90.5%	
...before age 18yrs	33	78.6%	
...after age 18yrs	15	35.7	
Physical abuse	34	81.0	
...before age 18yrs	33	78.6	
...after age 18 yrs	7	16.7	
Accident	4	9.5	
Suicide/homicide	5	11.9	
War/Torture/Imprisonment	1	2.4	
Borderline Personality Disorder	15	35.7	
Major Depressive Disorder	31	73.8	
Anxiety Disorders	7	16.7	
Eating Disorders	9	21.4	
Antidepressants	36	85.7	
Neuroleptic medication	17	40.5	
Mood stabilizer	18	42.9	
Other psychotropic medication	16	38.1	



	Mean	59.29	11.19	$SMD = 1.53$				
	SD	31.19	16.11					
Shame	N	42	42	$t = 9.57^{**}$	$Wilk's \Lambda = 0.60^{**}$	69.0%	11.9%	69.0%
	Mean	75.71	25.48	$SMD = 1.48$				
	SD	30.30	26.98					
Anger	N	42	42	$t = 0.78$	$Wilk's \Lambda = 0.97$	35.7%	54.8%	59.5%
	Mean	43.33	39.29	$SMD = 0.12$				
	SD	35.11	32.11					
Sadness	N	42	42	$t = 2.33^*$	$Wilk's \Lambda = 1.00$	47.6%	28.6%	47.6%
	Mean	62.38	49.52	$SMD = 0.36$				
	SD	30.19	30.84					
Radical	N	41	42	$t = -5.66^{**}$	$Wilk's \Lambda = 0.86^*$	38.1 %	7.3%	31.0%
Acceptance	Mean	22.20	55.95	$SMD = 0.88$				
	SD	24.75	30.37					

<sup>1</sup>*SMDs* of 0.2, 0.5, and 0.8 correspond to small, medium, and large effects. Significant on the level of \* $<.05$  (two-tailed) \*\* $<.01$  (two-tailed)




Figure 1. Bertin plots for the intensity ratings of each negative trauma-related emotion at the start of the exposure therapy phase and at the end of DBT-PTSD. Each row refers to one person while each column represents one specific emotion. The color code ranges from light grey (low ratings) to black (high ratings) and white signifies missing observations.