







RESEARCH ARTICLE

Clinical spectrum of individuals with pathogenic *NF1* missense variants affecting p.Met1149, p.Arg1276, and p.Lys1423: genotype–phenotype study in neurofibromatosis type 1

Magdalena Koczkowska^{1*} | Tom Callens¹ | Yunjia Chen¹ | Alicia Gomes¹ | Alesha D. Hicks¹ | Angela Sharp¹ | Eric Johns¹ | Kim Armfield Uhas² | Linlea Armstrong³ | Katherine Armstrong Bosanko⁴ | Dusica Babovic-Vuksanovic⁵ | Laura Baker⁶ | Donald G. Basel⁷ | Mario Bengala⁸ | James T. Bennett⁹ | Chelsea Chambers¹⁰ | Lola K. Clarkson¹¹ | Maurizio Clementi¹² | Fanny M. Cortés¹³ | Mitch Cunningham¹⁴ | M. Daniela D'Agostino¹⁵ | Martin B. Delatycki¹⁶ | Maria C. Digilio¹⁷ | Laura Dosa¹⁸ | Silvia Esposito¹⁹ | Stephanie Fox¹⁵ | Mary-Louise Freckmann²⁰ | Christine Fauth²¹ | Teresa Giugliano²² | Sandra Giustini²³ | Allison Goetsch²⁴ | Yael Goldberg²⁵ | Robert S. Greenwood²⁶ | Cristin Griffis⁷ | Karen W. Gripp⁶ | Punita Gupta²⁷ | Eric Haan²⁸ | Rachel K. Hachen²⁹ | Tamara L. Haygarth³⁰ | Concepción Hernández-Chico³¹ | Katelyn Hodge³² | Robert J. Hopkin³³ | Louanne Hudgins³⁴ | Sandra Janssens³⁵ | Kory Keller³⁶ | Geraldine Kelly-Mancuso³³ | Aaina Kochhar³⁷ | Bruce R. Korf¹ | Andrea M. Lewis³⁸ | Jan Liebelt³⁹ | Angie Lichty¹¹ | Robert H. Listernick²⁴ | Michael J. Lyons¹¹ | Isabelle Maystadt⁴⁰ | Mayra Martinez Ojeda⁴¹ | Carey McDougall⁴² | Lesley K. McGregor³⁹ | Daniela Melis⁴³ | Nancy Mendelsohn⁴⁴ | Malgorzata J.M. Nowaczyk⁴⁵ | June Ortenberg¹⁵ | Karin Panzer⁴⁶ | John G. Pappas⁴⁷ | Mary Ella Pierpont⁴⁸ | Giulio Piluso²² | Valentina Pinna⁴⁹ | Eniko K. Pivnick⁵⁰ | Dinel A. Pond⁴⁴ | Cynthia M. Powell⁵¹ | Caleb Rogers³⁶ | Noa Ruhrman Shahar²⁵ | S. Lane Rutledge^{1†} | Veronica Saletti¹⁹ | Sarah A. Sandaradura⁵² | Claudia Santoro⁵³ | Ulrich A. Schatz²¹ | Allison Schreiber⁵⁴ | Daryl A. Scott³⁸ | Elizabeth A. Sellars⁴ | Ruth Sheffer⁵⁵ | Elizabeth Siqveland⁴⁴ | John M. Slopis⁵⁶ | Rosemarie Smith⁵⁷ | Alberto Spalice⁵⁸ | David W. Stockton¹⁴ | Haley Streff³⁸ | Amy Theos⁵⁹ |

*Magdalena Koczkowska, PhD, is also affiliated with the Department of Biology and Medical Genetics, Medical University of Gdansk, Gdansk, Poland.

†This paper is dedicated to the memory of our wonderful colleague, S. Lane Rutledge, MD, who passed away on January 2, 2019.

Gail E. Tomlinson⁶⁰ | Grace Tran⁶¹ | Pamela L. Trapane⁶² | Eva Trevisson¹²  |
 Nicole J. Ullrich⁶³ | Jenneke Van den Ende⁶⁴ | Samantha A. Schrier Vergano⁶⁵ |
 Stephanie E. Wallace⁹ | Michael F. Wangler³⁸  | David D. Weaver³² | Kaleb H. Yohay⁶⁶ |
 Elaine Zackai⁴² | Jonathan Zonana⁶⁷ | Vickie Zurcher⁵⁴ | Kathleen B. M. Claes³⁵  |
 Marica Eoli⁶⁸ | Yolanda Martin³¹ | Katharina Wimmer²¹  | Alessandro De Luca⁴⁹  |
 Eric Legius⁶⁹ | Ludwine M. Messiaen¹ 

¹Department of Genetics, University of Alabama at Birmingham, Birmingham, Alabama

²Children's Healthcare of Atlanta at Scottish Rite, Atlanta, Georgia

³Department of Medical Genetics, BC Women's Hospital, University of British Columbia, Vancouver, British Columbia, Canada

⁴Division of Clinical Genetics and Metabolism, Arkansas Children's Hospital, University of Arkansas for Medical Sciences, Little Rock, Arkansas

⁵Department of Clinical Genomics, Mayo Clinic, Rochester, Minnesota

⁶Division of Medical Genetics, Al DuPont Hospital for Children, Wilmington, Delaware

⁷Children's Hospital of Wisconsin, Milwaukee, Wisconsin

⁸U.O.C Laboratorio di Genetica Medica, Dipartimento di Oncematologia, Fondazione Policlinico di Tor Vergata, Rome, Italy

⁹Division of Genetic Medicine, Department of Pediatrics, University of Washington, Seattle, Washington

¹⁰Department of Neurology, University of Virginia Medical Center, Charlottesville, Virginia

¹¹Greenwood Genetic Center, Greenwood, South Carolina

¹²Clinical Genetics Unit, Department of Women's and Children's Health, University of Padova, Padova, Italy

¹³Center for Rare Diseases, Clinica Las Condes, Santiago, Chile

¹⁴Division of Genetic, Genomic, and Metabolic Disorders, Detroit Medical Center, Children's Hospital of Michigan, Detroit, Michigan

¹⁵Division of Medical Genetics, McGill University Health Centre, Montréal, Quebec, Canada

¹⁶Bruce Lefroy Centre for Genetic Health Research, Murdoch Childrens Research Institute, Parkville, Victoria, Australia

¹⁷Medical Genetics Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

¹⁸SOC Genetica Medica, AOU Meyer, Florence, Italy

¹⁹Developmental Neurology Unit, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy

²⁰Department of Clinical Genetics, Royal North Shore Hospital, St Leonards, New South Wales, Australia

²¹Division of Human Genetics, Medical University of Innsbruck, Innsbruck, Austria

²²Department of Precision Medicine, Università degli Studi della Campania "Luigi Vanvitelli", Naples, Italy

²³Department of Dermatology and Venereology, Policlinico Umberto I, Sapienza University of Rome, Rome, Italy

²⁴Department of Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, Illinois

²⁵The Raphael Recanati Genetics Institute, Rabin Medical Center, Petah Tikva, Israel

²⁶Division of Child Neurology, University of North Carolina School of Medicine, Chapel Hill, North Carolina

²⁷Neurofibromatosis Diagnostic and Treatment Program, St. Joseph's Children's Hospital, Paterson, New Jersey

²⁸Adult Genetics Unit, Royal Adelaide Hospital, Adelaide, South Australia, Australia

²⁹Neurofibromatosis Program, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania

³⁰Carolinas HealthCare System, Levine Children's Specialty Center, Charlotte, North Carolina

³¹Department of Genetics, Hospital Universitario Ramón y Cajal, Institute of Health Research (IRYCIS) and Center for Biomedical Research-Network of Rare Diseases (CIBERER), Madrid, Spain

³²Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, Indiana

³³Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

³⁴Division of Medical Genetics, Stanford University School of Medicine, Stanford, California

³⁵Center for Medical Genetics, Ghent University Hospital, Ghent, Belgium

³⁶Department of Molecular and Medical Genetics, Oregon Health and Science University, Portland, Oregon

³⁷Department of Medical Genetics and Metabolism, Valley Children's Healthcare, Madera, California

³⁸Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas

³⁹The South Australian Clinical Genetics Service at the Women's and Children's Hospital, North Adelaide, South Australia, Australia

⁴⁰Center for Human Genetics, Institute of Pathology and Genetics (IPG), Gosselies, Belgium

⁴¹Division of Genetics and Genomics, Boston Children's Hospital, Boston, Massachusetts

⁴²Division of Human Genetics, Children's Hospital of Philadelphia, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania

- ⁴³Section of Pediatrics, Department of Translational Medical Sciences, Federico II University, Naples, Italy
- ⁴⁴Genomics Medicine Program, Children's Hospital Minnesota, Minneapolis, Minnesota
- ⁴⁵Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario, Canada
- ⁴⁶University of Iowa Stead Family Children's Hospital, Iowa City, Iowa
- ⁴⁷Division of Clinical Genetic Services, Department of Pediatrics, NYU School of Medicine, New York, New York
- ⁴⁸Department of Pediatrics and Ophthalmology, University of Minnesota, Minneapolis, Minnesota
- ⁴⁹Molecular Genetics Unit, IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo, Foggia, Italy
- ⁵⁰Department of Pediatrics and Department of Ophthalmology, University of Tennessee Health Science Center, Memphis, Tennessee
- ⁵¹Department of Genetics and Department of Pediatrics, University of North Carolina School of Medicine, Chapel Hill, North Carolina
- ⁵²Division of Clinical Genetics, Department of Paediatrics and Child Health, Children's Hospital at Westmead, University of Sydney, Sydney, New South Wales, Australia
- ⁵³Specialistic and General Surgery Unit, Department of Woman and Child, Referral Centre of Neurofibromatosis, Università degli Studi della Campania "Luigi Vanvitelli", Naples, Italy
- ⁵⁴Cleveland Clinic, Genomic Medicine Institute, Cleveland, Ohio
- ⁵⁵Department of Genetics and Metabolic Diseases, Hadassah-Hebrew University Medical Center, Jerusalem, Israel
- ⁵⁶Department of Neuro-Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas
- ⁵⁷Division of Genetics, Department of Pediatrics, Maine Medical Center, Portland, Maine
- ⁵⁸Child Neurology Division, Department of Pediatrics, Sapienza University of Rome, Rome, Italy
- ⁵⁹Department of Dermatology, University of Alabama at Birmingham, Birmingham, Alabama
- ⁶⁰Division of Pediatric Hematology-Oncology, Greehey Children's Cancer Research Institute, The University of Texas Health Science Center, San Antonio, Texas
- ⁶¹Department of Clinical Cancer Genetics, The University of Texas MD Anderson Cancer Center, Houston, Texas
- ⁶²Division of Pediatric Genetics, Department of Pediatrics, University of Florida College of Medicine, Jacksonville, Florida
- ⁶³Department of Neurology, Boston Children's Hospital, Boston, Massachusetts
- ⁶⁴Center for Medical Genetics, University of Antwerp and Antwerp University Hospital, Antwerp, Belgium
- ⁶⁵Division of Medical Genetics and Metabolism, Children's Hospital of the King's Daughters, Norfolk, Virginia
- ⁶⁶Department of Neurology, New York University School of Medicine, Langone Medical Center, New York, New York
- ⁶⁷Department of Molecular and Medical Genetics, Oregon Health and Science University, Portland, Oregon
- ⁶⁸Division of Molecular Neuro-Oncology, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy
- ⁶⁹Department of Human Genetics, KU Leuven, Leuven, Belgium

Correspondence

Ludwine M. Messiaen, PhD, Medical Genomics Laboratory, Department of Genetics, University of Alabama, 720 20th St. S., Birmingham, AL 35294.
Email: lmessiaen@uabmc.edu

Funding information

Medical Genomics Laboratory at UAB, Grant/Award Number: internal funds; Children's Tumor Foundation by the Isaac and Sadie Fuchs Genotype-Phenotype Study; Italian Health Ministry, Grant/Award Number: RF-2016-0236293

Abstract

We report 281 individuals carrying a pathogenic recurrent *NF1* missense variant at p.Met1149, p.Arg1276, or p.Lys1423, representing three nontruncating *NF1* hotspots in the University of Alabama at Birmingham (UAB) cohort, together identified in 1.8% of unrelated *NF1* individuals. About 25% (95% confidence interval: 20.5–31.2%) of individuals heterozygous for a pathogenic *NF1* p.Met1149, p.Arg1276, or p.Lys1423 missense variant had a Noonan-like phenotype, which is significantly more compared with the “classic” *NF1*-affected cohorts (all $p < .0001$). Furthermore, p.Arg1276 and p.Lys1423 pathogenic missense variants were associated with a high prevalence of cardiovascular abnormalities, including pulmonic stenosis (all $p < .0001$), while p.Arg1276 variants had a high prevalence of symptomatic spinal neurofibromas ($p < .0001$) compared with “classic” *NF1*-affected cohorts. However, p.Met1149-positive individuals had a mild phenotype, characterized mainly by pigmentary manifestations without externally visible plexiform neurofibromas, symptomatic spinal neurofibromas or symptomatic optic pathway gliomas. As up to 0.4% of unrelated individuals in the UAB cohort carries a p.Met1149 missense variant, this finding will contribute to more accurate stratification of a significant number of *NF1* individuals. Although clinically relevant genotype-phenotype correlations are rare in *NF1*, each affecting only a small percentage of individuals, together they impact counseling and management of a significant number of the *NF1* population.

KEYWORDS

genotype–phenotype correlation, NF1, p.Arg1276, p.Lys1423, p.Met1149

1 | INTRODUCTION

Neurofibromatosis type 1 (NF1; MIM# 162200) is one of the most common autosomal dominant disorders with a birth incidence of 1 in 2,000–3,000 (Evans et al., 2010; Lammert, Friedman, Kluwe, & Mautner, 2005; Uusitalo et al., 2015). It is characterized by a highly variable expressivity and age-dependent clinical features, including multiple café-au-lait macules (CALMs), skinfold freckling, Lisch nodules, cutaneous, plexiform and/or spinal neurofibromas, optic pathway gliomas (OPGs), neoplasms, skeletal abnormalities, and learning difficulties. The wide range of variable expression makes genetic counseling and anticipatory guidance challenging (Friedman, 2014).

NF1 results from loss-of-function pathogenic variants in the *NF1* tumor suppressor gene (MIM# 613113), located on chromosome 17q11.2, a locus with one of the highest spontaneous mutation rates across single-gene human genetic disorders (Friedman, 2014; Huson, Compston, Clark, & Harper, 1989). Indeed, the extreme diversity of the *NF1* mutational spectrum observed in the University of Alabama at Birmingham (UAB) cohort of approximately 8,000 unrelated NF1-affected individuals, all comprehensively analyzed with *NF1* transcript analysis, includes greater than 3,000 different constitutional variants, with only 31 having a prevalence of $\geq 0.4\%$ (Messiaen & Wimmer, 2012; Messiaen, in preparation). The constitutional *NF1* microdeletion represents the most frequent recurrent pathogenic variant (~3–8%) and therefore was first recognized to have a genotype–phenotype association with a distinct severe form of NF1 (Kehrer-Sawatzki, Mautner, & Cooper, 2017). Nontruncating pathogenic variants affecting only a single amino acid are more likely to be associated with genotype–phenotype correlations, as they may represent gain-of-function or hypomorphic variants. Although missense variants contribute to approximately 17% of all variants found in unrelated NF1-affected individuals in the UAB cohort,

very few are recurrent and therefore amenable to variant-specific genotype–phenotype analyses. Indeed, only six true nontruncating hotspots, each with a prevalence $\geq 0.4\%$, were found in the UAB cohort, i.e., missense variants at codons 844–848, 1149, 1276, 1423, and 1809 and a single amino acid deletion p.Met992del (Figure 1). To date, genotype–phenotype associations affecting pathogenic missense variants at *NF1* codons 1809 and 844–848 and p.Met992del have been reported, representing approximately 1.2, 0.8, and 0.9% of unrelated probands, respectively (Koczkowska et al., 2018; Koczkowska et al., 2019; Pinna et al., 2015; Rojnueangnit et al., 2015; Upadhyaya et al., 2007). Increased efforts towards the identification of additional clinically relevant genotype–phenotype correlations are needed. Therefore, we examined with priority those individuals carrying a pathogenic variant at one of the three remaining nontruncating hotspots, i.e. at p.Met1149, p.Arg1276 and p.Lys1423.

In this cross-sectional study, we investigated the clinical spectrum of 281 individuals from 237 unrelated families identified through clinical genetic testing as being heterozygous for a missense variant at p.Met1149, p.Arg1276, or p.Lys1423.

2 | METHODS

In total, 237 unrelated probands and 44 relatives, all carrying a pathogenic *NF1* missense variant at either p.Met1149 (50 probands and 19 relatives), p.Arg1276 (101 probands and 18 relatives), or p.Lys1423 (86 probands and 7 relatives), were included in this study (Figure 2). Briefly, blood samples from 220 individuals (184 probands and 36 relatives) were originally sent to the Medical Genomics Laboratory at the UAB for molecular *NF1* genetic testing to establish or confirm the diagnosis of NF1. Two individuals from the UAB cohort, carrying c.3445A>T (p.Met1149Leu) and c.4268A>T (p.Lys1423Met) were excluded from the genotype–phenotype





FIGURE 1 Spectrum of six mutational hotspots at *NF1* codons 844–848 (67/8,000), 992 (74/8,000), 1149 (34/8,000), 1276 (57/8,000), 1423 (52/8,000), and 1809 (99/8,000), affecting a total of 383/8,000 (4.8%) of unrelated probands in the University of Alabama at Birmingham (UAB) cohort, associated with mild (upper panel) or severe (lower panel) phenotypes. The figure was prepared using ProteinPaint application (Zhou et al., 2016). CSRD, cysteine-serine rich domain; GAP, GTPase-activating protein; GRD, GAP related domain; *NF1*, neurofibromatosis type 1; PH, pleckstrin homology-like domain; Sec14, Sec14 homology-like domain; Syn, syndecan binding domain; TBD, tubulin-binding domain

FIGURE 2 Spectrum of pathogenic *NF1* missense variants affecting p.Met1149, p.Arg1276, and p.Lys1423 in the studied cohort of 237 unrelated probands (a) and 44 relatives (b). Each number in the circle corresponds with the total number of individuals heterozygous for a specific variant. The figure was prepared using the ProteinPaint application (Zhou et al., 2016). GAP, GTPase-activating protein; GRD, GAP related domain; *NF1*, neurofibromatosis type 1; TBD, tubulin-binding domain



studies as these specific variants are still interpreted as “variant of uncertain significance” (VUS) or “likely pathogenic”, respectively, but not (yet) “pathogenic”, according to the here applied American College of Medical Genetics and Genomics (ACMG) recommendations (Richards et al., 2015). This initial study was expanded to include an additional 63 individuals (55 probands and 8 relatives), molecularly diagnosed in collaborating institutions (please see details in the Supporting Information Methods). The collaborating institutions and their referring physicians completed the same phenotypic checklist and as they did not have prior insight into the phenotypic associations emerging from the UAB cohort, their contributions were independent and unbiased. All variants identified in this study with the confirmed origin of the variants were submitted to the LOVD and ClinVar databases.

Comprehensive *NF1* molecular analysis, interpretation of variant pathogenicity and collection of clinical data (see details in Supporting Information Methods) was performed as previously described (Koczkowska et al., 2018; Koczkowska et al., 2019; Messiaen et al., 2000; Richards et al., 2015; Rojnueangnit et al., 2015). Two-tailed Fisher’s exact test with $p < .05$ considered as statistically significant was applied. The resulting p values were adjusted for multiple comparisons using the Benjamini–Hochberg (B-H) procedure with false discovery rates (FDR) at 0.05 and 0.01 (Thissen, Steinberg, & Kuang, 2002). These statistical analyses were performed with GraphPad software. The risk ratio was calculated with the

STATCALC module from the Epi Info™ package version 7.2.3.1 (<http://cdc.gov/epiinfo/index.html>).

This study was approved by the UAB Institutional Review Board and all individuals participating in this study or their legal guardians signed the informed consent forms for clinical genetic testing.

3 | RESULTS

3.1 | Description of the *NF1* missense variants affecting p.Met1149, p.Arg1276, and p.Lys1423

The pathogenic missense variants described here were identified in approximately 0.4% (p.Met1149), 0.7% (p.Arg1276), and 0.7% (p.Lys1423) of probands, together affecting 1.8% (95% confidence interval: 1.5–2.1%) of unrelated *NF1* individuals in the UAB cohort.

Twelve different missense variants (representing 14 different substitutions) at codons 1149, 1276, or 1423 are reported here in 237 unrelated individuals, with only 8/12 present in disease-associated public databases (as of December 2018; Figure S1) and 2/12 reported in $\geq 245,936$ alleles in the control Genome Aggregation database (gnomAD; Tables S1–S3).

RNA-based sequencing indicated that three substitutions at *NF1* codon 1423, involving the last three nucleotides of exon 32 (24), were associated with in-frame exon 32 (24) skipping during *NF1*

messenger RNA splicing (Table S4) and therefore were classified as splicing variants, not true missense.

All variants described in this study were classified according to the ACMG recommendations (Richards et al., 2015), with only pathogenic missense variants included in the genotype–phenotype study (Table S5).

3.2 | Clinical characterization of the p.Met1149-positive cohort

Detailed clinical descriptions of 70 individuals from 51 different families heterozygous for one of four different pathogenic *NF1* amino substitutions at p.Met1149 (38 probands with c.3445A>G [p.Met1149Val], a single proband with c.3445A>T [p.Met1149Leu], six probands with c.3446T>C [p.Met1149Thr] and six probands with c.3447G>A/C/T [p.Met1149Ile]) are presented in Tables S6 and S7 and Figure S2. UAB-R1495, carrying c.3445A>T (p.Met1149Leu), was excluded from the genotype–phenotype study as the interpretation of this specific variant is “variant of uncertain significance” according to the ACMG recommendations (see details in Table S5).

p.Met1149-positive individuals ≥ 9 years presented with a mild phenotype, including multiple CALMs (41/46) and skinfold freckles (29/44). Lisch nodules were reported in 3/29 individuals ≥ 9 years. No externally visible plexiform or symptomatic spinal neurofibromas were seen in this cohort (0/42 and 0/41 ≥ 9 years, respectively), except for an 18-year-old girl (UAB-R2593) having one possible spinal lesion with differential diagnosis of spinal neurofibroma versus cyst. Four of 22 adults reportedly had 2–6 cutaneous and/or subcutaneous neurofibromas, but none had been biopsied and histopathologically confirmed (Table S6). No symptomatic or asymptomatic OPG was found in all 58, including 23 individuals who underwent magnetic resonance imaging (MRI) screening. Other benign neoplasms reported in this cohort (10.5%, 6/57) included lipomas ($n = 4$), pilomatixoma ($n = 1$), and a lesion in the right temporal lobe ($n = 1$). The prevalence of skeletal abnormalities, mainly pectus abnormalities ($n = 8$) was 24.6% (15/61). Short stature and macrocephaly were observed in 15.2% and 42.2% of cases, respectively. Thirty-one p.Met1149-positive individuals had cognitive impairment and/or learning disabilities (47%, 31/66). Although a high number of individuals with Noonan-like features (29%) was reported (Table S8), pulmonic stenosis (PS) was found only in two individuals (UAB-R9777 and UAB-R13811FN.405). Other cardiovascular abnormalities included atrial septal defect ($n = 2$), hypertrophic cardiomyopathy ($n = 1$), and quadricuspid aortic valve ($n = 1$). Finally, 4/24 (16.7%) cases ≥ 9 years with a completed phenotypic checklist did not fulfill the National Institute of Health (NIH) diagnostic criteria and 6/24 (25%) when family history was excluded as a criterion (Table S9).

3.3 | Clinical characterization of the p.Arg1276- and p.Lys1423-positive cohorts

Detailed clinical descriptions of 119 individuals from 101 different families heterozygous for one of five different pathogenic *NF1* amino

acid substitutions at p.Arg1276 (14 probands with c.3826C>G [p.Arg1276Gly], a single proband with c.3826_3827delinsGA [p.Arg1276Glu], 80 probands with c.3827G>A [p.Arg1276Gln], three probands with c.3827G>C [p.Arg1276Pro], and three probands with c.3827G>T [p.Arg1276Leu]) and 94 individuals from 87 different families heterozygous for one of four different pathogenic *NF1* amino substitutions at p.Lys1423 (seven probands with c.4267A>C [p.Lys1423Gln], 77 probands with c.4267A>G [p.Lys1423Glu], two probands with c.4268A>C [p.Lys1423Thr], and a single proband with c.4268A>T [p.Lys1423Met]) are presented in Tables S10–S13 and Figures S3 and S4. UAB-R1753, carrying c.4268A>T (p.Lys1423Met), was excluded from the genotype–phenotype study as the interpretation of this specific variant is “likely pathogenic” according to the ACMG recommendations (see details in Table S5).

Individuals carrying p.Arg1276 or p.Lys1423 pathogenic missense variants presented with more severe phenotypes compared with the p.Met1149-positive cohort (Tables S11–S13). The most striking feature associated with p.Arg1276 was the presence of symptomatic spinal neurofibromas (18/97, 18.6% all ages). As many as 17/36 (47.2%) adults (≥ 19 years) had symptomatic spinal tumors and an additional 4/11 (36.4%) had multiple tumors on spinal roots without clinical symptoms (Table S14). A very high load of symptomatic or asymptomatic spinal neurofibromas, but few cutaneous neurofibromas and/or pigmentary manifestations, characteristic for so-called spinal NF, was seen in 5/21 adults (Figure S5), including one proband with a family history of spinal NF (UAB-R743). The p.Lys1423 cohort was characterized by a high prevalence of externally visible plexiform neurofibromas compared with the p.Arg1276 cohort (15/48 vs. 5/64 in ≥ 9 years; $p = .0022$; Tables S16–S18), but a lower prevalence of symptomatic spinal tumors (3/65 vs. 18/97 all ages; $p = .0091$; Tables S14, S15, and S18). Furthermore, 82.1% of adults with p.Lys1423 (23/28), but only 35% with p.Arg1276 (14/40) had ≥ 2 cutaneous neurofibromas ($p = .0002$; Table S18).

The prevalence of CALMs and skinfold freckling was similar in both cohorts (Tables S11 and S13). Lisch nodules were more frequently reported in the p.Lys1423-positive individuals (52.5%, 31/59) compared with the p.Arg1276-positive cases (24.1%, 19/70). Despite the severe phenotypes observed in p.Arg1276-positive cases, 3/31 individuals ≥ 9 years with a complete phenotypic checklist did not fulfill the NIH diagnostic criteria after excluding family history (Table S19).

Symptomatic OPGs were not found in the p.Arg1276-positive individuals (0/97) and were rare in the p.Lys1423 cohort (1/74; EUR-R49). An asymptomatic OPG was identified by MRI screening in 1/48 (2.1%) p.Arg1276-positive and 6/40 (15%) p.Lys1423-positive individuals. Neoplasms, other than OPGs and neurofibromas, were observed in 8/94 (8.5%) of the p.Arg1276-positive and 11/77 (14.3%) p.Lys1423-positive individuals, including 4/8 and 7/11, respectively who developed malignant lesions (Tables S10–S12).

Similarly to the p.Met1149 cohort, Noonan-like features were frequently reported in the p.Arg1276- and p.Lys1423-positive individuals (22/106 and 24/83, respectively). The prevalence may even be higher, as some individuals were noted as having possible

Noonan-like features (Tables S20 and S21), especially as Noonan-like features are more difficult to discern in adulthood (Allanson, Hall, Hughes, Preus, & Witt, 1985). Importantly, p.Arg1276 and p.Lys1423 cohorts had a high prevalence of cardiac/cardiovascular abnormalities (23.9%, 22/92 and 25%, 19/76, respectively; Tables S22 and S23), including PS (12%, 11/92 and 14.5%, 11/76, respectively). Short stature and macrocephaly were present in 17.5% (14/80) and 31.6% (24/76) of p.Arg1276-positive cases and in 41.2% (21/51) and 29.4% (15/51) of p.Lys1423-positive cases, respectively. The prevalence of cognitive impairment and/or learning disabilities was estimated at 43.8% (46/105) and 41.4% (36/87) in p.Arg1276 and p.Lys1423 cohorts, respectively.

3.4 | Comparison of clinical features of the studied cohorts with “classic” NF1 population, individuals carrying nonsense variants and cohorts of individuals with previously reported NF1 genotype–phenotype correlations

Detailed results of comparisons between phenotypes in the studied groups and the *NF1* codons 844-848-, 1809-, and p.Met992del-positive cohorts, “classic” NF1-affected cohorts, and the cohort of individuals with one of the 18 most recurrent nonsense variants identified in the UAB database are presented in Tables 1–3, Table S18, and Tables S24–S27.

All missense variants studied in the current research were associated with a high prevalence of Noonan-like phenotypes compared with the general NF1 population and/or the cohort carrying an *NF1* nonsense variant (all $p < .0001$, significant at FDR of 0.01 after B-H correction). Moreover, p.Arg1276- and p.Lys1423-positive individuals had a very significantly increased prevalence of cardiac/cardiovascular abnormalities, including PS, compared with the “classic” NF1 and nonsense variant cohorts (all $p < .0001$, significant at FDR of 0.01 after B-H correction).

Phenotype in p.Met1149-positive cases were similar to that of individuals with the p.Met992del and p.Arg1809 pathogenic variants, including paucity of superficial plexiform and histopathologically confirmed cutaneous and/or subcutaneous neurofibromas. No symptomatic spinal neurofibromas, symptomatic OPGs, or malignant neoplasms were found in this cohort (Table 1).

Compared with the general NF1 population, p.Arg1276-positive individuals showed a high prevalence of symptomatic spinal neurofibromas ($p < .0001$, significant at FDR of 0.01 after B-H correction) and a lower prevalence of cutaneous neurofibromas ($p < .0001$, significant at FDR of 0.01 after B-H correction).

Pathogenic *NF1* missense variants at p.Lys1423 may predispose to major external plexiform neurofibromas compared with “classic” NF1-affected cohorts (29.2% vs. 18.5%; $p = .0863$, but not significant).

Both p.Arg1276 and p.Lys1423 cohorts had a higher incidence of skeletal abnormalities (Table S28) compared with the general NF1 population (both $p \leq .0001$, significant at FDR of 0.01 after B-H correction).

There were no statistical differences in the prevalence of NF1 clinical features between cohorts of individuals heterozygous for p.Met1149, p.Arg1276, or p.Lys1423 referred to UAB and to the collaborating European institutions (Tables S29 and S30).

4 | DISCUSSION

NF1 and Noonan syndrome (NS; MIM# 163950) represent RASopathies, a group of phenotypically related conditions, caused by pathogenic variants in genes involved in the RAS/mitogen-activated protein kinase signaling transduction pathway. Due to some overlapping phenotypic features, a clinical diagnosis can be complicated. “Classic” NF1 cases presenting with Noonan-like features were classified as having neurofibromatosis-Noonan syndrome (NFNS; MIM# 601321), first described 30 years ago (Allanson, Hall, & Van Allen, 1985; Opitz & Weaver, 1985). Although few cases have been reported to carry both a pathogenic *NF1* and *PTPN11* (MIM# 176876) variant (Bertola et al., 2005; Nyström et al., 2009; Thiel et al., 2009), as a rule, *NF1* is the genetic cause underlying NFNS. De Luca et al. (2005) identified pathogenic variants spread across the *NF1* gene, including nonsense, missense, out-of-frame and small in-frame deletions, and one multiexon deletion in 16/17 unrelated NFNS individuals. Among four individuals with NFNS and PS, 2/4 carried an in-frame single amino acid deletion and 1/4 a pathogenic missense variant with 2/3 located in the GTPase-activating protein-related domain (GAP-GRD). Ben-Shachar et al. (2013) reported a higher prevalence of PS in NFNS individuals compared with the general NF1 population (9/35, 26% vs. 25/2322, 1.1%; $p < .001$), with NFNS cases carrying a nontruncating pathogenic variant having the highest risk ($p < .001$). However, this study did not provide more specific information related to location or type of nontruncating pathogenic variants and was not adjusted for multiple comparisons.

So far, only two specific *NF1* pathogenic variants have been associated with a statistically significant increased prevalence of Noonan-like features compared with “classic” NF1-affected cohorts, i.e., p.Arg1809 and p.Met992del, both located outside the GRD (Koczkowska et al., 2019; Pinna et al., 2015; Rojnueangnit et al., 2015; Upadhyaya et al., 2007). Here, we report that *NF1* pathogenic missense variants at p.Met1149, p.Arg1276, and p.Lys1423 also are associated with this distinct phenotype (all $p < .0001$, significant at FDR of 0.01 after B-H correction), with no pathogenic variants in other NS genes detected (Tables S6, S8, S10, S12, S20, and S21). A high prevalence of PS was associated with p.Arg1276, p.Lys1423 and p.Arg1809 compared to the general NF1 population as well as the UAB *NF1* nonsense variant cohort (all $P < 0.0001$, significant at FDR of 0.01 after B-H correction, Tables 2,3 and Table S18). These observations provide evidence that although PS overall is more frequently observed in individuals carrying one of the nontruncating pathogenic variants in the current and previous studies (48/625, 7.7%), the prevalence of PS and other cardiac/cardiovascular abnormalities is highest in the p.Arg1276- and p.Lys1423- positive individuals, both located in the GRD (at least 10-fold higher than in

TABLE 1 Comparison of clinical features of the cohort of individuals heterozygous for pathogenic NF1 missense variants affecting p.Met1149 with the cohorts of individuals with pathogenic NF1 missense variants affecting codons 1809 and 844-848, the NF1 p.Met992del as well as with large-scale previously reported cohorts of individuals with "classic" NF1

NF1 feature	N (%)		p Value (two-tailed Fisher's exact test)						
	p.Met1149	p.Met992del ^a	p.Arg1809 ^b	aa 844-848 ^c	Previously reported NF1 cohorts ^d	p.Met1149 versus p.Met992del	p.Met1149 versus p.Arg1809	p.Met1149 versus aa 844-848	p.Met1149 versus "classic" NF1
>5 CALMs	62/69 (89.9)	165/182 (90.7)	157/169 (92.9)	130/157 (82.8)	1,537/1,728 (89)				
Skinfold freckling	40/65 (61.5)	105/171 (61.4)	95/161 (59)	104/144 (72.2)	1,403/1,667 (84.2)				<0.0001** ↘
Lisch nodules	3/44 (6.8)	16/139 (11.5)	12/120 (10)	42/98 (42.9)	729/1,237 (58.9)			<0.0001** ↘	<0.0001** ↘
Major external plexiform neurofibromas ^e	0/42 (0)	0/125 (0)	0/105 (0)	36/92 (39.1)	120/648 (18.5)			<0.0001** ↘	0.0005 ** ↘
Cutaneous neurofibromas ^f	0-3/24 (0-12.5) ^g	0-1/57 (0-1.8) ^g	0/57 (0)	47/69 (68.1)	656/723 (90.7)			<0.0001** ↘	<0.0001** ↘
Subcutaneous neurofibromas ^f	0-3/22 (0-13.6) ^g	0-3/36 (0-8.3) ^g	0-5/57 (0-8.8) ^g	33/65 (50.8)	297/515 (57.7)			<0.0001** ↘	<0.0001** ↘
Symptomatic spinal neurofibromas	0/59 (0)	1/165 (0.6)	0/76 (0)	13/127 (10.2)	36/2,058 (1.8)				
Symptomatic OPGs ^h	0/58 (0)	0/170 (0)	0/139 (0)	12/136 (8.8)	64/1,650 (3.9)				
Asymptomatic OPGs ⁱ	0/23 (0)	1/41 (2.4)	0/38 (0)	18/63 (28.6)	70/519 (13.5)			0.0023* ↘	
Other malignant neoplasms ^j	0/57 (0)	1/126 (0.8) ^k	2/155 (1.3) ^l	13/139 (9.4)	18/523 (3.4)				
Skeletal abnormalities	15/61 (24.6)	30/172 (17.4)	21/126 (16.7)	48/144 (33.3)	144/948 (15.2)				
Scoliosis ^f	2/20 (10)	7/57 (12.3)	6/48 (12.5)	20/64 (31.3)	51/236 (21.6)				
Cognitive impairment and/or learning disabilities	31/66 (47)	58/176 (33)	80/159 (50.3)	56/138 (40.6)	190/424 (44.8)				
Noonan-like phenotype ^m	18/62 (29)	19/166 (11.5)	46/148 (31.1)	10/134 (7.5)	57/1,683 (3.4)		0.0023* ↗	0.0001** ↗	<0.0001** ↗

(Continues)

TABLE 1 (Continued)

NF1 feature	N (%)		p Value (two-tailed Fisher's exact test)					
	p.Met1149	p.Met992del ^a	p.Arg1809 ^b	aa 844-848 ^c	Previously reported NF1 cohorts ^d	p.Met1149 versus p.Arg1809	p.Met1149 versus aa 844-848	p.Met1149 versus "classic" NF1
Short stature ^e	5/33 (15.2)	16/118 (13.6)	32/111 (28.8)	15/91 (16.5)	109/684 (15.9)			
Macrocephaly	19/45 (42.2)	30/132 (22.7)	31/107 (29)	36/98 (36.7)	239/704 (33.9)			
Pulmonic stenosis	2/52 (3.9)	8/160 (5)	14/132 (10.6)	2/113 (1.8)	25/2,322 (1.1)			
Cardiovascular abnormalities	5/52 (9.6)	16/160 (10)	21/118 (17.8)	16/113 (14.2)	54/2,322 (2.3)			

Note: Statistically significant *p* values with FDR of 0.05 (indicated by*) and 0.01 (indicated by**) after correction for multiple testing using Benjamini–Hochberg procedure (see details in Table S24). After applying the Benjamini–Hochberg correction, $p \leq .0023$ and $p \leq .0005$ remained statistically significant at FDR of 0.05 and 0.01, respectively. The black arrows indicate the statistically significant differences of the NF1 clinical features prevalence between the p.Met1149 group and the cohort(s) used for the comparison, with the up and down arrows representing an increase and a decrease of the prevalence in the p.Met1149 group, respectively.

Abbreviations: CALM, café-au-lait macule; FDR, false discovery rate; MRI, magnetic resonance imaging; OPG, optic pathway glioma.

^aBased on data from Upadhyaya et al. (2007) and Koczkowska et al. (2019).

^bBased on data from Pinna et al. (2015), Rojueangnit et al. (2015), Ekvall et al. (2014), Nyström et al. (2009) and Santoro et al. (2015).

^cBased on data from Koczkowska et al. (2018).

^dPrevious NF1 cohorts used for the comparison: Huson, Harper, and Compston (1988), Huson, Compston, and Harper (1989), Listernick, Charrow, Greenwald, and Mets (1994), Friedman and Birch (1997), Clossen et al. (1998), McGaughan et al. (1999), Thakkar, Feigen, and Mautner (1999), Lin et al. (2000), Blazo et al. (2004), Khosrotehrani et al. (2005), Plotkin et al. (2012), and/or Blanchard et al. (2016).

^eIn individuals ≥ 9 years old.

^fIn individuals ≥ 19 years old.

^gIndividuals with few (2–6) cutaneous and/or subcutaneous "neurofibromas," none were biopsied and therefore none have been histologically confirmed.

^hThe absence of symptomatic OPGs was determined by ophthalmological examination and/or by MRI.

ⁱIncluding only individuals without signs of symptomatic OPGs who underwent MRI examination.

^jOnly malignant neoplasms, not including OPGs and neurofibromas, have been taken into account.

^kA single case of neuroblastoma ($n = 1$) was found in the NF1 p.Met992del cohort, no follow-up information on this individual was available.

^lBreast cancer ($n = 1$) and Ewing sarcoma ($n = 1$) were found in the NF1 p.Arg1809 cohort, no follow-up information on these individuals was available.

^mAn individual was classified as having a Noonan-like phenotype when at least two of the following features were present: short stature, low set ears, hypertelorism, midface hypoplasia, webbed neck, pectus abnormality, and/or pulmonic stenosis.

ⁿAs no specific growth curves are available for the Hispanic and Asian populations, Hispanic and Asian individuals were excluded as having short or normal stature.

TABLE 2 Comparison of clinical features of the cohort of individuals heterozygous for pathogenic NF1 missense variants affecting p.Arg1276 with the cohorts of individuals with pathogenic NF1 missense variants affecting codons 1809 and 844-848, the NF1 p.Met992del as well as with large-scale previously reported cohorts of individuals with "classic" NF1

NF1 feature	N (%)		p Value (two-tailed Fisher's exact test)						
	p.Arg1276	p.Met992del ^a	p.Arg1809 ^b	aa 844-848 ^c	Previously reported NF1 cohorts ^d	p.Arg1276 versus p.Met992del	p.Arg1276 versus p.Arg1809	p.Arg1276 versus aa 844-848	p.Arg1276 versus "classic" NF1
>5 CALMs	111/119 (93.3)	165/182 (90.7)	157/169 (92.9)	130/157 (82.8)	1,537/1,728 (89)				
Skinfold freckling	74/112 (66.1)	105/171 (61.4)	95/161 (59)	104/144 (72.2)	1,403/1,667 (84.2)				<0.0001** ↘
Lisch nodules	19/70 (24.1)	16/139 (11.5)	12/120 (10)	42/98 (42.9)	729/1,237 (58.9)	0.0059* ↗	0.0038* ↗		<0.0001** ↘
Major external plexiform neurofibromas ^e	5/64 (7.8)	0/125 (0)	0/105 (0)	36/92 (39.1)	120/648 (18.5)	0.0040* ↗	0.0070* ↗	<0.0001** ↘	
Cutaneous neurofibromas ^f	14/40 (35)	0-1/57 (0-1.8) ^g	0/57 (0)	47/69 (68.1)	656/723 (90.7)	<0.0001** ↗	<0.0001** ↗	0.0012* ↘	<0.0001** ↘
Subcutaneous neurofibromas ^f	21/37 (56.8)	0-3/36 (0-8.3) ^g	0-5/57 (0-8.8) ^g	33/65 (50.8)	297/515 (57.7)	<0.0001** ↗	<0.0001** ↗		
Symptomatic spinal neurofibromas	18/97 (18.6) ^h	1/165 (0.6)	0/76 (0)	13/127 (10.2)	36/2,058 (1.8)	<0.0001** ↗	<0.0001** ↗		<0.0001** ↗
Symptomatic OPGs ⁱ	0/97 (0)	0/170 (0)	0/139 (0)	12/136 (8.8)	64/1,650 (3.9)			0.0016* ↘	
Asymptomatic OPGs ^j	1/48 (2.1)	1/41 (2.4)	0/38 (0)	18/63 (28.6)	70/519 (13.5)			0.0002** ↘	
Other malignant neoplasms ^k	4/94 (4.3) ^l	1/126 (0.8) ^m	2/155 (1.3) ⁿ	13/139 (9.4)	18/523 (3.4)				
Skeletal abnormalities	32/100 (32)	30/172 (17.4)	21/126 (16.7)	48/144 (33.3)	144/948 (15.2)	0.0070* ↗	0.0076* ↗		0.0001** ↗
Scoliosis ^f	8/35 (22.9)	7/57 (12.3)	6/48 (12.5)	20/64 (31.3)	51/236 (21.6)				
Cognitive impairment and/or learning disabilities	46/105 (43.8)	58/176 (33)	80/159 (50.3)	56/138 (40.6)	190/424 (44.8)				
Noonan-like phenotype ^o	22/106 (20.8)	19/166 (11.5)	46/148 (31.1)	10/134 (7.5)	57/1,683 (3.4)			0.0037* ↗	<0.0001** ↗

(Continues)

TABLE 2 (Continued)

NF1 feature	N (%)		p Value (two-tailed Fisher's exact test)				
	p.Arg1276	p.Arg1809 ^b	aa 844-848 ^c	Previously reported NF1 cohorts ^d	p.Arg1276 versus p.Arg1809	p.Arg1276 versus aa 844-848	p.Arg1276 versus "classic" NF1
Short stature ^e	14/80 (17.5)	16/118 (13.6)	32/111 (28.8)	109/684 (15.9)			
Macrocephaly	24/76 (31.6)	30/132 (22.7)	31/107 (29)	239/704 (33.9)			
Pulmonic stenosis	11/92 (12)	8/160 (5)	14/132 (10.6)	25/2,322 (1.1)		0.0034* ↗	<0.0001** ↗
Cardiovascular abnormalities	22/92 (23.9)	16/160 (10)	21/118 (17.8)	54/2,322 (2.3)	0.0055** ↗		<0.0001** ↗

Note: Statistically significant *p* values with FDR of 0.05 (indicated by *) and 0.01 (indicated by **) after correction for multiple testing using Benjamini–Hochberg procedure (see details in Table S25). After applying the Benjamini–Hochberg correction, *p* ≤ .0076 and *p* ≤ .0002 remained statistically significant at FDR of 0.05 and 0.01, respectively. The black arrows indicate the statistically significant differences of the NF1 clinical features prevalence between the p.Arg1276 group and the cohort(s) used for the comparison, with the up and down arrows representing an increase and a decrease of the prevalence in the p.Arg1276 group, respectively.

Abbreviations: CALM, café-au-lait macule; FDR, false discovery rate; MPNST, malignant peripheral nerve sheath tumor; MRI, magnetic resonance imaging; OPG, optic pathway glioma.

^aBased on data from Upadhyaya et al. (2007) and Koczkowska et al. (2019).

^bBased on data from Pinna et al. (2015), Rojueangnit et al. (2015), Ekvall et al. (2014), Nyström et al. (2009), and Santoro et al. (2015).

^cBased on data from Koczkowska et al. (2018).

^dPrevious NF1 cohorts used for the comparison: Huson et al. (1988), Huson, Compston, Clark et al. (1989), Huson, Compston, and Harper (1989), Listernick et al. (1994), Friedman and Birch (1997), Cnossen et al. (1998), McGaughan et al. (1999), Thakkar et al. (2000), Blazo et al. (2004), Khosrotehrani et al. (2005), Plotkin et al. (2012), and/or Blanchard et al. (2016).

^eIn individuals ≥9 years old.

^fIn individuals ≥19 years old.

^gIndividuals with few (2–6) cutaneous and/or subcutaneous "neurofibromas," none were biopsied and therefore none have been histologically confirmed.

^hThe overall prevalence of symptomatic spinal neurofibromas in all individuals was 18.6% (18/97) but in adults 47.2% (17/36 ≥ 19 years old).

ⁱThe absence of symptomatic OPGs was determined by ophthalmological examination and/or by MRI.

^jIncluding only individuals without signs of symptomatic OPGs who underwent MRI examination.

^kOnly malignant neoplasms, not including OPGs and neurofibromas, have been taken into account.

^lAstrocytoma (*n* = 2), colon cancer (*n* = 1) and MPNST (*n* = 1) were found in the NF1 p.Arg1276 cohort.

^mA single case of neuroblastoma (*n* = 1) was found in the NF1 p.Arg1276 cohort; no follow-up information on this individual was available.

ⁿBreast cancer (*n* = 1) and Ewing sarcoma (*n* = 1) were found in the NF1 p.Arg1809 cohort, no follow-up information on these individuals was available.

^oAn individual was classified as having a Noonan-like phenotype when at least two of the following features were present: short stature, low set ears, hypertelorism, midface hypoplasia, webbed neck, pectus abnormality, and/or pulmonic stenosis.

^pAs no specific growth curves are available for the Hispanic and Asian populations, Hispanic and Asian individuals were excluded as having short or normal stature.

TABLE 3 Comparison of clinical features of the cohort of individuals heterozygous for pathogenic NF1 missense variants affecting p.Lys1423 with the cohorts of individuals with pathogenic NF1 missense variants affecting codons 1809 and 844-848, the NF1 p.Met992del as well as with large-scale previously reported cohorts of individuals with "classic" NF1

NF1 feature	N (%)		p Value (two-tailed Fisher's exact test)						
	p.Lys1423	p.Met992del ^a	p.Arg1809 ^b	aa 844-848 ^c	Previously reported NF1 cohorts ^d	p.Lys1423 versus p.Met992del	p.Lys1423 versus p.Arg1809	p.Lys1423 versus aa 844-848	p.Lys1423 versus "classic" NF1
>5 CALMs	86/91 (94.5)	165/182 (90.7)	157/169 (92.9)	130/157 (82.8)	1,537/1,728 (89)				
Skinfold freckling	65/85 (76.5)	105/171 (61.4)	95/161 (59)	104/144 (72.2)	1,403/1,667 (84.2)		0.0074* ↗		
Lisch nodules	31/59 (52.5)	16/139 (11.5)	12/120 (10)	42/98 (42.9)	729/1,237 (58.9)	<0.0001** ↗	<0.0001** ↗		
Major external plexiform neurofibromas ^e	14/48 (29.2)	0/125 (0)	0/105 (0)	36/92 (39.1)	120/648 (18.5)	<0.0001** ↗	<0.0001** ↗		
Cutaneous neurofibromas ^f	23/28 (82.1)	0-1/57 (0-1.8) ^g	0/57 (0)	47/69 (68.1)	656/723 (90.7)	<0.0001** ↗	<0.0001** ↗		
Subcutaneous neurofibromas ^f	13/23 (56.5)	0-3/36 (0-8.3) ^g	0-5/57 (0-8.8) ^g	33/65 (50.8)	297/515 (57.7)	<0.0001** ↗	<0.0001** ↗		
Symptomatic spinal neurofibromas	3/65 (4.6)	1/165 (0.6)	0/76 (0)	13/127 (10.2)	36/2,058 (1.8)				
Symptomatic OPGs ^h	1/74 (1.4)	0/170 (0)	0/139 (0)	12/136 (8.8)	64/1,650 (3.9)				
Asymptomatic OPGs ⁱ	6/40 (15)	1/41 (2.4)	0/38 (0)	18/63 (28.6)	70/519 (13.5)				
Other malignant neoplasms ^j	7/77 (9.1) ^k	1/126 (0.8) ^l	2/155 (1.3) ^m	13/139 (9.4)	18/523 (3.4)	0.0052* ↗	0.0070* ↗		
Skeletal abnormalities	34/83 (41)	30/172 (17.4)	2/126 (1.6) ⁿ	48/144 (33.3)	144/948 (15.2)	<0.0001** ↗	0.0002** ↗		<0.0001** ↗
Scoliosis ^f	10/27 (37)	7/57 (12.3)	6/48 (12.5)	20/64 (31.3)	51/236 (21.6)				
Cognitive impairment and/or learning disabilities	36/87 (41.4)	58/176 (33)	80/159 (50.3)	56/138 (40.6)	190/424 (44.8)				
Noonan-like phenotype ⁿ	24/83 (28.9)	19/166 (11.5)	46/148 (31.1)	10/134 (7.5)	57/1,683 (3.4)	0.0011** ↗	<0.0001** ↗		<0.0001** ↗

(Continues)

TABLE 3 (Continued)

NF1 feature	N (%)		p Value (two-tailed Fisher's exact test)					
	p.Lys1423	p.Met992del ^a	p.Arg1809 ^b	aa 844-848 ^c	Previously reported NF1 cohorts ^d	p.Lys1423 versus p.Arg1809	p.Lys1423 versus aa 844-848	p.Lys1423 versus "classic" NF1
Short stature ^e	21/51 (41.2)	16/118 (13.6)	32/111 (28.8)	15/91 (16.5)	109/684 (15.9)	0.0002** ↗	0.0022* ↗	<0.0001** ↗
Macrocephaly	15/51 (29.4)	30/132 (22.7)	31/107 (29)	36/98 (36.7)	239/704 (33.9)			
Pulmonic stenosis	11/76 (14.5)	8/160 (5)	14/132 (10.6)	2/113 (1.8)	25/2,322 (1.1)		0.0010** ↗	<0.0001** ↗
Cardiovascular abnormalities	19/76 (25)	16/160 (10)	21/118 (17.8)	16/113 (14.2)	54/2,322 (2.3)	0.0053** ↗		<0.0001** ↗

Note: Statistically significant *p* values with FDR of 0.05 (indicated by *) and 0.01 (indicated by **) after correction for multiple testing using Benjamini–Hochberg procedure (see details in Table S26). After applying the Benjamini–Hochberg correction, $p \leq .0074$ and $p \leq .0011$ remained statistically significant at FDR of 0.05 and 0.01, respectively. The black arrows indicate the statistically significant differences of the NF1 clinical features prevalence between the p.Lys1423 group and the cohort(s) used for the comparison, with the up and down arrows representing an increase and a decrease of the prevalence in the p.Lys1423 group, respectively.

Abbreviations: CALM, café-au-lait macule; FDR, false discovery rate; MRI, magnetic resonance imaging; OPG, optic pathway glioma.

^aBased on data from Upadhyaya et al. (2007) and Koczkowska et al. (2019).

^bBased on data from Pinna et al. (2015), Rojueangnit et al. (2015), Ekvall et al. (2014), Nyström et al. (2009), and Santoro et al. (2015).

^cBased on data from Koczkowska et al. (2018).

^dPrevious NF1 cohorts used for the comparison: Huson et al. (1988), Huson, Compston, Clark et al. (1989), Huson, Compston, and Harper (1989), Listernick et al. (1994), Friedman and Birch (1997), Cnossen et al. (1998), McGaughan et al. (1999), Thakkar et al. (1999), Lin et al. (2000), Blazo et al. (2004), Khosrotehrani et al. (2005), Plotkin et al. (2012), and/or Blanchard et al. (2016).

^eIn individuals ≥ 9 years old.

^fIn individuals ≥ 19 years old.

^gIndividuals with few (2–6) cutaneous and/or subcutaneous "neurofibromas," none were biopsied and therefore none have been histologically confirmed.

^hThe absence of symptomatic OPGs was determined by ophthalmological examination and/or by MRI.

ⁱIncluding only individuals without signs of symptomatic OPGs who underwent MRI examination.

^jOnly malignant neoplasms, not including OPGs and neurofibromas, have been taken into account.

^kAstrocytoma ($n = 1$), rhabdomyosarcoma ($n = 1$), hypothalamic glioma ($n = 2$), brainstem glioma ($n = 1$), colon adenocarcinoma ($n = 1$), and cerebellar glioblastoma ($n = 1$) were found in the NF1 p.Lys1423 cohort.

^lA single case of neuroblastoma ($n = 1$) was found in the NF1 p.Met992del cohort; no follow-up information on this individual was available.

^mBreast cancer ($n = 1$) and Ewing sarcoma ($n = 1$) were found in the NF1 p.Arg1809 cohort; no follow-up information on these individuals was available.

ⁿAn individual was classified as having a Noonan-like phenotype when at least two of the following features were present: short stature, low set ears, hypertelorism, midface hypoplasia, webbed neck, pectus abnormality, and/or pulmonic stenosis.

^oAs no specific growth curves are available for the Hispanic and Asian populations, Hispanic and Asian individuals were excluded as having short or normal stature.

TABLE 4 Risk ratio calculations with 95% CI for the comparison of NF1 clinical features in the studied p.Met1149-, p.Arg1276-, and p.Lys1423-positive cohorts with the “classic” NF1 population

NF1 feature	Risk ratio (95% CI)		
	p.Met1149 versus “classic” NF1	p.Arg1276 versus “classic” NF1	p.Lys1423 versus “classic” NF1
Skinfold freckling	0.73 (0.60–0.89)	0.78 (0.69–0.90)	
Lisch nodules	0.12 (0.04–0.35)	0.46 (0.31–0.68)	
Cutaneous neurofibromas	0.14 (0.05–0.40)	0.38 (0.25–0.59)	
Subcutaneous neurofibromas	0.24 (0.08–0.68)		
Symptomatic spinal neurofibromas		10.61 (6.26–17.98)	
Skeletal abnormalities		2.11 (1.52–2.91)	2.70 (2.00–3.64)
Noonan-like phenotype	8.57 (5.38–13.65)	6.13 (3.90–9.62)	8.54 (5.59–13.03)
Short stature			2.58 (1.78–3.74)
Pulmonic stenosis		11.10 (5.64–21.87)	13.44 (6.87–26.30)
Cardiovascular abnormalities		10.28 (6.56–16.12)	10.75 (6.72–17.20)

Abbreviation: CI, confidence interval.

the “classic” NF1 population; Table 4 and Tables S31 and S32), whereas is not significantly increased in individuals carrying pathogenic variants affecting codons 844–848 and 1149. As cardiovascular abnormalities may be congenital or become symptomatic at a young age and are associated with morbidity and mortality (Friedman et al., 2002; Lin et al., 2000), all individuals carrying one of the aforementioned pathogenic *NF1* missense variants should receive a detailed cardiac examination.

Watson syndrome (WS; MIM# 193520) was described in 1967 and is characterized by CALMs, PS, and intellectual disability, and is allelic to NF1 (Ben-Shachar et al., 2013; Watson, 1967). Among 24 cases with PS found in this report, one 12-year-old individual (UAB-R1256) presented solely with greater than five CALMs, PS, developmental delay, severe scoliosis, and short stature. Additionally, two children less than 4 years old (UAB-R37401FN.202 and UAB-R8917) had greater than five CALMs, PS, abnormal development with speech delay, and Noonan-like facial features as the only features. *NF1* c.3827G>A (p.Arg1276Gln), identified in the largest WS family from the original study (Ben-Shachar et al., 2013; Watson, 1967) was also found in UAB-R1256. Although 40% of the p.Arg1276-positive individuals ≥ 9 years old (29/73) had a severe phenotype with cardiac/cardiovascular abnormalities and/or spinal neurofibromas, 26% of cases ≥ 9 years old (19/73) presented with only pigmentary manifestations and a variable amount of learning disabilities (Table S6). Therefore, further studies aimed at identifying potential modifying genes, unlinked to the *NF1* locus, in these individuals should be considered.

Familial spinal neurofibromatosis (FSNF; MIM# 162210) is another rare subtype of NF1, characterized by the presence of multiple histopathologically confirmed neurofibromas along spinal nerve roots with few, if any, cutaneous neurofibromas, but often multiple firm subcutaneous tumors and CALMs (Korf, 2015; Pulst, Riccardi, Fain, & Korenberg, 1991; Ruggieri et al., 2015). Although pathogenic *NF1* missense and splicing variants are more commonly found in these individuals (Kluwe, Tatagiba, Fünsterer, & Mautner,

2003; Messiaen, Riccardi, & Peltonen, 2003; Upadhyaya et al., 2009), no specific recurrent variants associated with symptomatic spinal neurofibromas were identified, except recently for pathogenic missense variants affecting p.Gly848 (Koczkowska et al., 2018). In the current study, we identified p.Arg1276Gln being the second specific *NF1* variant associated with a high prevalence of symptomatic spinal neurofibromas (11-fold higher than in the general NF1 population; Table 4), found in 55.6% of individuals greater than 19 years (15/27; Table S10), including 4/15 with so-called spinal form of NF1 (Table S14 and Figure S5). An additional 4/6 p.Arg1276Gln-positive adults who underwent routine MRI screening had asymptomatic spinal tumors, and such individuals may possibly be eligible for treatment with MEK inhibitors. Similar to p.Gly848-positive individuals (Koczkowska et al., 2018), fewer individuals ≥ 19 years developed cutaneous neurofibromas compared with “classic” cohorts ($p < .0001$, significant at FDR of 0.01 after B-H correction; Table 2). Cutaneous neurofibromas were observed in only one-third of p.Arg1276Gln-positive cases with spinal tumors (7/19), ranging in number from 2 to 99, while the majority of NF1-affected individuals ≥ 30 years old with “classic” NF1 exhibit greater than 100 neurofibromas (Mautner et al., 2008). Only EUR-R32, carrying a different pathogenic variant, p.Arg1276Pro, presented with greater than 100 cutaneous and subcutaneous lesions at age of 60 years (Table S14). Among 22 individuals with spinal tumors, 16 cases had normal development in line with previous findings (Korf, 2015).

NF1 p.Arg1276 and p.Lys1423 are highly conserved residues lying in a shallow pocket in the central catalytic domain forming the RAS-binding region with the p.Arg1276 residue also called the arginine “finger,” being the most essential catalytic element for RasGAP activity (Scheffzek, Ahmadian, & Wiesmüller, 1998). Functional studies have shown that pathogenic variants affecting these residues result in a dramatic reduction of GAP activity (Pouillet, Lin, Esson, & Tamanoi, 1994; Scheffzek et al., 1998; Ahmadian, Kiel, Stege, & Scheffzek, 2003). Moreover, pathogenic missense variants at residue p.Lys1423 affects the stability of the neurofibromin/RAS

complex by disrupting an intramolecular salt-bridge of the GRD (Ahmadian et al., 2003).

In this study, we report an additional *NF1* variant, p.Met1149Val, associated specifically with a mild form of *NF1* without externally visible plexiform or symptomatic spinal neurofibromas, symptomatic OPGs or malignant neoplasms similar to the phenotypes associated with variants at p.Arg1809 and p.Met992del (Koczkowska et al., 2019; Pinna et al., 2015; Rojnueangnit et al., 2015; Upadhyaya et al., 2007). As only six probands were identified so far with either the p.Met1149Ile or p.Met1149Thr (Table S7), even larger datasets, as well as follow-up of individuals enrolled in the current study, are required to further refine a mild phenotype associated with the other substitutions at p.Met1149. Therefore, we have now shown that approximately 2.5% of all unrelated *NF1* pathogenic variant-positive probands of the UAB cohort develop a mild phenotype associated with nontruncating recurrent variants at one of three residues, p.Met992, p.Arg1809, and p.Met1149. Identification of this additional genotype–phenotype correlation will benefit the genetic counseling of these individuals and allow their stratification as they may not require stringent follow-up care. Moreover, as only 75% of p.Met1149-positive individuals fulfilled the NIH diagnostic criteria when family history was excluded as a criterion (Table S9), molecular analysis significantly facilitates the *NF1* diagnostic process and distinguishes Legius syndrome (MIM# 611431) from the p.Met992-del, p.Arg1809, and now p.Met1149 phenotypes with clinically overlapping features.

5 | CONCLUSION

Although single case reports may suggest an association of p.Arg1276 with spinal neurofibromas (Korf, Henson, & Stemmer-Rachamimov, 2005; Upadhyaya et al., 2009) or p.Lys1423 with Noonan-like features (De Luca et al., 2005) (two unrelated individuals described with each phenotype), only large datasets of postpubertal individuals, carrying the same constitutional pathogenic variant and with the phenotype recorded in a standardized way allow the establishment of clinically relevant genotype–phenotype correlations. International multicenter studies involving close collaborations between *NF1* clinicians and molecular geneticists are required to unfold such associations and to enhance correct variant interpretation. Our findings demonstrate genotype–phenotype correlations at the *NF1* codons p.Met1149, p.Arg1276, and p.Lys1423. Although each of the reported *NF1* variants associated with a specific clinical presentation (Kehrer-Sawatzki et al., 2017; Koczkowska et al., 2018; Koczkowska et al., 2019; Pinna et al., 2015; Rojnueangnit et al., 2015; Upadhyaya et al., 2007) affects only a small percentage of *NF1* individuals, together, including the current results, they affect counseling and management of approximately 10% of the *NF1* population. As the number of *NF1* genotype–phenotype correlations continues to accumulate, genotype-driven personalized medicine will reach a turning point in *NF1* by improving the disease surveillance and stratification of *NF1*-affected individuals.

ACKNOWLEDGMENT

We thank the individuals and their families for participating in this study.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Eva Trevisson  <http://orcid.org/0000-0002-5380-6265>

Michael F. Wangler  <http://orcid.org/0000-0001-5245-5910>

Kathleen B. M. Claes  <http://orcid.org/0000-0003-0841-7372>

Katharina Wimmer  <http://orcid.org/0000-0002-0732-7538>

Alessandro De Luca  <http://orcid.org/0000-0002-4408-8062>

Ludwine M. Messiaen  <http://orcid.org/0000-0002-1136-0154>

REFERENCES

- Ahmadian, M. R., Kiel, C., Stege, P., & Scheffzek, K. (2003). Structural fingerprints of the Ras-GTPase activating proteins neurofibromin and p120GAP. *Journal of Molecular Biology*, 329(4), 699–710.
- Allanson, J. E., Hall, J. G., Hughes, H. E., Preus, M., & Witt, R. D. (1985). Noonan syndrome: The changing phenotype. *American Journal of Medical Genetics*, 21(3), 507–514.
- Allanson, J. E., Hall, J. G., & Van Allen, M. I. (1985). Noonan phenotype associated with neurofibromatosis. *American Journal of Medical Genetics*, 21(3), 457–462.
- Ben-Shachar, S., Constantini, S., Halleivi, H., Sach, E. K., Upadhyaya, M., Evans, G. D., & Huson, S. M. (2013). Increased rate of missense/in-frame mutations in individuals with *NF1*-related pulmonary stenosis: A novel genotype-phenotype correlation. *European Journal of Human Genetics*, 21(5), 535–539.
- Bertola, D. R., Pereira, A. C., Passetti, F., de Oliveira, P. S. L., Messiaen, L., Gelb, B. D., ... Krieger, J. E. (2005). Neurofibromatosis-Noonan syndrome: Molecular evidence of the concurrence of both disorders in a patient. *American Journal of Medical Genetics*, 136(3), 242–245.
- Blanchard, G., Lafforgue, M. P., Lion-François, L., Kemlin, I., Rodriguez, D., Castelnau, P., ... Chaix, Y. (2016). Systematic MRI in *NF1* children under six years of age for the diagnosis of optic pathway gliomas. Study and outcome of a French cohort. *European Journal of Paediatric Neurology*, 20(2), 275–281.
- Blazo, M. A., Lewis, R. A., Chintagumpala, M. M., Frazier, M., McCluggage, C., & Plon, S. E. (2004). Outcomes of systematic screening for optic pathway tumors in children with neurofibromatosis type 1. *American Journal of Medical Genetics*, 127A(3), 224–229.
- Cnossen, M. H., de Goede-Bolder, A., van den Broek, K. M., Waasdorp, C. M. E., Oranje, A. P., Stroink, H., ... Niermeijer, M. F. (1998). A prospective 10 year follow up study of patients with neurofibromatosis type 1. *Archives of Disease in Childhood*, 78(5), 408–412.

- De Luca, A., Bottillo, I., Sarkozy, A., Carta, C., Neri, C., Bellacchio, E., ... Dallapiccola, B. (2005). *NF1* gene mutations represent the major molecular event underlying neurofibromatosis-Noonan syndrome. *The American Journal of Human Genetics*, 77(6), 1092–1101.
- Ekvall, S., Sjörs, K., Jonzon, A., Vihinen, M., Annerén, G., & Bondeson, M. L. (2014). Novel association of neurofibromatosis type 1-causing mutations in families with neurofibromatosis-Noonan syndrome. *American Journal of Medical Genetics, Part A*, 164A(3), 579–587.
- Evans, D. G., Howard, E., Giblin, C., Clancy, T., Spencer, H., Huson, S. M., & Laloo, F. (2010). Birth incidence and prevalence of tumor-prone syndromes: Estimates from a UK family genetic register service. *American Journal of Medical Genetics, Part A*, 152A(2), 327–332.
- Friedman, J. M. (2014). Neurofibromatosis 1. In Pagon, R. E., Adam, M. P., & Ardinger, H. H. (Eds.), *GeneReviews*[®]. Seattle, WA: University of Washington. <http://www.ncbi.nlm.nih.gov/books/NBK1109/>. Accessed June 21, 2018.
- Friedman, J. M., Arbiser, J., Epstein, J. A., Gutmann, D. H., Huot, S. J., Lin, A. E., ... Korf, B. R. (2002). Cardiovascular disease in neurofibromatosis 1: Report of the NF1 cardiovascular task force. *Genetics in Medicine*, 4(3), 105–111.
- Friedman, J. M., & Birch, P. H. (1997). Type 1 neurofibromatosis: A descriptive analysis of the disorder in 1,728 patients. *American Journal of Medical Genetics*, 70(2), 138–143.
- Huson, S. M., Compston, D. A., Clark, P., & Harper, P. S. (1989). A genetic study of von Recklinghausen neurofibromatosis in south east Wales. I. Prevalence, fitness, mutation rate, and effect of parental transmission on severity. *Journal of Medical Genetics*, 26(11), 704–711.
- Huson, S. M., Compston, D. A., & Harper, P. S. (1989). A genetic study of von Recklinghausen neurofibromatosis in south east Wales. II. *Guidelines for genetic counselling. J Med Genet*, 26(11), 712–721.
- Huson, S. M., Harper, P. S., & Compston, D. A. S. (1988). Von recklinghausen neurofibromatosis: A clinical and population study in south-east wales. *Brain*, 111(Pt 6), 1355–1381.
- Kehrer-Sawatzki, H., Mautner, V. F., & Cooper, D. N. (2017). Emerging genotype-phenotype relationships in patients with large *NF1* deletions. *Human Genetics*, 136(4), 349–376.
- Khosrotehrani, K., Bastuji-Garin, S., Riccardi, V. M., Birch, P., Friedman, J. M., & Wolkenstein, P. (2005). Subcutaneous neurofibromas are associated with mortality in neurofibromatosis 1: A cohort study of 703 patients. *American Journal of Medical Genetics, Part A*, 132A(1), 49–53.
- Kluwe, L., Tatagiba, M., Fünsterer, C., & Mautner, V. F. (2003). *NF1* mutations and clinical spectrum in patients with spinal neurofibromas. *Journal of Medical Genetics*, 40(5), 368–371.
- Koczkowska, M., Callens, T., Gomes, A., Sharp, A., Chen, Y., Hicks, A. D., ... Messiaen, L. M. (2019). Expanding the clinical phenotype of individuals with a 3-bp in-frame deletion of the *NF1* gene (c.2970_2972del): An update of genotype-phenotype correlation. *Genetics in Medicine*, 21(4), 867–876.
- Koczkowska, M., Chen, Y., Callens, T., Gomes, A., Sharp, A., Johnson, S., ... Messiaen, L. M. (2018). Genotype-phenotype correlation in *NF1*: Evidence for a more severe phenotype associated with missense mutations affecting *NF1* codons 844–848. *The American Journal of Human Genetics*, 102(1), 69–87.
- Korf, B. R. (2015). Spinal neurofibromatosis and phenotypic heterogeneity in *NF1*. *Clinical Genetics*, 87(5), 399–400.
- Korf, B. R., Henson, J. W., & Stemmer-Rachamimov, A. (2005). Case 13-2005: A 48-year-old man with weakness of the limbs and multiple tumors of spinal nerves. *New England Journal of Medicine*, 352(17), 1800–1808.
- Lammert, M., Friedman, J. M., Kluwe, L., & Mautner, V. F. (2005). Prevalence of neurofibromatosis 1 in German children at elementary school enrollment. *Archives of Dermatology*, 141(1), 71–74.
- Lin, A. E., Birch, P. H., Korf, B. R., Tenconi, R., Niimura, M., Poyhonen, M., ... Friedman, J. M. (2000). Cardiovascular malformations and other cardiovascular abnormalities in neurofibromatosis 1. *American Journal of Medical Genetics*, 95(2), 108–117.
- Listernick, R., Charrow, J., Greenwald, M., & Mets, M. (1994). Natural history of optic pathway tumors in children with neurofibromatosis type 1: A longitudinal study. *The Journal of Pediatrics*, 125(1), 63–66.
- Mautner, V. F., Asuagbor, F. A., Dombi, E., Fünsterer, C., Kluwe, L., Wenzel, R., ... Friedman, J. M. (2008). Assessment of benign tumor burden by whole-body MRI in patients with neurofibromatosis 1. *Neuro-Oncology*, 10(4), 593–598.
- McGaughran, J. M., Harris, D. I., Donnai, D., Teare, D., MacLeod, R., Westerbeek, R., ... Evans, D. G. (1999). A clinical study of type 1 neurofibromatosis in north west England. *Journal of Medical Genetics*, 36(3), 197–203.
- Messiaen, L. (2003). Independent *NF1* mutations in two large families with spinal neurofibromatosis. *Journal of Medical Genetics*, 40(2), 122–126.
- Messiaen, L. M., Callens, T., Mortier, G., Beysen, D., Vandenbroucke, I., Van Roy, N., ... Papee, A. D. (2000). Exhaustive mutation analysis of the *NF1* gene allows identification of 95% of mutations and reveals a high frequency of unusual splicing defects. *Human Mutation*, 15(6), 541–555.
- Messiaen, L. M., & Wimmer, K. (2012). Mutation analysis of the *NF1* gene by cDNA-based sequencing of the coding region. In K. S. G. Cunha, & M. Geller (Eds.), *Advances in neurofibromatosis research* (pp. 89–108). Hauppauge, NY: Nova Science Publishers.
- Nyström, A. M., Ekvall, S., Strömberg, B., Holmström, G., Thuresson, A. C., Annerén, G., & Bondeson, M. L. (2009). A severe form of Noonan syndrome and autosomal dominant café-au-lait spots—evidence for different genetic origins. *Acta Paediatrica*, 98(4), 693–698.
- Opitz, J. M., & Weaver, D. D. (1985). The neurofibromatosis-Noonan syndrome. *American Journal of Medical Genetics*, 21(3), 477–490.
- Pinna, V., Lanari, V., Daniele, P., Consoli, F., Agolini, E., Margiotti, K., ... De Luca, A. (2015). p.Arg1809Cys substitution in neurofibromin is associated with a distinctive *NF1* phenotype without neurofibromas. *European Journal of Human Genetics*, 23(8), 1068–1071.
- Plotkin, S. R., Bredella, M. A., Cai, W., Kassarian, A., Harris, G. J., Esparza, S., ... Mautner, V. F. (2012). Quantitative assessment of whole-body tumor burden in adult patients with neurofibromatosis. *PLoS One*, 7(4), e35711.
- Poulet, P., Lin, B., Esson, K., & Tamanoi, F. (1994). Functional significance of lysine 1423 of neurofibromin and characterization of a second site suppressor which rescues mutations at this residue and suppresses RAS2Val-19-activated phenotypes. *Molecular and Cellular Biology*, 14(1), 815–821.
- Pulst, S. M., Riccardi, V. M., Fain, P., & Korenberg, J. R. (1991). Familial spinal neurofibromatosis: Clinical and DNA linkage analysis. *Neurology*, 41(12), 1923–1923.
- Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., ... Rehm, H. L. (2015). Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in Medicine*, 17(5), 405–423.
- Rojnueangnit, K., Xie, J., Gomes, A., Sharp, A., Callens, T., Chen, Y., ... Messiaen, L. (2015). High incidence of Noonan syndrome features including short stature and pulmonic stenosis in patients carrying *NF1* missense mutations affecting p.Arg1809: Genotype-phenotype correlation. *Human Mutation*, 36(11), 1052–1063.
- Ruggieri, M., Polizzi, A., Spalice, A., Salpietro, V., Caltabiano, R., D'Orazi, V., ... Nicita, F. (2015). The natural history of spinal neurofibromatosis: A critical review of clinical and genetic features. *Clinical Genetics*, 87(5), 401–410.
- Santoro, C., Maietta, A., Giugliano, T., Melis, D., Perrotta, S., Nigro, V., & Piluso, G. (2015). Arg¹⁸⁰⁹ substitution in neurofibromin:

- Further evidence of a genotype-phenotype correlation in neurofibromatosis type 1. *European Journal of Human Genetics*, 23(11), 1460–1461.
- Scheffzek, K. (1998). Structural analysis of the GAP-related domain from neurofibromin and its implications. *EMBO Journal*, 17(15), 4313–4327.
- Thakkar, S. D., Feigen, U., & Mautner, V. F. (1999). Spinal tumours in neurofibromatosis type 1: An MRI study of frequency, multiplicity, and variety. *Neuroradiology*, 41(9), 625–629.
- Thiel, C., Wilken, M., Zenker, M., Sticht, H., Fahsold, R., Gusek-Schneider, G. C., & Rauch, A. (2009). Independent *NF1* and *PTPN11* mutations in a family with neurofibromatosis-Noonan syndrome. *American Journal of Medical Genetics*, 149(6), 1263–1267.
- Thissen, D., Steinberg, L., & Kuang, D. (2002). Quick and easy implementation of the Benjamini-Hochberg procedure for controlling the false positive rate in multiple comparisons. *Journal of Educational and Behavioral Statistics*, 27(1), 77–83.
- Upadhyaya, M., Huson, S. M., Davies, M., Thomas, N., Chuzhanova, N., Giovannini, S., ... Messiaen, L. (2007). An absence of cutaneous neurofibromas associated with a 3-bp inframe deletion in exon 17 of the *NF1* gene (c.2970–2972 delAAT): Evidence of a clinically significant *NF1* genotype-phenotype correlation. *The American Journal of Human Genetics*, 80(1), 140–151.
- Upadhyaya, M., Spurlock, G., Kluwe, L., Chuzhanova, N., Bennett, E., Thomas, N., ... Mautner, V. (2009). The spectrum of somatic and germline *NF1* mutations in *NF1* patients with spinal neurofibromas. *Neurogenetics*, 10(3), 251–263.
- Uusitalo, E., Leppävirta, J., Koffert, A., Suominen, S., Vahtera, J., Vahlberg, T., ... Peltonen, S. (2015). Incidence and mortality of neurofibromatosis: A total population study in Finland. *Journal of Investigative Dermatology*, 135(3), 904–906.
- Watson, G. H. (1967). Pulmonary stenosis, cafe-au-lait spots, and dull intelligence. *Archives of Disease in Childhood*, 42(223), 303–307.
- Zhou, X., Edmonson, M. N., Wilkinson, M. R., Patel, A., Wu, G., Liu, Y., ... Zhang, J. (2016). Exploring genomic alteration in pediatric cancer using ProteinPaint. *Nature Genetics*, 48(1), 4–6.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Koczkowska M, Callens T, Chen Y, et al. Clinical spectrum of individuals with pathogenic *NF1* missense variants affecting p.Met1149, p.Arg1276, and p.Lys1423: genotype-phenotype study in neurofibromatosis type 1. *Human Mutation*. 2020;41:299–315.
<https://doi.org/10.1002/humu.23929>