This item is the archived peer-reviewed author-version of:

Sensitization profiles to peanut allergens in Belgium : cracking the code in infants, children and adults

Reference:

Faber Margaretha, Donné Inne, Herrebosch Evelien, Sabato Vito, Hagendorens Margo, Bridts Christiaan, De Clerck Luc S., Ebo Didier.- Sensitization profiles to peanut allergens in Belgium : cracking the code in infants, children and adults
Acta clinica Belgica - ISSN 0001-5512 - (2016), p. 1-7

Full text (Publishers DOI): <http://dx.doi.org/doi:10.1080/17843286.2015.1109170>

To cite this reference: <http://hdl.handle.net/10067/1328580151162165141>

Sensitization profiles to peanut allergens in Belgium; cracking the code in infants, children and adults

Margaretha A. Faber¹, Inne Donné¹, Evelien Herrebosch¹, Vito Sabato¹, Margo M. Hagendorens^{1,2}, Chris H. Bridts¹, Luc S. De Clerck¹, Didier G. Ebo¹

¹ Department of Immunology-Allergology-Rheumatology, Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp (Belgium) and, Antwerp University Hospital, Antwerp (Belgium)

² Department of Pediatrics, University of Antwerp and Antwerp University Hospital, Antwerp (Belgium)

Key words: Ara h 2; Component Resolved Diagnosis; Food allergy; Peanut allergy

Conflict of interest: none

Corresponding author;

Professor Dr. D. G. Ebo,

Department of Immunology, Allergology, Rheumatology, University of Antwerp (Faculty of Medicine and Health Science)

Campus Drie Eiken T.595

Universiteitsplein 1

2610 Antwerpen, Belgium.

Email: immuno@uantwerpen.be

18 **Acknowledgement**

19 The authors thank Mrs. Christel Mertens for her technical skills and Karin Van Cotthem for
20 her contribution to the sIgE determination. Vito Sabato is a Clinical Researcher of the
21 Research Foundation Flanders (FWO: 1700614N). Didier Ebo is a Senior Clinical Researcher
22 of the Research Foundation Flanders (FWO: 1800614N).

23

24

25

26

27

28

29

30

31

32

33

34

35

Abstract

Objectives

Peanut allergy shows distinct clinical patterns that can be predicted by component resolved diagnosis. However data about peanut sensitization profiles in populations with a broad age-stratification are scarce.

Methods

Sera of 89 peanut allergic patients (age 1 – 70 years), 21 infants (< 1y) with atopic dermatitis (AD) sensitized to peanut, 24 age matched peanut tolerant individuals with positive sIgE to peanut and 15 healthy individuals were tested for sIgE reactivity to rAra h 1, rAra h 2, rAra h 3, rAra h 8, rAra h 9 and rBet v 1 (FEIA ImmunoCAP, Thermo Fisher Scientific).

Results

In infants with AD, Ara h 1, Ara h 2 and Ara h 3 enabled to explain 14/21 (67%) of peanut sensitizations. No sensitization to Ara h 8 or Bet v 1 was observed. Patients with generalized reactions were more frequently sensitized to Ara h 1, Ara h 2 and Ara h 3 compared to patients with an oral allergy syndrome (OAS) and peanut tolerant patients. Sensitization to Ara h 8 was significantly more observed in patients with an OAS. Ara h 2 showed to be the best marker to distinguish patients with generalized reactions from patients with an OAS and/or peanut sensitized patients but tolerating the legume.

Conclusion

Sensitization to Ara h 1, Ara h 2 and Ara h 3 can have early onset and is predominantly associated with a more severe outcome. Ara h 2 is the best marker of a generalized peanut allergy.

Introduction

Peanut (*Arachis hypogaea*) constitutes an increasing cause of food allergy in children and adults^{1, 2}. The clinical presentation of peanut allergy is highly variable and can vary from localized allergic reactions such as an oral allergy syndrome (OAS) to severe generalized reactions, which can be life-threatening^{3, 4}. Given the severity and impact of peanut allergy, correct diagnosis is mandatory.

In the last decade, different studies have shown that additional quantification of specific IgE (sIgE) antibodies to the different allergenic components of peanut by component resolved diagnosis (CRD) is more reliable to predict clinical outcomes than only conventional assays measuring sIgE to whole peanut extracts⁵⁻¹¹.

Overall, Ara h 2, a member of the prolamin superfamily, has been described as the most important component for discrimination between peanut allergic patients with generalized symptoms and peanut tolerant individuals^{5, 6, 10-15}. Other clinically important peanut components, also predominantly associated with generalized allergic reactions, are Ara h 1 and Ara h 3, both members of the cupin superfamily and Ara h 6, another member of the prolamin superfamily that is displaying a high sequence identity with Ara h 2^{10, 12, 16}.

In southern Europe, Ara h 9, the non-specific lipid transfer protein (ns-LTP) of peanut, has been described as a relevant peanut allergen and sensitization to this allergen is oftentimes observed to be associated with severe allergic reactions¹⁷. In contrast, in northern Europe patients frequently present with a mild OAS resulting from a cross-reactivity between Bet v 1 the major allergen component from birch (*Betula verrucosa*) pollen and its homologue Ara h 8 from peanut, both proteins belonging to the pathogenesis related (PR10) protein group^{11, 17}.

Besides geographic differences, it has been observed that food allergy can also exhibit distinct age-dependent sensitization profiles^{11, 18}. However, for the time being, the large majority of studies assessing the peanut sensitization profile has been performed in pediatric populations whereas data in adults remain scarce¹⁹. Therefore, in this study we aimed (1) at evaluating the sensitization patterns to five peanut components in a population ranging from infants (< 1 year of age), over preschool- and school-children to adults with different clinical presentations, and (2) to assess the predictive value of sIgE quantification against the various peanut components.

118

119 **Methods**

120 *Subjects*

121 A total of 150 individuals were enrolled via the outpatients' clinics of Allergology and
122 Pediatrics of the Antwerp University Hospital.

123 The study group consisted of infants less than 1 year old with atopic dermatitis (AD), all with
124 a sIgE to whole peanut extract ≥ 0.35 kUa/L (FEIA ImmunoCAP, Thermo Fisher Scientific,
125 Uppsala, Sweden) were selected. Second, patients with a definite history of immediate
126 allergic symptoms due to consumption of raw and/or processed peanut and a sIgE to whole
127 peanut extract ≥ 0.35 kUa/L were included. These patients were stratified into 6 groups
128 according to the extent of the clinical reaction (OAS or generalized reaction) and age ($< 7y$, 7-
129 18y; $> 18y$) as detailed elsewhere¹⁸. Third, peanut tolerant individuals with a sIgE to whole
130 peanut extract ≥ 0.35 kUa/L were included and finally non-allergic age-matched individuals
131 were enrolled. At the time of inclusion, all peanut tolerant individuals indicated to consume
132 peanut regularly without allergic symptoms.

133 The diagnosis of a generalized peanut allergy was based upon a compelling history of an
134 adverse reaction upon roasted or processed peanut. Allergic symptoms could involve skin or
135 subcutaneous tissues (e.g. generalized erythema, urticaria, or angioedema) and/or the
136 respiratory tract (e.g. dyspnea, stridor, wheezing, chest/throat tightness or cyanosis) and/or the
137 gastrointestinal system (e.g. nausea, vomiting or abdominal pain) and/or the cardiovascular
138 system (dizziness, diaphoresis, hypotension, confusion, loss of consciousness)²⁰. The
139 diagnosis of an OAS to peanut relied upon a compelling history of repetitive pruritus and/or
140 angioedema of the lips, tongue and/or palate on consumption of peanut.

Challenges were not conducted neither in patients with a compelling history of generalized reactions due to the severity of the reported symptoms and the potential risk of eliciting serious reactions, nor in OAS, since challenges were deemed unnecessary as the clinical history in such cases was highly reliable and symptoms were easily recognized and described by the patient.

The local ethics committee approved this study (B300201316182). Patients, healthy controls or their representatives approved an informed consent in accordance with the Declaration of Helsinki.

Total and specific IgE

Total IgE and sIgE to the whole peanut extract, the recombinant (r) peanut components rAra h 1, rAra h 2, rAra h 3, rAra h 8 and rAra h 9 and recombinant birch pollen major allergen rBet v 1 were quantified by the FEIA ImmunoCAP technique (Thermo Fisher Scientific) according to the manufacturer's instructions.

Calculation of predictive values

We calculated predictive values of different peanut components and a combination thereof to evaluate their ability to discriminate patients with generalized allergic reactions from patients with OAS and individuals tolerant to peanut but demonstrating a positive sIgE to this legume.

Statistical analysis

Data are expressed as median and range. IBM SPSS 20 software was used for data analysis. Evaluation of the diagnostic performance of the different peanut components in discriminating peanut allergic patients from healthy control individuals was performed by

using receiver operating characteristic (ROC) curve analysis. Non-parametric tests and χ^2 analysis were used where appropriate. A $P < 0.05$ was regarded as statistically significant.

Results

Patient characteristics

Table 1 summarizes the demographics of patients and healthy control individuals. In total, 68 patients exhibited a generalized reaction to peanut ingestion, whereas 21 patients reported an OAS to peanut. We were unable to enroll sufficient numbers of children younger than 7 years old with an OAS for robust analysis.

ROC analysis

ROC analyses between patients with generalized reactions upon peanut consumption and healthy control individuals generated a cut-off value of 0.10 kUa/L for all components. As a consequence, 0.10 kUa/L was applied as a decision threshold for further analysis of the sensitization profiles and for calculation of predictive values.

Sensitization to Ara h 1

As displayed in Figure 1, sensitization towards Ara h 1 was mainly observed in patients with generalized reactions as compared to OAS patients (χ^2 analysis, $P < 0.05$). None of the patients with generalized reactions were monosensitized to Ara h 1. Sensitization to Ara h 1 can have an early onset as it was observed in 12/21 (57%) infants with AD. Three out of 24 peanut tolerant individuals (13%) with a sensitization to whole peanut extract, showed sIgE reactivity to Ara h 1.

183

184 **Sensitization to Ara h 2**

185 As demonstrated in Figure 1., similarly to Ara h 1, sensitization to Ara h 2 was significantly
 186 more observed in patients with generalized allergic reactions compared to patients with an
 187 isolated OAS (χ^2 analysis, $P < 0.05$). Monosensitization to Ara h 2 was seen in only 10/68 (7%)
 188 patients with generalized reactions. With respect to age, sensitization to Ara h 2 was
 189 significantly more frequent in preschool and school children than in adults with generalized
 190 reactions (χ^2 analysis, $P < 0.05$). As a matter of fact all 32 preschool children, 24/25 (96%)
 191 schoolchildren and 8/11 (73%) adults with generalized reactions showed sIgE reactivity to
 192 Ara h 2.

193 Six of the 21 infants with AD (29%) were sensitized to Ara h 2. Five out of twentyfour peanut
 194 tolerant individuals with a positive sIgE to peanut (21%) showed IgE reactivity to rAra h 2.

195 **Sensitization to Ara h 3**

196 Sensitization to Ara h 3 was also predominantly observed in patients with generalized
 197 reactions compared to patients with an OAS (χ^2 analysis, $P < 0.05$). None of the patients with
 198 generalized reactions showed a monosensitization to Ara h 3.

199 Sensitization to the legumin Ara h 3 was observed in 8/21 (38 %) infants with AD (Fig. 1).
 200 Furthermore, 3 patients in the group of peanut tolerant but sensitized individuals (13%)
 201 showed sIgE antibodies to Ara h 3.

202 **Sensitization to Bet v 1 and its peanut homologue Ara h 8**

203 Unlike sensitization to Ara h 1, Ara h 2 and Ara h 3, sensitization to the Bet v 1 homologue,
 204 Ara h 8, was significantly more observed in patients with an OAS versus patients with

generalized reactions upon peanut (χ^2 analysis , $P < 0.05$). However, it should be noticed that only 3 patients with generalized reactions were monosensitized to the Bet v 1 homologue, the remainder patients demonstrated a cosensitization to Ara h 1 and/or Ara h 2 and/or Ara h 3 and/or Ara h 9.

Sensitization to Ara h 8 is significant correlated with sensitization to Bet v 1 (Pearson correlation coefficient = 0.81). In the group of peanut tolerant patients 20/24 (83%) showed a positive sIgE to Ara h 8 and 22/24 (92%) were sensitized to Bet v 1. In the group of infants with AD and demonstrating a positive sIgE to crude peanut no sensitization to Bet v 1 and Ara h 8 was demonstrable.

Sensitization to Ara h 9

Sensitization to Ara h 9 was found in 9 out of the 68 patients (13%) with generalized reaction and in 4/21 (19%) of patients with an OAS. Neither in the group of patients with generalized reactions nor in the group of patients with an OAS there was a monosensitization to the ns-LTP of peanut demonstrable. In the group of peanut tolerant individuals 5/24 (21%) were sensitized to Ara h 9. One infant with AD was sensitized to the ns-LTP of peanut.

Sensitization to Ara h 1 and/or Ara h 2 and/or Ara h 3

When evaluating sIgE to Ara h 1 and/or Ara h 2 and/or Ara h 3, a positive result was found in all 46 preschool children, in 24/25 (96%) school children and in 9/11 (82%) of adults with generalized reactions. Furthermore at least 1 out of 3 storage proteins were recognized in 14/21 infants with AD.

In contrast, only 3/21 of patients with an OAS (14 %) ($P < 0.05$) and 6/24 peanut tolerant patients with a positive sIgE to peanut (25%) ($P < 0.05$) were sensitized to Ara h 1 and/or Ara h 2 and/or Ara h 3.

Predictive values of Ara h1 and/or Ara h 2 and/or Ara h 3

Because sensitization to Ara h 1, Ara h 2 and Ara h 3 was significantly more seen in patients with generalized reactions, we wondered whether these components or a combination thereof could be predictive for a more severe clinical outcome. Therefore, positive and negative predictive values (PPV and NPV) were calculated for these components between patients with generalized allergic reactions versus patients with an OAS and peanut tolerant patients with a positive sIgE to peanut.

As demonstrated in Table 2, applying the ROC-generated decision thresholds of 0.10 kUa/L we found highest overall PPV and NPV for Ara h 2. As a matter of fact, the NPV of Ara h 2 was found to be absolute in preschool children. Furthermore, combining the sIgE results of Ara h 1 and/or Ara h 2 and/or Ara h 3 did not significantly increase the PPV and/or NPV of the component resolved diagnosis technique.

Correlation between different peanut components

Figure 2. shows a hierarchic cluster analysis of the sensitization to different peanut components. As expected, sensitization to the major birch pollen allergen Bet v 1, and sensitization to the Bet v 1 homologue Ara h 8 is highly correlated. In the group of seed storage proteins there is a high correlation between sensitization to Ara h 1 and sensitization to Ara h 2 (Pearson correlation coefficient = 0.83), furthermore sensitization to Ara h 3 is correlated to sensitization to Ara h 1.

Discussion

To our knowledge this is the first study that investigates peanut sensitization profiles using 5 components in an age-differentiated population composed out of infants, preschool- and school-children and adults. It reveals several particularities. First it confirms that patients with peanut allergy can display various sensitization profiles with distinct clinical outcomes. Sensitization to the 2S albumin, Ara h 2 and to members of the bicupin family, such as the vicilin Ara h 1 and legumin Ara h 3, is predominantly, but certainly not uniquely, associated with more generalized reactions. Whereas a sensitization to the Bet v 1 homologue Ara h 8, is generally associated with an OAS⁵⁻¹⁵.

Second, this study reveals that these sensitization profiles also exhibit significant age variations. Sensitization to Ara h 1, Ara h 2 and Ara h 3, unlike sensitization to the Bet v 1 homologue Ara h 8, can already start in infants without an overt peanut exposure as two-thirds of our group of infants with atopic dermatitis < 1 year of age with a positive sIgE to peanut extract was sensitized to Ara h 1 and/or Ara h 2 and/or Ara h 3. The exact mechanisms and routes of exposure for these early sensitizations remain elusive but probably relate to cutaneous exposure to peanut allergen present in for example house dust or in peanut derived cosmetic products^{21, 22} and/or sensitization due to peanut exposure during pregnancy or breastfeeding^{23, 24}. In contrast, the observation that sensitization to the Bet v 1 homologue, Ara h 8 is absent in infants is likely to be explained by the fact that sensitization to the major birch pollen allergen generally starts at later age in childhood²⁵. In any way, longitudinal follow-up studies seem mandatory in order to ascertain the clinical significance and prognosis of a sensitization against Ara h 1, Ara h 2 and Ara h 3 in infants without overt peanut allergy. These studies should focus on the fact whether infants with atopic dermatitis demonstrating such a sensitization pattern are at particular risk to develop generalized peanut allergies or do

outgrow their sensitization during childhood. In addition, longitudinal studies are also mandatory to explain the observation that adults seem less sensitized to Ara h 2 as compared to children. From our cross-sectional data it cannot be concluded whether this lower prevalence of Ara h 1 and Ara h 2 sensitization is due to a genuine outgrow^{26, 27} or merely reflects a loss of medical follow-up, i.e. adults with established peanut allergy not attending the clinics anymore.

Third, it is well-known that peanut allergy can constitute a severe condition with potentially fatal outcome⁴. On the other hand, overdiagnosis of peanut allergy with the set-up of unnecessary dietary prevention measures should be avoided. Therefore, correct identification of patients with more severe allergic reactions upon peanut is of paramount importance.

The most significant finding of our study is that Ara h 2 is the best marker to distinguish patients with generalized reactions from patients with an OAS and/or peanut sensitized patients but tolerating the legume. As a matter of fact, the negative predictive value of Ara h 2 was absolute in preschool children, meaning that the absence of sIgE reactivity to Ara h 2 precludes a more severe generalized reaction in the youngest peanut allergic children. Furthermore, this study shows that addition of sIgE measurement to Ara h 1 and Ara h 3, does not significantly improve the capacity of the component resolved diagnosis technique to identify patients with generalized reactions compared to sIgE measurement to Ara h 2 alone.

In contrast, monosensitization to the Bet v 1 homologue Ara h 8 in our cohort was predominantly associated with an OAS. Alternatively, it should be kept in mind that in a region where birch trees are endemic that sIgE antibodies to Bet v 1 homologues were confirmed to be the most prevalent cause of clinically irrelevant sIgE results and not to contribute to discriminate between birch pollen allergic individuals with or without peanut allergy²⁸.

A sensitization rate of $\geq 10\%$ to Ara h 9 was observed in the group of peanut allergic individuals with generalized reactions as well as in the group of patients with an OAS upon peanut, although sensitization to the ns-LTP of peanut was also observed in the group of peanut tolerant patients. To be conclusive about clinical correlation, more studies are needed.

A potential criticism on our study could be the absence of double-blind placebo controlled challenges (DBPCFC) with peanut. However, from the literature it appears that even DBPCFC's are not absolutely predictive for the clinical outcome, as both false negative and false positive food challenges have been described in approximately 5% and 13% of cases 29-31. Second, almost half of our population are preschool children aged less than 7 years in whom DBPCFC remains particularly dangerous. Finally, in patients with a compelling history of OAS upon peanut, challenges were deemed unnecessary as the clinical history in such cases is highly reliable and symptoms are easily recognized and described by the patients.

Another potential criticism could be that skin tests have not been systematically carried out in this study, therefore no robust and reliable statistical analysis in order to evaluate the diagnostic value of skin prick tests could be done. However, it is the authors' experience that in children, not sensitized to pollen, skin prick tests can add to the diagnosis, particularly in cases where quantification of sIgE yields borderline positive results. In contrast, in adolescent and adults, skin testing frequently yields clinically irrelevant responses mainly because of the extensive cross-reactivity with birch pollen components such as the major birch pollen allergen Bet v 1 and profilin (Bet v 2). To our knowledge no purified or recombinant peanut components are available for skin testing.

In conclusion, this study emphasizes the importance of a broad age stratification when assessing sensitization profiles in food allergy. First it reveals that, unlike Ara h 8, sensitization to Ara h 1, 2 and 3 can occur in infants without overt peanut allergy. Second, it

confirms that sensitization to Ara h 1,2 and 3 is predominantly associated with a more severe outcome, whereas sensitization to Ara h 8 is generally associated with an OAS. Finally, and most importantly, our study indicates that Ara h 2 is the best marker to identify peanut allergic patients in risk of generalized reactions.

REFERENCES

1. Sicherer SH, Munoz-Furlong A, Godbold JH, Sampson HA. US prevalence of self-reported peanut, tree nut, and sesame allergy: 11-year follow-up. *The Journal of allergy and clinical immunology*. 2010;125(6):1322-6.
2. Bunyavanich S, Rifas-Shiman SL, Platts-Mills TA, Workman L, Sordillo JE, Gillman MW, et al. Peanut allergy prevalence among school-age children in a US cohort not selected for any disease. *The Journal of allergy and clinical immunology*. 2014;134(3):753-5.
3. Burks AW. Peanut allergy. *Lancet*. 2008;371(9623):1538-46.
4. Bock SA, Munoz-Furlong A, Sampson HA. Further fatalities caused by anaphylactic reactions to food, 2001-2006. *The Journal of allergy and clinical immunology*. 2007;119(4):1016-8.
5. Codreanu F, Collignon O, Roitel O, Thouvenot B, Sauvage C, Vilain AC, et al. A novel immunoassay using recombinant allergens simplifies peanut allergy diagnosis. *International archives of allergy and immunology*. 2011;154(3):216-26.
6. Nicolaou N, Murray C, Belgrave D, Poorafshar M, Simpson A, Custovic A. Quantification of specific IgE to whole peanut extract and peanut components in prediction of peanut allergy. *The Journal of allergy and clinical immunology*. 2011;127(3):684-5.
7. Klemans RJ, Broekman HC, Knol EF, Bruijnzeel-Koomen CA, Otten HG, Pasmans SG, et al. Ara h 2 is the best predictor for peanut allergy in adults. *The journal of allergy and clinical immunology In practice*. 2013;1(6):632-8 e1.
8. Lieberman JA, Glaumann S, Batelson S, Borres MP, Sampson HA, Nilsson C. The utility of peanut components in the diagnosis of IgE-mediated peanut allergy among distinct populations. *The journal of allergy and clinical immunology In practice*. 2013;1(1):75-82.
9. Dang TD, Tang M, Choo S, Licciardi PV, Koplin JJ, Martin PE, et al. Increasing the accuracy of peanut allergy diagnosis by using Ara h 2. *The Journal of allergy and clinical immunology*. 2012;129(4):1056-63.
10. Eller E, Bindslev-Jensen C. Clinical value of component-resolved diagnostics in peanut-allergic patients. *Allergy*. 2013;68(2):190-4.
11. Ballmer-Weber BK, Lidholm J, Fernandez-Rivas M, Seneviratne S, Hanschmann KM, Vogel L, et al. IgE recognition patterns in peanut allergy are age dependent: perspectives of the EuroPrevall study. *Allergy*. 2015.
12. Ebisawa M, Moverare R, Sato S, Maruyama N, Borres MP, Komata T. Measurement of Ara h 1-, 2-, and 3-specific IgE antibodies is useful in diagnosis of peanut allergy in Japanese children. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2012;23(6):573-81.
13. Klemans RJ, Liu X, Knulst AC, Knol MJ, Gmelig-Meyling F, Borst E, et al. IgE binding to peanut components by four different techniques: Ara h 2 is the most relevant in peanut allergic children and adults. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2013;43(8):967-74.
14. Ackerbauer D, Bublin M, Radauer C, Varga EM, Hafner C, Ebner C, et al. Component-resolved IgE profiles in Austrian patients with a convincing history of peanut allergy. *International archives of allergy and immunology*. 2015;166(1):13-24.

15. Kukkonen AK, Pelkonen AS, Makinen-Kiljunen S, Voutilainen H, Makela MJ. Ara h 2 and Ara 6 are the best predictors of severe peanut allergy: A double-blind placebo-controlled study. *Allergy*. 2015.
16. Klemans RJ, Knol EF, Bruijnzeel-Koomen CA, Knulst AC. The diagnostic accuracy of specific IgE to Ara h 6 in adults is as good as Ara h 2. *Allergy*. 2014;69(8):1112-4.
17. Vereda A, van Hage M, Ahlstedt S, Ibanez MD, Cuesta-Herranz J, van Odijk J, et al. Peanut allergy: Clinical and immunologic differences among patients from 3 different geographic regions. *J Allergy Clin Immunol*. 2011;127(3):603-7.
18. Faber MA, De Graag M, Van Der Heijden C, Sabato V, Hagendorens MM, Bridts CH, et al. Cor a 14: Missing Link in the Molecular Diagnosis of Hazelnut Allergy? *International archives of allergy and immunology*. 2014;164(3):200-6.
19. Klemans RJ, van Os-Medendorp H, Blankestijn M, Bruijnzeel-Koomen CA, Knol EF, Knulst AC. Diagnostic accuracy of specific IgE to components in diagnosing peanut allergy: a systematic review. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2014.
20. Brown SG. Clinical features and severity grading of anaphylaxis. *The Journal of allergy and clinical immunology*. 2004;114(2):371-6.
21. Brough HA, Liu AH, Sicherer S, Makinson K, Douiri A, Brown SJ, et al. Atopic dermatitis increases the effect of exposure to peanut antigen in dust on peanut sensitization and likely peanut allergy. *The Journal of allergy and clinical immunology*. 2015;135(1):164-70 e4.
22. Lack G, Fox D, Northstone K, Golding J, Avon Longitudinal Study of P, Children Study T. Factors associated with the development of peanut allergy in childhood. *The New England journal of medicine*. 2003;348(11):977-85.
23. McGowan EC, Bloomberg GR, Gergen PJ, Visness CM, Jaffee KF, Sandel M, et al. Influence of early-life exposures on food sensitization and food allergy in an inner-city birth cohort. *The Journal of allergy and clinical immunology*. 2015;135(1):171-8 e4.
24. Sicherer SH, Wood RA, Stablein D, Lindblad R, Burks AW, Liu AH, et al. Maternal consumption of peanut during pregnancy is associated with peanut sensitization in atopic infants. *The Journal of allergy and clinical immunology*. 2010;126(6):1191-7.
25. Flinterman AE, Akkerdaas JH, den Hartog Jager CF, Rigby NM, Fernandez-Rivas M, Hoekstra MO, et al. Lipid transfer protein-linked hazelnut allergy in children from a non-Mediterranean birch-endemic area. *The Journal of allergy and clinical immunology*. 2008;121(2):423-8 e2.
26. Fleischer DM. The natural history of peanut and tree nut allergy. *Current allergy and asthma reports*. 2007;7(3):175-81.
27. Fleischer DM, Conover-Walker MK, Christie L, Burks AW, Wood RA. The natural progression of peanut allergy: Resolution and the possibility of recurrence. *The Journal of allergy and clinical immunology*. 2003;112(1):183-9.
28. Ebo DG, Bridts CH, Verweij MM, De Knop KJ, Hagendorens MM, De Clerck LS, et al. Sensitization profiles in birch pollen-allergic patients with and without oral allergy syndrome to apple: lessons from multiplexed component-resolved allergy diagnosis. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology*. 2010;40(2):339-47.
29. Vlieg-Boerstra BJ, van der Heide S, Bijleveld CM, Kukler J, Duiverman EJ, Dubois AE. Placebo reactions in double-blind, placebo-controlled food challenges in children. *Allergy*. 2007;62(8):905-12.
30. Asero R, Fernandez-Rivas M, Knulst AC, Bruijnzeel-Koomen CA. Double-blind, placebo-controlled food challenge in adults in everyday clinical practice: a reappraisal of their limitations and real indications. *Current opinion in allergy and clinical immunology*. 2009;9(4):379-85.
31. Oole-Groen CJ, Brand PL. Double-blind food challenges in children in general paediatric practice: useful and safe, but not without pitfalls. *Allergologia et immunopathologia*. 2014;42(4):269-74.

		No. (Female)	Age (years) Median (range)	Total IgE (kU/L) Median (range)
Peanut allergic patients with generalized reactions (sIgE peanut extract $\geq$ 0.35 kUa/L)	Preschool	32 (10)	3.9 (0.7 - 6.7)	202 (10 - 13450)
	School	25 (11)	11.3 (7.1 - 17.9)	441 (20 - 28500)
	Adults	11 (4)	21.5 (18.0 – 38.4)	642 (58 - 6500)
	Total	68 (25)		
Peanut allergic patients with localized reactions (OAS) (sIgE peanut extract $\geq$ 0.35 kUa/L)	Preschool	1 (1)	*	*
	School	7 (3)	13.4 (6.3 - 16.0)	422 (211 - 833)
	Adults	14 (12)	42.3 (20.1 - 69.5)	133 (20 - 1702)
	Total	22 (16)		
Infants with atopic dermatitis (sIgE peanut extract $\geq$ 0.35 kUa/L)	$\leq$ 1 year	21 (7)	0.6 (0.5 - 1.0)	159 (9 - 4149)
Peanut tolerant individuals (sIgE peanut extract $\geq$ 0.35 kUa/L)	Preschool	4 (2)	5.7 (4.9 - 6.3)	606 (270 - 4292)
	School	8 (5)	10.8 (8.2 - 14.3)	746 (531 - 2206)
	Adults	12 (8)	35.4 (18.1 - 56.2)	311 (76 - 5800)
	Total	24 (15)		
Healthy control individuals (sIgE peanut extract < 0.35 kUa/L)	Preschool	5 (4)	6.4 (2.8 - 6.5)	28 (16 - 278)
	School	5 (2)	10.0 (8.2 - 15.8)	182 (8 - 290)
	Adults	5 (5)	32.4 (22.0 - 44.3)	7 (2 - 73)
	Total	15 (11)		
TOTAL		150 (74)		

Table 1. Demographics and characteristics

*No sufficient number of preschool children with an Oral Allergy Syndrome (OAS) were included for robust analysis.

440

441

442

Table 2. Performance characteristics of specific IgE to Ara h 1, Ara h 2 and Ara h 3.

To differentiate:	Age (years)	rAra h 1		rAra h 2		rAra h 3		rAra h 1 and/or rAra h 2 and/or rAra h 3	
	</								

OAS: Oral Allergy Syndrome, PPV: Positive Predictive Value, NPV: Negative Predictive Value