

This item is the archived peer-reviewed author-version of:

Interspecific variation in the tetradactyl manus of modern tapirs (**Perissodactyla: Tapirus**) exposed using geometric morphometrics

Reference:

Maclaren Jamie, Nauw elaerts Sandra.- Interspecific variation in the tetradactyl manus of modern tapirs (**Perissodactyla: Tapirus**) exposed using geometric morphometrics

Journal of morphology - ISSN 0362-2525 - 278:11(2017), p. 1517-1535

Full text (Publisher's DOI): https://doi.org/10.1002/JMOR.20728

To cite this reference: https://hdl.handle.net/10067/1455380151162165141

uantwerpen.be

Institutional repository IRUA

Page 1 of 53

Journal of Morphology

1	
2	
3	
4	
5	
6	
7	
0	
0	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
21	
22 22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
27	
20	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52 52	
03	
54 55	
55	
56	
57	
58	
59	
60	

1	Interspecific variation in the tetradactyl manus of modern tapirs
2	(Perissodactyla: Tapirus) exposed using geometric morphometrics
3	Authors: Jamie A. MacLaren ¹ * and Sandra Nauwelaerts ^{1 & 2}
4	
5	Affiliation:
6	¹ Department of Biology, Universiteit Antwerpen, Building D, Campus Drie Eiken,
7	Universiteitsplein, Wilrijk, Antwerp, 2610, Belgium
8	² Centre for Research and Conservation, Koninklijke Maatschappij voor Dierkunde (KMDA),
9	Koningin Astridplein 26, Antwerp, 2018, Belgium
10	
11	* Corresponding author
12	
13	Corresponding Author: Jamie A. MacLaren, Room D1.41, Building D, Campus Drie Eiken,
14	Universiteitsplein 1, 2610 Antwerp, Belgium
15	Correspondence Email: jamie.maclaren@uantwerpen.be
16	Correspondence Telephone: (+32) 32 65 1994
17	
18	
19	Short Title: Shape variation in the tapir manus
20	.L
20	

MacLaren 2

Abstract

The distal forelimb (autopodium) of guadrupedal mammals is a key morphological unit involved in locomotion, body support and interaction with the substrate. The manus of the tapir (Perissodactyla: *Tapirus*) is unique within modern perissodactyls, as it retains the plesiomorphic tetradactyl (four-toed) condition also exhibited by basal equids and rhinoceroses. Tapirs are known to exhibit anatomical mesaxonic symmetry in the manus, although interspecific differences and biomechanical mesaxony has yet to be rigorously tested. Here, we investigate variation in the manus morphology of four modern tapir species (Tapirus indicus, T. bairdii, T. pinchaque and T. terrestris) using a geometric morphometric approach. Autopodial bones were laser scanned to capture surface shape and morphology was quantified using 3D landmark analysis. Landmarks were aligned using Generalised Procrustes Analysis, with discriminant function and partial least square analyses performed on aligned coordinate data to identify features that significantly separate tapir species. Overall, our results support the previously held hypothesis that *T. indicus* is morphologically separate from neotropical tapirs; however, previous conclusions regarding function from morphological differences are shown to require reassessment. We find evidence indicating that T. bairdii exhibits reduced reliance on the lateral fifth digit compared to other tapirs. Morphometric assessment of the metacarpophalangeal joint and the morphology of the distal facets of the lunate lend evidence towards not only high loading on the lateral digits of both the large T. indicus (large body mass) and the small, long limbed T. pinchaque (ground impact). Our results support other recent studies on *T. pinchaque*, suggesting subtle but important adaptations to a compliant but inclined habitat. In conclusion, we demonstrate further evidence that the modern tapir forelimb is a variable locomotor unit with a range of interspecific features tailored to habitual and biomechanical needs of each species.

44 Keywords:- discriminant function – locomotion – manus – mesaxonic symmetry – tetradactyl

Introduction

 MacLaren 3

Modern tapirs (Tapiridae; *Tapirus* Brünnich) are enigmatic, forest-dwelling representatives of the order Perissodactyla (odd-toed ungulates) (Steiner & Ryder 2011; Cozzuol et al. 2013; Ruiz-García et al. 2012). In addition to equids (horses, asses and zebras) and rhinoceroses, tapirs represent the last members of a formerly highly speciose order of small to very large herbivores (Norman & Ashley 2000: Janis 1989). The tetradactyl (four-toed) manus of the modern tapir is a unique feature in extant perissodactyls, with equids and rhinoceroses having reduced their functional digit number to one and three respectively (MacFadden 1992); the earliest ancestors of rhinoceroses, tapirs and equids also displayed a tetradacyl manus (Holbrook 2001). The small, basal members of the Perissodactyla (e.g. Propalaeotherium, Hyracotherium, Heptodon) are interpreted as forest-dwelling browsers with a 'primitive' digital condition, bearing three toes on the hind foot (tridactyly) and four on the forefoot (Hellmund 2005; Wood et al. 2010; Radinsky 1965; Holbrook 2001). This plesiomorphic characteristic of the tapir manus, among other features of tapir anatomy, has contributed to the traditional interpretation of tapirs as 'living fossils' (Janis 1984; Hershkovitz 1954; Padilla et al. 2010; Schoch 1989). Consequently, extant tapirs have been the object of numerous morphological and ecological comparisons to extinct tetradactyl perissodactyls (including Radinsky 1965; Janis 1984; Holbrook 2001, 2009). However, these studies often treat *Tapirus* either as a single morphological unit (e.g. Holbrook 1999, 2001), or compare only one or two species of *Tapirus* with extinct tetradactyl perissodactyls (e.g. Simpson 1945; Radinsky 1965). Recent studies on the extinct tapirs of North America are beginning to increase species counts when performing comparative analyses, albeit with predominantly qualitative techniques (Hulbert 1995; Hulbert 2005; Hulbert 2010; Hulbert et al. 2009; Holanda et al. 2012). Using Tapirus as a solitary morphological unit is greatly beneficial for phylogenetic comparisons with more basal tapiromorph perissodactyls, e.g. Lophiodon (Holbrook 2009) and Colodon (Colbert 2005), as it does not require exhaustive character comparisons across all species of tapir through time. However, to test evolutionary questions on the

comprehensive understanding of limb variation in potential modern analogues is essential. One such

functional morphology of the postcranial skeleton in basal, tetradactyl perissodactyls, a

question concerns the true axis of symmetry in the mesaxonic autopodium.

Page 4 of 53

MacLaren 4

Perissodactyls, including tetradactyl, tridactyl and monodactyl taxa, possess mesaxonic symmetry in their manus (Klaits 1971); the axis of symmetry passes through the third digit. The term 'mesaxonic' has been used to describe autopodia in a variety of tetrapod groups. Anatomical and morphometric studies determine a mesaxonic autopodium to exhibit a third digit that is longer than all the others, flanked by digits two and four, which are shorter than digit three but of comparable length to one another (Reghem et al. 2012; Lockley 2009; Brown & Yalden 1973; Tougard et al. 2001; Rajkumar & Klein 2014). Other studies approach the subject of mesaxony from a more functional and biomechanical standpoint, suggesting that mesaxonic symmetry is not exclusively defined by longer third digits, but that the central third digit is loaded most greatly during locomotion. Lateral digits are then loaded approximately equally (Brown & Yalden 1973; Klaits 1971; Holbrook 2001), with the third digit acting as the centre of rotation during lift-off of the foot (Klaits 1971). The first, anatomical definition of mesaxonic symmetry has been known to be true for perissodactyls for many years (Earle 1893; Earle 1896; Simpson 1945; Gregory 1929); the second, biomechanical interpretation has yet to be explored in all living perissodactyl groups. Understanding the comparative morphology of the manus in modern tapirs, which are known to exhibit anatomical mesaxonic symmetry, may reveal osteological evidence for variation in load application across the four manual digits that also support the biomechanical interpretation of mesaxonic symmetry. Unfortunately the majority of tapir postcranial research has centred on qualitative descriptions, with little by way of quantitative morphological investigation required for proper functional interpretations.

Journal of Morphology

MacLaren 5

Previous qualitative studies of modern tapir postcranial morphology have revealed interspecific differences, almost exclusively between the lowland tapir (T. terrestris L.) and the Malayan tapir (T. *indicus* Desmarest) (Earle 1893; Gregory 1929). Results often align, with T. *indicus* shown to possess longer, heavier and more graviportally adapted limb elements compared to T. terrestris in all analyses (Earle 1893; Osborn 1929; Hulbert 1995). In addition, T. terrestris has been stated to have a smaller lateral toe (fifth metacarpal) relative to tapirs of greater body size, e.g. T. indicus, T. havsii Leidy (Earle 1893; Osborn 1929; Hulbert 1995). When interpreted functionally, the graviportal adaptations of the upper arm, carpus and the metacarpals in T. indicus have been suggested to imply greater loading on the forelimb, and in turn greater reliance on the lateral digits than the smaller T. terrestris (Earle 1893). Many of these qualitative observations may have functional consequences and also associated changes in surrounding bones which have not yet been quantified.

Quantitative comparisons of tapir postcrania have recently been undertaken, with results suggesting that differences in both forelimb (MacLaren and Nauwelaerts 2016; Nauwelaerts et al. 2016) and hindlimb (Hawkins 2011) pertain to subtle variations in locomotor ecology across extant tapir species. These quantitative studies corroborate qualitative observations on the large Malayan tapir (T, T)*indicus*), demonstrating that this species exhibits subtle adaptations to the upper forelimb bones consistent with increased necessity for gravitational support (MacLaren and Nauwelaerts 2016). The mountain tapir (Tapirus pinchaque Roulin) has been shown to possess more gracile upper forelimb and lower hindlimb bones (MacLaren & Nauwelaerts 2016; Hawkins 2011), and morphological features pertaining to proximal shock absorption and increased stride frequency (MacLaren and Nauwelaerts 2016). The upper forelimb morphologies of the Baird's (T. bairdii Gill) and lowland (T. *terrestris*) tapirs have been shown to differ significantly from both *T. pinchaque* and *T. indicus*,

```
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
```

MacLaren 6

despite presenting only subtle osteological differences from one another (corroborating qualitative
observations of these species by Simpson 1945; MacLaren and Nauwelaerts 2016). From the results
of the few quantitative studies on tapir limbs that have been performed, ecological conclusions have
been drawn (Hawkins 2011; MacLaren & Nauwelaerts 2016; Nauwelaerts et al. 2016). Here, we
present a quantitative assessment of the autopodium of extant tapirs to further our understanding of
interspecific differences in the locomotor apparatus of modern tapirs.

Using results and interpretations from qualitative studies on the perissodactyl carpus and metacarpus, combined with recent quantitative results on tapir postcranial anatomy (Hawkins 2011; MacLaren & Nauwelaerts 2016), we will investigate several hypotheses concerning tapir autopodial variation. Firstly, we will quantitatively test the hypotheses presented by Earle (1893), Osborn (1929) and Simpson (1945), detailing differences in the morphology of the carpals and metacarpals between T. *indicus* and *T. terrestris*. Furthermore, due to its larger average body dimensions and mass (de Thoisy et al. 2014), we hypothesise that *T. indicus* will display shape differences in keeping with greater loading on the autopodium compared to all other extant species across all autopodial bones. Recent work on the limb morphology of modern tapirs has shown distinct morphological differences between the mountain tapir (T. pinchaque) and other neotropical tapirs (Hawkins 2011; MacLaren & Nauwelaerts 2016); in keeping with these results, we predict significant differences in the autopodial anatomy of *T. pinchaque* relative to other neotropical tapirs. Finally, we hypothesise that mean average carpal and metacarpal shapes for T. terrestris and T. bairdii will not show significant differences, based on results from both qualitative (Simpson 1945) and quantitative (MacLaren and Nauwelaerts 2016) studies. By testing these hypotheses, we aim to shed light on potential differences in the mesaxonic manus of modern tapirs (Klaits 1971), and infer biomechanical outcomes based on any variation revealed. We will use discriminant function analyses to identify features of the

Journal of Morphology

MacLaren 7

autopodium that contribute to accurate discrimination between species, and aim to formulate functional interpretations from these discriminant features. Material and Methods **Specimens** As study material, 22 disarticulated forelimbs (dry bones) of tapirs were collected from museums in Europe and the USA (see Supplementary Information 1 & 2). Four species of modern tapir (Tapirus terrestris, T. pinchaque, T. bairdii and T. indicus) were collected for analysis, with multiple specimens accounting for intraspecific variation. Whenever possible, morphologically mature specimens were scanned (adult; SI 1), as defined by the complete ossification of the epiphyses, including the scapular cartilage (Liebich et al. 2007; Simpson 1945). Specimens without fully ossified dorsal borders (sub-adult; SI 1) were also included. Sexual dimorphism has been described as non-significant for morphological comparisons in tapirs, and therefore was not considered as a limiting factor for specimens (Simpson 1945). Seven carpals and all four metacarpals were included in the study (Figure 1). Sesamoids and phalanges were not included in this study due to poor sample sizes for these elements. The bones were split into three groupings: the proximal carpal row, distal carpal row, and metacarpals. The proximal row included the pisiform (accesorium), cuneiform (ulnare), lunate (intermedium) and scaphoid (radiale). The distal row included the trapezoid (carpale *II*), magnum (*carpale III*) and unciform (*carpale IV*). The trapezium (*carpale I*) was observed in the juvenile *T. indicus* after dissection and is known to be exhibited in living perissodactyls, although with little consistency (Constantinescu et al. 2012); the trapezium was omitted from this analysis as few scanned specimens possessed it or had the bone preserved for study. All available metacarpals (MC2, MC3, MC4, MC5) were included in the analysis (Figure 1).

MacLaren 8

A dissection was performed on the limbs of a juvenile *Tapirus indicus* to supplement functional interpretations from published tapir osteology and myology (Pereira 2013; Campbell 1936; Murie 1871). The juvenile tapir was provided by the Royal Zoological Society of Antwerp (KMDA). Muscular and ligamentous attachment sites available from the dissection and published literature assisted in the identification of osteological features and interpreting functional outcomes. Veterinary accounts of equid and rhinocerotid osteology and myology (Clayton et al. 2013; Constantinescu et al. 2012; Budras et al. 2003; Liebich et al. 2007; Yalden 1971) were used where necessary to assist identification and interpretations.

174 Scanning

The disarticulated carpus and metacarpals from one forelimb of each specimen were laser scanned using a FARO ScanArm Platinum V2 system with integrated FARO Laser Line Probe (up to 50 µm resolution). Bones were balanced on supports positioned on regions of the specimen surface on which landmarks could not be placed (e.g. shaft of metacarpal). A three-dimensional virtual point cloud was produced for each autopodial bone, visualised in GeoMagic (GeoMagic Qualify v.10, Morrisville, NY, USA). Any outlying surfaces in the point clouds (e.g. incidental scanning of support structures) were digitally removed to focus only on surface information from the bones. Point clouds were then converted into polygon-based surface models, ranging in detail from 200k to 500k polygons, dependent upon bone and the detail necessary around articular surfaces.

Geometric Morphometrics

Landmark-based geometric morphometrics has been extensively used and is an appropriate
technique for quantifying differences in shape between three-dimensional objects (Rohlf & Slice
1990; Gould 2014). The technique is based on series of discrete, biologically or operationally

Page 9 of 53

Journal of Morphology

MacLaren 9

3 4	189	homologous points (landmarks) placed onto a succession of objects (Zelditch et al. 2012). Type II
5 6	190	landmark points (representing maxima and minima) were used in this study to define the shape of the
7 8	191	carpals and metacarpals. Landmark placement on representative bones in this analysis are visualised
9 10	192	in Figure 2 (carpals) and Figure 3 (metacarpals). To aid in the description of discriminant features,
11 12 12	193	landmarks were annotated with subscript denominations pertaining to the bone the landmark
13 14 15	194	describes (as in MacLaren and Nauwelaerts 2016) (defined in Supplementary Information 3: Table
16 17	195	S1). Surface models were imported into Landmark Editor v.3.0 software (Wiley 2006) for three-
18 19	196	dimensional landmark application. Raw landmark coordinates were then exported into MorphoJ
20 21 22	197	v1.06d (Klingenberg 2011) and aligned using Generalised Procrustes Analysis (GPA). This
22 23 24	198	technique removed the effect of size, location and orientation and aligned raw coordinate
25 26	199	configurations based on geometric centre (centroid), minimising inter-landmark distance (Rohlf &
27 28	200	Slice 1990; Adams et al. 2004; Zelditch et al. 2012). Resultant Procrustes coordinates and centroid
29 30	201	sizes were then exported into SPSS v.23 (IBM Corp. 2013) for discriminant analyses and post-hoc
31 32 33	202	testing. Centroid size represents an intrinsic size measure that can be used to scale a configuration of
34 35	203	landmarks, for example to assess metric distances between landmarks. Centroid sizes for adult
36 37	204	specimens were retained for size comparisons. A multivariate analysis of variance (MANOVA) was
38 39	205	performed on the Procrustes coordinates calculated for each bone. MANOVA was used to test
40 41 42	206	differences in the means of the groups (species), and the observed power of our MANOVA using
42 43 44	207	small sample sizes was retrieved from the analysis. The MANOVA and power analysis was
45 46	208	performed in SPSS v.23.
47 48 49	209	
50 51 52	210	Discriminant Function Analysis
53 54 55	211	A linear discriminant function analysis (DFA) was performed on the Procrustes coordinates (x, y, z)
55 56 57 58	212	for all bones. DFA was used to determine what combination of continuous variables for each bone

best discriminated between the four species. DFAs were performed in SPSS v.23 (IBM Corp. 2013) using a forward step-wise method for Procrustes coordinate input; this removed independent variables that were not significant for discrimination. A classification table was produced by predicting group membership and cross-validating by jack-knifing the dataset. Sensitivity and specificity tests were also performed. To assess differences in group means, we employed the Wilk's lambda test (0 - 1; 0 = highest likelihood of inequality, 1 = high likelihood of group means being equal). For visualisation of results, linear discriminant function plots were produced based on the first two discriminant functions (DF1 & DF2), which accounted for the highest percentage of variance. Territorial maps were added to demonstrate how groups were divided and where cut-off values were placed dependent on DF1 and DF2 scores. The third discriminant function accounted for between 0.3% and 11.6% of total variance; DF3 is reported in the results, but is not plotted in discriminant function plots. Cut-off values between groups were based on the weighted mean of the discriminant score for each group centroid. Classification tables and territorial maps were produced in SPSS v.23 (IBM Corp. 2013), and resultant discriminant function plots were configured in R Studio (R Core Development Team 2008).

229 Partial Least Squares (PLS) analysis

Within the carpal complex there are a variety of bones with multiple facets interacting with one
another. Changes in morphology in one joint facet which may signal a shift in mechanical
capabilities should correspond to similar changes in adjoining bones. In order to assess whether
bones and joint facets covary in morphology with neighbouring carpals, we utilised a two-block
partial least square analysis (2B-PLS) accompanied by a permutation test (10000 repetitions) to test
for significance of covariance (Rohlf & Corti 2000; Fadda & Corti 2001). The RV coefficient of
integration (multivariate generalization of squared Pearson correlation coefficient; Klingenberg

Journal of Morphology

MacLaren 11

2009) produced by PLS analyses is used to predict degree of covariation between two blocks of data (in this case, landmark coordinates and facet areas); it is measured between 0 (no covariation) and 1 (complete covariation)(Klingenberg 2009). This analysis was used for adjoining carpals which demonstrated key features which both discriminated between species and pertained to possible biomechanical differences during locomotion. When area measurements were tested for covariance (as opposed to landmark coordinates) the raw area data were log-transformed prior to 2B-PLS analysis. The PLS analyses were performed in MorphoJ v1.06d (Klingenberg 2011), with graphical representations compiled in R Studio (R Core Development Team 2008).

Joint Facet Comparisons

To compliment carpal shape differences detected using 3D landmark analysis, the relative areas of joint surfaces were also calculated. Variation in joint surfaces (facets) have been reported in tapirs through qualitative comparisons (Simpson 1945; Earle 1893; Osborn 1929); here we used landmark analyses to detect differences in the shape of facets using only the landmarks that define the joint facet in question. In addition, we calculated relative areas of a series of joint surfaces of the scaphoid, lunate and unciform to quantify previous qualitative claims about interactions between carpals within the autopodium (Osborn 1929; Earle 1893; Simpson 1945). We identified and tested two inter-carpal facet relationships: the distal lunate facet ratio (highlighted by Earle 1893) and the unciformmagnum facet ratio (again pertaining to the lunate, Osborn 1929; Simpson 1945). The distal lunate possesses two large facets: anteriorly the unciform facet and posteriorly the volar magnum facet. The anterior magnum facet is found alongside the unciform facet. In the densely packed carpus, a relatively larger facet intuitively implies greater loading rather than greater mobility, with a larger surface area available for force transmission. The three-dimensional scans of the carpal bones provided smooth articular surfaces for quantitative comparison between carpals. Facet areas were

MacLaren 12

2
3
4
5
6
0
1
8
9
10
11
12
13
14
14
15
16
17
18
19
20
21
22
22
23
24
25
26
27
28
20
20
30
31
32
33
34
35
36
37
38
20
39
40
41
42
43
44
45
46
17
יד 10
40
49
50
51
52
53
54
55
56
50
ວ/ 50
58
59
60

1

261	calculated by pruning the full 3D laser scans of bones until only the joint facet under study remained;
262	this was performed in GeoMagic (GeoMagic Qualify v.10, Morrisville, NY, USA). These reduced
263	surface scans were then imported into MeshLab (Cignoni et al. 2008) to calculate surface areas.
264	Ratios were formulated by dividing the posterior or anterior magnum facet area (whichever was
265	appropriate) by the combined total of the unciform joint facet and the respective magnum facets. To
266	test for covariation between the joint facets of the distal lunate (both between anterior unciform-
267	magnum and posterior unciform-magnum), a two-block partial least square analysis (2B-PLS) was
268	performed on the log-transformed area data. 2B-PLS analyses and 10000 permutations were
269	performed in MorphoJ v1.06d (Klingenberg 2011).
270	

271 Distal Metacarpophalangeal Facet Variation

The distal metacarpal shape was further investigated using a subset of landmarks to test for interspecific differences exclusively in the distal joint surface. Eight landmarks, homologous for all four metacarpals across tapir species, were selected, describing the palmar distal joint facet (metacarpophalangeal joint). In order to test for differences concurrently between both metacarpals and species, a Procrustes ANOVA was performed in MorphoJ. This analysis was used to complement and inform functional interpretations of morphological changes in the distal metacarpals pertaining to interactions with the proximal phalanges (pastern) and the proximal sesamoids.

279

280 Results

Overall, linear discriminant functions successfully discriminated between the four extant species of tapir for all autopodial bones. Jack-knifed classification tables for all bones in the autopodium are presented in Table 1, reporting sensitivity and specificity of the analyses. Accuracy of jack-knifed species classification for autopodials exceeded 75% accuracy for all cases. *T. indicus* is classified

Journal of Morphology

MacLaren 13

100% accurately for all carpal bones, whereas T. terrestris is the most frequently misclassified species (six different bones). T. bairdii is the most accurately classified neotropical tapir, with only the cuneiform demonstrating inaccuracy in classification (Table 1). Power analyses revealed high statistical power for all MANOVAs (mean power = 0.88 ± 0.08); full tabulated results can be found in the Supplementary Information 4: Table S2. We are therefore confident in the power of this analysis and the morphological differences between the taxa. Here, we describe results of discriminant function analysis for all autopodials (proximal carpal row, distal carpal row, and metacarpals). Descriptions of landmarks affecting discrimination can be found in the Supplementary Information.

295 Proximal Carpal Row

The proximal carpal row (scaphoid, lunate, cuneiform and pisiform; Figure 1) contains the bones that interface with the radius and ulna; the scaphoid, lunate and cuneiform also articulate with the distal carpal row. Interspecific classification in the proximal carpal row ranges from 100% accuracy (scaphoid) to 75% accuracy (cuneiform) (Figure 4), with bones that articulate with the radius (scaphoid and lunate) showing more accurate classification than those articulating with the ulna (cuneiform and pisiform) (Table 1). Here we present results for discriminant function analysis on the bones of the proximal carpal row:

Scaphoid

The scaphoid is the largest carpal of the tapir proximal carpal row, and articulates proximally with the radius, medially with the lunate, and distally with the magnum, trapezoid and in some cases the trapezium. The first two discriminant functions (DF1 and DF2) based on scaphoid landmarks account for 97.9% of variance (Figure S1a). The features that most greatly influence accurate species classification include the anteroposterior morphology of the palmar lunate facet (scLm 20), and the

MacLaren 14

upper margin of the trapezoid-magnum facet, defined by _{sc}Lm 11. *T. bairdii* shows the greatest distinction in scaphoid morphology from other tapirs. In *T. bairdii*, _{sc}Lm 11 is placed more distally and _{sc}Lm 20 is more posterior than in other species. Average species centroid sizes for the scaphoid

show that *T. indicus* possess the largest scaphoid, with *T. pinchaque* displaying the smallest (Table
2).

Lunate

The lunate, or semi-lunar, represents the central carpal in the proximal carpal row. The proximal surface articulates with the radius, medially and laterally it articulates with the scaphoid and cuneiform respectively. Distally the lunate has three articular facets: one to the unciform and two to the magnum (one dorsal, one palmar). Discriminant function 1 accounts for 93.1% of variation (Figure S1b); interspecific discrimination along DF1 is most greatly influenced by placement of _{lu}Lm6, the edge of the palmar magnum facet closest to the dorsal facet. Dorsal deflection and elongation of this facet in T. indicus brings the distal lunate facets into closer proximity to each other. Proximodistal expansion of the entire dorsal surface of the lunate is observed along DF2 between neotropical taxa (Figure S1b), driven by placement of _{lu}Lms 3 & 18. Average centroid sizes differ to those of the scaphoid, revealing T. indicus to possess the largest lunates and T. terrestris the smallest (Table 2).

Cuneiform

The cuneiform, triquetrum or ulnar carpal, is the most lateral bone in the proximal carpal row; it articulates proximally with the pisiform and ulna, medially with the lunate, and distally with the unciform (Figure 1). The cuneiform is the most poorly discriminated bone in the autopodium, with one in four bones being misclassified (Figure 4). The first two discriminant functions describe 92.4% of total variance (Figure S1c). One or more specimens of all neotropical species are misclassified as *T. indicus*, with additional misclassification between *T. bairdii* and *T. terrestris*; the first discriminant

Journal of Morphology

MacLaren 15

function successfully separates only *T. pinchaque* from the other species (Figure S1c; see also
Supplementary Information 5: Table S3). The most discriminatory feature is the shape of the
mediodistal facet articulating with the lunate (_{py}Lm 3); the orientation of the cuneiform (defined by
_{py}Lm 3 and 4) also contributes to successful discrimination of *T. pinchaque*. Average centroid sizes
for the cuneiform show a similar pattern to that of the scaphoid, with *T. indicus* displaying the largest
and *T. pinchaque* the smallest cuneiform carpals (Table 2).

339 Pisiform

The pisiform, or accessory carpal, is the most palmar bone in the carpus and facilitates the passage of flexor tendons through the carpal tunnel. The pisiform articulates distally with the cuneiform and proximally via two facets with the ulna. The first two discriminant functions describe 89.1% of variance, with each species occupying a discrete region of canonical variate-space (Figure S1d). Placement of _{pi}Lm 4 (distal extremity of ulnar facet) shows variation along DF1, with *T. indicus* displaying a distinct morphology from T. bairdii. DF2 discriminates between T. terrestris and T. *pinchaque* with the placement of _{pi}Lm12 (accessory ulnar facet) discriminating between these two taxa. As with the scaphoid and cuneiform, T. indicus demonstrates the largest centroid sizes for the pisiform, and *T. pinchaque* the smallest centroid size (Table 2).

350 Distal Carpal Row

The distal carpal row (trapezoid, magnum, unciform) is the most successfully classified group of autopodials (mean classification accuracy = 98.4%). Within this grouping, both the trapezoid and unciform achieved 100% interspecific classification, whereas the magnum was classified with 95.2% accuracy. Graphical results are presented in Figure S2.

355 Trapezoid

MacLaren 16

articulation with the scaphoid, a lateral facet for the magnum, a distal facet for the second metacarpal (MC2) and a small medial facet for articulation with the trapezium (first carpal; not present in all specimens and therefore not included in the analysis). The first discriminant function accounts for 93.6% of interspecific variation, and successfully discriminates the four taxa (Figure S2a). Separation along this function is influenced by the landmarks describing the palmar region ($_{tr}$ Lm 7) and the laterodistal margin (trLm 10) of the of the magnum facet. The difference in the magnum facet morphology is greatest between T. bairdii and T. indicus. The centroid size for the trapezoid mirrors that of the scaphoid, cuneiform and pisiform (Table 2).

The trapezoid, or second carpal, is the smallest carpal in the tapir autopodium. It has a proximal

365 Magnum

The magnum, capitate or third carpal, is the central carpal of the tapir autopodium. The magnum articulated proximally with the scaphoid and lunate (via two facets), proximolaterally with the unciform, medially with the trapezoid and distally with a small facet for the second metacarpal (MC2) and a large facet for the third metacarpal (MC3). The first discriminant function accounts for 95.8% of magnum variation (Figure S2b). T. indicus and T. bairdii are the most easily discriminated taxa along DF1; however, specimens of *T. pinchaque* and *T. terrestris* are misclassified along DF1. Landmarks that most heavily influence interspecific discrimination along DF1 include maLm 7 (expansion-contraction of the unciform facet) and maLm 15 (defining the concavity of the dorsal trapezoid facet). T. bairdii exhibits a highly concave trapezoid joint plane. As with the majority of the carpals (excluding the lunate), T. indicus displays the largest average magnum centroid size and *T. pinchaque* exhibits the smallest (Table 2).

377 Unciform

The unciform, unciform or fourth carpal, is the largest carpal in the distal carpal row; it articulates proximally with the lunate and unciform, medially with the magnum, and distally with the third,

Journal of Morphology

MacLaren 17

fourth and fifth metacarpals (MC3, MC4 & MC5). The first two discriminant functions describe 88.4% of total variance (Figure S2c). The placement of taxa along DF1 is greatly affected by haLm 3 (lateral morphology of the MC5 joint facet) and haLm 10 (anteroposterior expansion or constriction of the lunate facet). T. indicus displays an expanded lunate-cuneiform facet relative to neotropical taxa. DF2 (32.9%) is most greatly influenced by haLm 8, which tracks a relative expansion of the medial edge of the cuneiform facet. DF2 is also influenced by $_{ha}Lm 5$, which describes the antero-posterior constriction (T. bairdii) and expansion (T. indicus and T. pinchaque) of the MC4 facet, and by extension the entire distal unciform. The average centroid size is once again greatest in T. indicus and smallest in T. pinchaque (Table 2).

390 Metacarpals

Overall results for the metacarpals suggest that the lateral bones exhibit marginally less interspecific variation than the medial metacarpals. Mcs 2 and 3 were classified 100% accurately after jackknifing, whereas Mcs 4 and Mc 5 exhibited occasional misclassification. Centroid sizes for the metacarpals do not follow the same pattern as in the carpals.

Metacarpal 2

Metacarpal 2 (MC2), or the second metacarpal, is the most medial hand bone in the tapir
autopodium. It has a proximal articulation with the trapezoid and trapezium (absent in this analysis),
a lateral articulation with MC3, and a distal articular facet for the proximal phalange and proximal
sesamoids. DF1 accounts for 99.8% or variation, with *T. indicus* greatly separated from the
neotropical taxa (Figure S3a). Separation along DF1 is heavily influenced by the placement of 2Lm
26 (palmar margin of trapezoid facet), and also 2Lms 4 & 5, which describe the morphology of the
proximolateral sesamoid joint facet, in addition to affecting the dorsopalmar depth of the metacarpal

MacLaren 18

head. The largest average centroid size for the MC2 is found in *T. indicus*, with the smallest

404 exhibited by *T. terrestris*. *T. pinchaque* exhibits the second largest MC2 centroid size (Table 2).

Metacarpal 3

Metacarpal 3 (MC3), or the third metacarpal (cannon bone in equids) is the largest metacarpal in the tapir autopodium. Proximally it articulates with the trapezoid, magnum and unciform, proximomedially with MC2 and proximolaterally with MC4; MC3 articulates distally with the proximal sesamoids and phalange for digit three. The first two discriminant functions account for 92.0% of variation (Figure S3b). Landmarks that contribute most greatly to interspecific classification along DF1 include ₃Lm 19 (proximodistal depth of the magnum facet) and ₃Lm 23 (palmar edge of the MC2 joint facet). Classification along DF2 is dominated by ₃Lm 17 (describing the breadth of the unciform joint facet). Average centroid size for MC3 suggests that T. indicus have the largest metacarpal; both T. terrestris and T. bairdii display very similar average centroid sizes, smaller than the other two species (Table 2).

Metacarpal 4

Metacarpal 4 (MC4), or the fourth metacarpal, is the intermediate metacarpal between the central third and lateral fifth. MC4 articulates with the unciform proximally, MC3 proximomedially and MC5 proximolaterally; as with other metacarpals, MC4 distally articulates with the corresponding proximal phalange and paired sesamoids. MC4 is accurately classified for 90.9% of specimens (Table 1), with the first discriminant function accounting for 94.8% of variation (Figure S3c). The landmarks which contribute most greatly toward interspecific discrimination describe the concave shape of the unciform facet ($_{4}$ Lm 15) and the expansion-contraction of the medial margin of the metacarpophalangeal joint facet (4Lm 8). The MC4s of the neotropical tapirs show very similar average centroid sizes, with *T. terrestris* marginally displaying the smallest (Table 2).

426 Metacarpal 5

Journal of Morphology

MacLaren 19

The fifth metacarpal (MC5) is the most lateral hand bone, and the smallest metacarpal in the tapir autopodium. Proximally MC5 articulates with the unciform, proximomedially with MC4, and distally with the proximal phalange and paired sesamoids of digit five. The first two discriminant functions account for 94.5% of interspecific variation. Along DF1, three morphotypes are separated (Figure S3d). Landmarks that display high loading on DF1 include those describing the morphology of the lateral sesamoid facet ($_5Lms 3 \& 5$), which divides modern tapirs into three morphotypes (Figure S3d). T. bairdii displays a notably smaller average centroid size than other neotropical taxa, with T. indicus exhibiting the largest average centroid size for MC5 (Table 2).

435 Metacarpophalangeal Facet

The metacarpophalangeal joint of the tapir metacarpal comprises of three principal regions: the medial sesamoid facet, lateral sesamoid facet and metacarpal sagittal ridge. The subset of eight landmarks describe the proximopalmar margin of the metacarpophalangeal joint, incorporating the sesamoid facets and sagittal ridge. Results for the subset of eight landmarks for all four metacarpals demonstrated notable interspecific differences. Procrustes ANOVA results detected significant differences (p < 0.01) between individual species and between the four metacarpals in the morphology of the distopalmar metacarpal facet (Table 3).

444 Joint Facet Ratios

445 Unciform–Magnum Facet Ratio (HMF Ratio)

The unciform-magnum facet ratios show a different pattern to that of the distal lunate facets. *T. pinchaque* displays the largest average anterior magnum facet, and shows a significant difference to *T. indicus* in the ratio of unciform to anterior magnum facets (p = 0.014; Table 4). *T. indicus* displays
the greatest range of ratios, with one outlying specimen exhibiting a ratio comparable to *T. pinchaque* (Figure 6). *T. indicus* displays no significant difference to *T. terrestris*, and *T. bairdii* and

MacLaren 20

T. terrestris display similar HMF ratios. Covariation analyses based on 2B-PLS analysis of the 452 unciform and anterior magnum facet areas do not support covariation between these facets. PLS1 453 axes account for 100% of covariation; however, PLS axes do not correlate highly (RV = 0.363) 454 (Figure 5a; Table 6). Overall correlation is weak and not statistically significant after permutation 455 (RV = 0.132; p = 0.117).

457 Distal Lunate Facet Ratio (DLF Ratio)

The comparison between the distal lunate facets (DLFs) show that there is a spectrum of variation across neotropical species (Figure 7). T. bairdii demonstrates the greatest difference between anterior and posterior distal facets (Figure 7), showing a significant difference to *T. terrestris* (Table 5) which exhibits the smallest difference between facet areas. T. indicus and T. pinchaque demonstrate near identical mean values for distal facet area ratios (T. indicus: 0.663 ± 0.036 ; T. pinchaque: $0.663 \pm$ 0.031). Covariation analyses based on 2B-PLS analysis of the unciform and posterior magnum facet areas support a covariation relationship between these facets. Again, PLS1 axes account for 100% of covariation, with a strong positive co-relationship between PLS axes (RV = 0.738) (Figure 5b; Table 6). Overall correlation is fairly strong (RV = 0.545) and statistical significance from the permutation test is very high (p < 0.001).

469 Partial Least Squares analyses (2B-PLS)

470 Results from discriminant function analyses suggest bones along the medial autopodium (scaphoid,
471 trapezoid and MC2) are most accurately discriminated across all tapir species (Figure 4). To
472 investigate specific articulations in the medial autopodium, 2B-PLS was performed between the
473 trapezoid and magnum (examining the joint facet between the two bones) and the respective facet
474 morphologies of the trapezoid and scaphoid. Overall 2B-PLS analyses between the trapezoid and

Journal of Morphology

MacLaren 21

magnum revealed a strong covariation in joint facet morphology (RV = 0.778), with high statistical significance from permutation test (p < 0.001). The fist PLS axes account for over 80% of covariance between the bones (Figure 5c; Table 6), which is also highly significant following permutation testing (p < 0.001). Coordinates which most greatly influence covariation for PLS1 include trLm8 (anterior concave edge of magnum facet), maLm14 and maLm15 (anterior and posterior concave margins of trapezoid facet).

The overall 2B-PLS analyses between the trapezoid and scaphoid shows modest covariation in joint facet shape (RV = 0.415), albeit with no statistical significance after permutation test (p = 0.089). The fist PLS axes account for over 60% of covariance between the bones (Figure 5d; Table 6), which does exhibit high statistical significance with permutation testing (p = 0.005). Procrustes coordinates which most greatly influence covariation for PLS1 include trLm3, 4 and 6 (proximal extremities of both anterior and posterior margins), and scLm11 and scLm13 (anterior margin of trapezoid facet and deepest point on the concave facet for the trapezoid).

489 Discussion

Variation in the carpal and metacarpal arrangement within Perissodactyla has been studied with various qualitative techniques, with both morphological and functional conclusions being drawn at the genus level (Tapirus) (Klaits 1972; Osborn 1929; Earle 1893; Simpson 1945; Holbrook 2001). However, the comparative morphology and interspecific variation within the manus of the genus *Tapirus* has only briefly been touched upon in previous studies (Earle 1893; Simpson 1945; Osborn 1929), and has not taken all extant taxa into account. Interspecific variation in tapir autopodials may reflect subtle variation in locomotor style, and possibly variation in application of loading forces on the anatomically mesaxonic manus of tapirs. Here we discuss the major osteological differences in the autopodium of extant *Tapirus*, and their implications for locomotor variability in this group.

MacLaren 22

499	
500	Facets of the lunate
501	Throughout previous comparisons between tapir postcrania, several key differences in the
502	autopodium have been postulated. In particular, clear differences between T. indicus and other
503	modern tapirs have been suggested (Earle 1893; Osborn 1929). We find strong support for this
504	distinction between T. indicus and other modern tapirs. However, our findings do not correlate with
505	the specific conclusions from previous qualitative studies (Earle 1893; Osborn 1929). For example,
506	Earle noted little to no contact between the lunate and magnum (Figure 1) in <i>T. terrestris</i> when
507	compared to <i>T. indicus</i> , and that the approximately equal facets for unciform and magnum in <i>T</i> .
508	indicus allows equal transmission of force to the medial and lateral digits (Earle 1893).
509	Our investigation reveals that the lunate contact with the magnum in <i>T. indicus</i> possesses the
510	smallest facet (on average) relative to the unciform joint (Figure 6), which is in direct contrast to the
511	findings of Earle (1893). A relatively larger unciform facet on the lunate would conceivably enable
512	greater force transmission to the unciform and the digits beneath it (the lateral digits) in <i>T. indicus</i> .
513	As such, our results for the lunate facets suggest that <i>T. indicus</i> may not exhibit biomechanical
514	mesaxonic symmetry, in favour of increased loading on lateral digits. Additionally, results for T.
515	terrestris suggest no significant difference to T. indicus in the distal lunate facets (Figure 6; Table 3),
516	which also contrasts with Earle's findings. Finally, we found no statistically significant support for
517	covariation between the areas of these facets across the four tapir species. Individual variation in
518	facet size may be a key factor here, as demonstrated by the large error bars for this ratio in T. indicus
519	(Figure 6). These findings lead us to conclude that, contrary to the deductions of Earle (1893), tapirs
520	with an enlarged unciform facet will not necessarily display reduction in their anterior magnum
521	facet.

Journal of Morphology

MacLaren 23

Our study suggests that T. pinchaque exhibits the largest anterior magnum facet of the lunate (Figure 6), which combined with a large unciform facet enables a more even spread of loading forces to the anterior carpal row and both MC3 and MC4. Although we find no statistical evidence that there is a strong correlation between these facets in our sample, a morphological similarity to extinct tetradactyl perissodactyls is nevertheless present. The carpal arrangement is reminiscent of early, functionally tetradactyl perissodactyls (e.g. Lophiodon & Hyrachyus) (Osborn 1929), and supports quantitative results from scapulo-humeral morphology suggesting T. pinchaque displays a number of osteological features in common with Eocene perissodactyls (MacLaren & Nauwelaerts 2016). It should also be emphasised that our results for the distal lunate facets and anterior magnum-unciform ratios suggest only very small differences in overall area (~10% between largest and smallest). However, we did find significant covariation in the distal lunate facets, suggesting that the lunate articulation with the posterior magnum is linked to changes in area of unciform facet and vice versa. We believe that extrapolating differences in loading regime and further functional outcomes from these small differences would involve over-interpretation of the data. We also stress that the morphological conclusions from Earle (1893) and Osborn (1929) remain on the whole accurate, although their functional interpretations require rigorous re-examination (as recommended by Klaits 1971) with modern quantitative kinematic methods before any solid conclusions on locomotor function can be made.

Mobility of the pisiform

The accessory carpal (pisiform) of tapirs is flattened dorsopalmarly and curves inwards toward the medial border of the autopodium. The curvature of the pisiform enables the passage of the flexor tendons of the *m. flexor digitorum sublimis* + *profundus* through the carpal tunnel (Campbell 1936; Murie 1871; Bressou 1961), and the spatulate tip of the bone is the site for attachment of the *m*.

MacLaren 24

flexor carpi ulnaris (proximal) and *m. abductor digiti minimi* (distal). A recent quantitative analysis revealed two different morphologies for the pisiform facet of the ulna in Tapirus (MacLaren & Nauwelaerts 2016); T. terrestris and T. indicus demonstrated mediolaterally broad pisiform facets on the posterior ulna, whereas T. bairdii and T. pinchaque exhibited more proximodistally elongate facets. Results from the present analysis of the pisiform (accessory carpal) suggest a similar pattern of morphological disparity, especially between T. terrestris and T. pinchaque (Figure 8), further corroborating previous analyses revealing differences in forearm osteology between these closely related taxa (MacLaren & Nauwelaerts 2016). In T. terrestris, the pisiform facet is sub-rhomboidal with approximately parallel edges, and the articulating facet of the ulna is semi-circular in lateral view (Figure 8a; also MacLaren & Nauwelaerts 2016). This offers the pisiform of T. terrestris a relatively smaller surface with which to articulate compared to the other neotropical tapirs, while concurrently allowing a greater range of mobility for the pisiform during carpus flexion. The flatter, more elongate pisiform facet for the ulna may limit the functional capabilities of the lateral autopodium in T. pinchaque and T. bairdii, whereas T. terrestris does not appear to be under such mechanical constraints. In addition, the insertion area for the *m. flexor carpi ulnaris* on the proximoposterior edge of the pisiform (Figure 8i) is accentuated in neotropical taxa (most greatly so in T. bairdii), whereas T. indicus shows no great proximal expansion. The prominent insertion point in neotropical taxa offers a greater surface area for tendon attachment, suggesting increased resistance to carpal over-extension (by the antagonistic *m. flexor carpi ulnaris*). By contrast, the broader distal edge of the pisiform in T. indicus offers greater attachment surface for the m. abductor digiti minimi (abductor of the fifth digit) (Campbell 1936; Murie 1871). In addition to an enlarged attachment site for the *m. adductor digiti minimi* on the volar process of the magnum, this morphological feature implies that T. indicus has greater muscular control over the fifth digit, allowing it to splay the toes to support greater mass on soft substrates. This result supports previous claims that T. indicus utilises the fifth digit to a greater extent than its living neotropical relatives

Journal of Morphology

MacLaren 25

(Earle 1893; Gregory 1929). To assess whether this morphology is common to all large tapir species. similar analyses on extinct taxa of high estimated body mass (e.g. T. haysii, T. augustus) will be necessary. Medial digit loading Typically, the lateral and medial digits within a mesaxonic autopodium will be loaded approximately equally (Klaits 1971; Holbrook 2001). The lateral digits and corresponding unciform (fourth carpal) have been shown to display morphological differences in modern tapirs; therefore, corresponding morphologies in the trapezoid (second carpal) may be expected. One of the most discriminatory features of the trapezoid was the morphology of the joint facet with the magnum (third carpal). The corresponding facet on the magnum was also highly discriminatory (visualised in Figure S2b). For simplicity, we discuss the articulation from here as the trapezoid-magnum facet. The anterior border of the facet is highly concave in T. bairdii, affording relatively less surface for articulation between the bones in this species (Figure 9a). By contrast, T. indicus displays a much less concave anterior or posterior border to the facet, enlarging the relative area of the facet (Figure 9b). This morphology is mirrored in the trapezoid, and the relationship is strongly supported with results from PLS analyses for covariance. Landmarks defining the concave margins of the facet on both trapezoid and magnum contribute most greatly toward the high covariation coefficient. The high covariance between these bones implies a tightly associated morphological relationship between trapezoid and magnum. In the larger T. indicus, the less concave margins and relatively greater articular surface suggest greater immobility across this joint. In addition, we find that the scaphoid facet for the trapezoid is more concave, thus allowing less mobility for the trapezoid within the T. indicus carpus; this finding should be treated with some caution, as this feature was not revealed to be statistically significant after covariation analysis (p = 0.08). Despite the poor co-variation between the trapezoid and

MacLaren 26

scaphoid, evidence from the trapezoid-magnum facet implies the morphology of T. indicus is adapted for greater loading on the medial digit than other modern tapirs, allowing greater force transmission through the medial carpus. This conclusion is further supported by results for metacarpal morphology in this study (see below) and conclusions from previous qualitative assessments (Gregory 1929; Earle 1893). When combined with other subtle differences in the carpal complex of *T. indicus*, the adaptation of the trapezoid-magnum joint suggests the medial manus of *T. indicus* may be loaded more heavily relative to other tapirs, despite maintaining anatomical mesaxonic symmetry. By contrast, T. bairdii displays the least concave trapezoid facet on the scaphoid, with both T. terrestris and T. pinchaque displaying very similar trapezoid-magnum (Figure 9c-d) and trapezoid-scaphoid facet morphologies, intermediate between T. indicus and T. bairdii. This similarity may signify a phylogenetic aspect to this morphological difference (Ruiz-García et al. 2016; Ruiz-García et al. 2012). Further investigation into the carpal morphology of closely related South American taxa (e.g. the extinct *T. cristatellus* and *T. rondoniensis*) may shed more light on the evolutionary history of this morphology.

Metacarpophalangeal facet variation

The tapir metacarpals display anatomical mesaxonic symmetry (axis of symmetry passing through the third digit), both in absolute length and in average centroid size (Table 2). Whereas the central third metacarpal exhibits discriminant variation in the proximal joint surfaces, the second, fourth and fifth metacarpals are most successfully discriminated interspecifically using landmarks describing the palmar metacarpopalangeal joint (Figure 10). The metacarpophalangeal joint includes three main regions (the medial sesamoid facet, lateral sesamoid facet and metacarpal sagittal ridge), all of which are described in part by the landmark analysis. The palmar section of the metacarpophalangeal joint facet articulates with the proximal phalange in addition to the paired sesamoid bones, which slot

Journal of Morphology

MacLaren 27

either side of the sagittal ridge (Liebich et al. 2007; Constantinescu et al. 2012). Quantitative comparisons of this facet across the four metacarpals demonstrated that the facet morphology of each type of metacarpal is significantly different, as are the differences between species (p < 0.001) (Table 3). Two taxa stand out as notably different in their palmar metacarpophalangeal joint morphology: *T. indicus* and *T. pinchaque*.

T. indicus demonstrates a suite of adaptations for increased forelimb loading, as has been shown in previous literature and in this study (Earle 1893; Gregory 1929; Hulbert 1995; MacLaren & Nauwelaerts 2016). The palmar facet of the metacarpals also shows adaptations for increased bone-bone contact, with T. indicus demonstrating a relatively broad facet on all metacarpals (Figure 10), in addition to adaptations to the medial and lateral carpus enabling dissipation of compressive forces (Figure 9). Furthermore, the sagittal ridge of the metacarpals in *T. indicus* is elongated proximally, with mediolaterally broad sesamoid facets, offering large sesamoids a greater surface area with which to articulate. We interpret this as an adaptation contributing to load distribution across each metacarpal, and by extension the entire foot (Easton & Kawcak 2007). Interestingly, this morphology of the palmar sagittal ridge is mirrored in *T. pinchaque*, which is on average the smallest and least massive of the neotropical taxa. As it is unlikely that T. pinchaque would require increased sesamoid-metacarpal contact to overcome high loading due to increased mass (i.e. graviportalism), we hypothesise that this shift in morphology in *T. pinchaque* is consistent with conferring greater stability to each toe (Hildebrand 1985) and spreading the forces more evenly during limb loading (Easton & Kawcak 2007). We also infer that, as this feature is seen in all the metacarpals of T. *pinchaque*, that the distal forelimb of this species has developed increased stability in all its digits. Impact of the fifth digit on the substrate would greatly benefit the animal, especially under potentially high loading conditions such as running up a steep, forested incline. Increased loading and necessity for stability in this comparatively small tapir may be due to a number of factors. Reduced reliance on the digital pad in favour of the toes, as is seen through equid evolution

MacLaren 28

(Thomason 1985; MacFadden 1992), would cause a shift in loading forces to the toes and may account for increased sesamoid facets and necessity for toe stability in *T. pinchaque*. No quantitative comparisons of toepad size has been reported in *T. pinchaque*, and so this interpretation remains speculative until further investigation has been undertaken. In addition, moving up or down sub-alpine habitats and over uneven, high altitude wet-grassland (Padilla et al. 2010; Downer 1996) would necessitate increased digital stability; this supports previous quantitative results on forelimb morphology and biomechanics in this species (MacLaren and Nauwelaerts 2016). We find further support for previous quantitative works (MacLaren & Nauwelaerts 2016; Hawkins 2011) in the overall shape of T. pinchaque metacarpals, which demonstrate a more gracile morphology than those of other extant taxa. Overall, the adaptations observed in *T. pinchaque* in this and other osteological studies hint at the retention or re-exploration of putatively 'primitive' perissodactyl forelimb traits (e.g. equal force distribution from lunate to unciform and magnum; functional fifth digit; gracile long bones), while concurrently developing novel adaptations to both the upper and lower forelimb (e.g. large supraspinous fossa as potential proximal shock absorber; braced resting stance; strongly keeled metacarpophalangeal joints increasing stability for the phalanges during locomotion) (this study; MacLaren and Nauwelaerts 2016). Finally, when compared to the metacarpals of other extant taxa, *T. bairdii* demonstrates the least proximal enlargement of the palmar sagittal ridges, more notably on MC2 and MC4 (Figure 10). T.

bairdii also demonstrates a compressed proximal carpal row, indicative of resistance to compressive

663 forces in large quadrupeds (Prothero 2005), which may represent an adaptation towards

664 graviportalism not seen in the upper forelimb of this species. The manus of *T. bairdii* demonstrates

greater potential for mobility of the medial digit (MC2; Figure 9), and small centroid size of MC5

666 compared to that of other neotropical species (Table 2). From these adaptations, we posit that *T*.

bairdii, despite its large size, has reduced functionality of the most lateral digit in favour of the

second, third and fourth digits, strongly supporting both anatomical and biomechanical mesaxonic

59 60

Journal of Morphology

MacLaren 29

2		
3	669	symmetry in this taxon. In contrast, the anatomical features of the autopodium in T. indicus
- 5 6	670	demonstrate adaptations for broader force dissipation across the four digits of the manus; as such, T .
7 8	671	indicus is the only extant tapir that may not adhere to both anatomical and biomechanical
9 10	672	interpretations of mesaxonic symmetry. Kinematic and kinetic research will be necessary to shed
11 12 13	673	greater light upon actual limb loading regimes in this enigmatic, and variable, group of mammals.
14 15	674	
16 17 18	675	Conclusions
19 20	676	From both qualitative and quantitative comparisons of the tanir fore foot, we formulated several
21	070	From both quantative and quantitative comparisons of the taph fore-root, we formulated several
22 23 24	677	hypotheses regarding the morphology of modern tapir carpals and metacarpals, investigating whether
24 25 26	678	the anatomy of the tapir autopodium supports both anatomical and biomechanical interpretations of
27 28	679	mesaxonic symmetry. Discriminant function results support our hypothesis that T. indicus is
29 30	680	morphologically separate from neotropical tapirs, although conclusions from previous studies
31 32	681	regarding carpal morphology and function are shown to require rigorous reassessment. Interspecific
33 34	682	differences among neotropical taxa do not align with previous quantitative comparisons of the
35 36 27	683	forelimb (MacLaren and Nauwelaerts 2016), with T. bairdii rather than T. pinchaque displaying the
37 38 39	684	most divergent neotropical morphologies. Interspecific comparisons and covariance analyses of the
40 41	685	autopodials offer evidence that T. indicus has adapted its metapodials and distal carpals to cope with
42 43	686	higher loading forces than other tapirs, which supports all previous assessments on tapir limb
44 45	687	morphology. Morphometric analysis suggests that T. bairdii places greater reliance on digits II, III
46 47 48	688	and IV than other species, with a decreased load predicted for digit V due to size and distal joint
49 50	689	morphology. Testing this will require further analysis of the kinematics of locomotion in living
51 52	690	tapirs. Conversely, T. indicus and T. pinchaque demonstrate osteological evidence for functional use
53 54	691	of the fifth digit, widely considered as redundant in neotropical tapirs (Earle 1893; Osborn 1929).
55 56 57 58	692	These and other features of the autopodium lead us to believe that both <i>T. indicus</i> and <i>T. pinchaque</i>

MacLaren 30

have retained a fully functional tetradactyl manus, and *Tapirus* as a genus may not necessarily display both anatomical and biomechanical mesaxonic symmetry as has previously been supposed. We conclude that the tetradactyl tapir manus should be considered as a variable locomotor unit with a spectrum of functional adaptations, rather than simply a lingering plesiomorphy. Acknowledgments The authors would like to thank Chris Conroy (MVZ), Christiane Funk (ZMB), Eleanor Hoeger (AMNH), Pepijn Kamminga (RMNH), Josephine Lesur (MNHN), Luc Tyteca (MEO) and Frank Zachos (NHMW) for provision of museum specimens for this study. The authors would also like to thank two anonymous reviewers who provided excellent feedback and ideas for improving the study. Finally we would like to thanks Peter Aerts for assistance in finalising the revised manuscript. This study was financially supported by a FWO (Fonds Wetenschappelijk Onderzoek) scholarship awarded to JM and BOF-UA (Special Research Fund for Universiteit Antwerpen) research grant awarded to SN. References Adams, D.C., Rohlf, F.J. & Slice, D.E., 2004. Geometric morphometrics: Ten years of progress following the "revolution." Italian Journal of Zoology, 71(1), pp.5-16. Bressou, C., 1961. La myologie du tapir (Tapirus indicus L.). Mammalia, 25(3), pp.358-400. Brown, J.C. & Yalden, D.W., 1973. The description of mammals - 2 Limbs and locomotion of terrestrial mammals. Mammal Review, 3(4), pp.107-134. Budras, K.-D., Sack, W.O. & Rock, S., 2003. Thoracic Limb. In K.-D. Budras, W. O. Sack, & S.

Rock, eds. *Anatomy of the Horse*. Stuttgart: Schlutersche, pp. 2–13.

Journal of Morphology

MacLaren 31

2 3	716	Campbell, B., 1936. The comparative myology of the forelimb of the hippopotamus, pig and tapir.
4 5 6	717	Americal Journal of Anatomy, 59(2), pp.201–247.
7 8 9	718	Cignoni, P. et al., 2008. MeshLab: an open-source mesh processing tool. In V. Scarano, R. De
10 11	719	Chiara, & U. Erra, eds. Eurographics Italian Chapter Conference. Salerno, pp. 1-8.
12 13 14	720	Clayton, H.M., Chateau, H. & Back, W., 2013. Forelimb function. In W. Back & H. M. Clayton, eds.
15 16 17	721	Equine Locomotion. London: Saunders Elsevier, pp. 99-125.
17 18 19	722	Colbert, M., 2005. The facial skeleton of the early Oligocene Colodon (Perissodactyla, Tapiroidea).
20 21 22	723	Palaeontologia Electronica.
23 24	724	Constantinescu, G.M. et al., 2012. Illustrated Veterinary Anatomical Nomenclature Third. G. M.
25 26 27	725	Constantinescu & O. Schaller, eds., Stuttgart: Enke.
28 29	726	Cozzuol, M.A. et al., 2013. A new species of tapir from the Amazon. Journal of Mammalogy, 94(6),
30 31 32	727	pp.1331–1345.
33 34	728	Downer, C.C., 1996. The mountain tapir, endangered "flagship" species of the high Andes. Oryx,
35 36 37	729	30(1), pp.45–58.
38 39 40	730	Earle, C., 1893. Some points in the comparative osteology of the tapir. Science, 21(526), p.118.
41 42 43	731	Earle, C., 1896. Tapirs past and present. American Journal of Science, 4(104), pp.934–935.
44 45	732	Easton, K.L. & Kawcak, C.E., 2007. Evaluation of increased subchondral bone density in areas of
46 47 48	733	contact in the metacarpophalangeal joint during joint loading in horses. American Journal of
49 50	734	Veterinary Research, 68(8), pp.816–821.
51 52 53	735	Fadda, C. & Corti, M., 2001. Three-dimensional geometric morphometrics of Arvicanthis:
53 54 55	736	implications for systematics and taxonomy. Journal of Zoological Systematics and Evolutionary
56 57 58 59 60	737	<i>Research</i> , 39(4), pp.235–245.

2	720	Gould EDH 2014 To 3D or not to 3D that is the question: do 3D surface analyses improve the
4	/ 50	Gould, F.D.H., 2014. 10 5D of not to 5D, that is the question. do 5D surface analyses improve the
5 6 7	739	ecomorphological power of the distal femur in placental mammals? <i>PloS one</i> , 9(3), p.e91719.
8 9	740	Gregory, W.K., 1929. Mechanics of locomotion in the evolution of limb structure as bearing on the
10 11	741	form and habits of the titanotheres and the related odd-toed ungulates. In H. F. Osborn, ed. The
12 13	742	titanotheres of ancient Wyoming, Dakota and Nebraska. Washington D.C.: United States
14 15 16	743	Government Printing Office, pp. 727–756.
17 18	744	Hawkins, P.L., 2011. Variation in the Modified First Metatarsal of a Large Sample of Tapirus
19 20 21	745	polkensis, and the Functional Implication for Ceratomorphs. East Tennessee State University.
22 23 24	746	Hellmund, M., 2005. A three-dimensional skeletal reconstruction of the Middle Eocene
25 26	747	Propalaeotherium hassiacum Haupt 1925 (Equidae, Perissodactyla, Mammalia) and a modern
27 28	748	synoptic painting of some individuals within their habitat. In Current Research in Vertebrate
29 30	749	Palaeontolgy 3rd Annual Meeting of the European Association of Vertebrate Palaeontologists
31 32 33	750	(EAVP). Hessisches Landesmuseum Darmstadt, p. 15.
34 35	751	Hershkovitz, P., 1954. Mammals of Northern Colombia, Preliminary Report No. 7: Tapirs (genus
36 37 28	752	Tapirus) with a systematic review of American species. Proceedings of the United States
39 40	753	National Museum, 103, pp.465–496.
41 42 42	754	Hildebrand, M., 1985. Walking and running. In M. Hildebrand et al., eds. Functional Vertebrate
43 44 45	755	Morphology. Cambridge: Harvard University Press, pp. 38–57.
40 47 48	756	Holanda, E.C., Ribeiro, A.M. & Ferigolo, J., 2012. New material of Tapirus (Perissodactyla:
49 50	757	Tapiridae) from the Pleistocene of southern Brazil. Revista Mexicana de Ciencias Geologicas,
51 52 53	758	29(2), pp.308–318.
54 55	759	Holbrook, L., 2001. Comparative osteology of early Tertiary tapiromorphs (Mammalia,
56 57 58 59	760	Perissodactyla). Zoological Journal of the Linnean Society, 132(1), pp.1–54.

Page 33 o	53 Journal of Morphology
1	MacLaren 33
2 3 76	Holbrook, L.T., 2009. Osteology of <i>Lophiodon</i> Cuvier, 1822 (Mammalia, Perissodactyla) and its
4 5 76 6	phylogenetic implications. Journal of Vertebrate Paleontology, 29(1), pp.212–230.
7 8 76 9	Holbrook, L.T., 1999. The phylogeny and classification of tapiromorph perissodactyls (Mammalia).
10 76 11	<i>Cladistics</i> , 15(3), pp.331–350.
13 76 14	Hulbert, R.C., 2010. A new early Pleistocene tapir (Mammalia: Perissodactyla) from Florida, with a
15 76 16	review of Blancan tapirs from the state. Bulletin of the Florida Museum of Natural History,
17 18 76 19	49(3), pp.67–126.
20 21 76	Hulbert, R.C. et al., 2009. Cranial morphology and systematics of an extraordinary sample of the
22 23 76 24	Late Neogene dwarf tapir, <i>Tapirus polkensis</i> (Olsen). <i>Journal of Paleontology</i> , 83(2), pp.238–
25 77 26	262.
27 28 77 29	Hulbert, R.C., 2005. Late Miocene Tapirus (Mammalia, Perissodactyla) from Florida, with
30 77 31	description of a new species, <i>Tapirus webbi</i> . Bulletin of the Florida Museum of Natural History,
32 77 33 34	45(4), pp.465–494.
35 77 36	Hulbert, R.C., 1995. The giant tapir, <i>Tapirus haysii</i> , from Leisey Shell Pit 1A and other Florida
37 77 38 20	Irvingtonian localities. <i>Bulletin of the Florida Museum of Natural History</i> , 37(16), pp.515–551.
40 77 41	5 IBM, 2013. IBM SPSS Statistics for Windows.
42 43 77 44	Janis, C., 1984. Tapirs as Living Fossils. In N. Eldredge & S. M. Stanley, eds. Living Fossils.
45 77 46 47	Casebooks in Earth Sciences. New York, NY: Springer New York, pp. 80–86.
48 77 49	Janis, C.M., 1989. A climatic explanation for patterns of evolutionary diversity in ungulate
50 78 51 78	mammals. <i>Palaeontology</i> , 32(3), pp.463–481.
53 78 54	Klaits, B.G., 1971. The moving mesaxonic manus: a comparison of tapirs and rhinoceroses.
55 78 56 57 57 58 59	<i>Mammalia</i> , 36(1), pp.126–145.
60	John Wiley & Sons

MacLaren 34

783 Klaits, B.G., 1972. The moving mesaxonic manus - a comparison of tapirs and rhinoceroses.

- *Mammalia*, 36(1), pp.126–145.
- 785 Klingenberg, C.P., 2011. MorphoJ: an integrated software package for geometric morphometrics.

Molecular ecology resources, 11(2), pp.353–7.

787 Klingenberg, C.P., 2009. Morphometric integration and modularity in configurations of landmarks:

tools for evaluating a priori hypotheses. *Evolution & Development*, 11(4), pp.405–421.

789 Liebich, H.-G., Konig, H.E. & Maierl, J., 2007. Forelimb or thoracic limb (membra thoracica). In H.

790 E. Konig & H.-G. Liebich, eds. *Veterinary Anatomy of Domestic Animals: Textbook and Colour*

- *Atlas*. Stuttgart: Schlutersche, pp. 145–214.
- The Total To
- widespread convergence in developmental dynamics. *Geological Quarterly*, 53(4), pp.415–432.

794 MacFadden, B.J., 1992. Fossil Horses: Systematics, Paleobiology and Evolution of the Family

Equidae, Cambridge: Cambridge University Press.

- 796 MacLaren, J.A. & Nauwelaerts, S., 2016. A three-dimensional morphometric analysis of upper
- forelimb morphology in the enigmatic tapir (Perissodactyla: *Tapirus*) hints at subtle variations
 in locomotor ecology. *Journal of Morphology*.

799 Murie, J., 1871. The Malayan tapir. *Journal of Anatomy and Physiology*, 6(1), pp.131–512.

- Nauwelaerts, S., Vangeel, K. & Aerts, P., 2016. Loading distribution over the four fingers of the
 tapir during locomotion. In Washington D.C.
- Norman, J.E. & Ashley, M. V., 2000. Phylogenetics of Perissodactyla and tests of the molecular
 clock. *Journal of Molecular Evolution*, 50(1), pp.11–21.

804 Osborn, H.F., 1929. Theories as to the origin, ancestry, and adaptive radiation of the titanotheres and

Journal of Morphology

MacLaren 35

1		
2 3 4	805	other odd-toed ungulates. In H. F. Osborn, ed. The titanotheres of ancient Wyoming, Dakota
5 6	806	and Nebraska. Washington D.C.: United States Government Printing Office, pp. 757-804.
7 8 9	807	Padilla, M., Dowler, R.C. & Downer, C.C., 2010. Tapirus pinchaque (Perissodactyla: Tapiridae).
10 11	808	Mammalian Species, 42(1), pp.166–182.
12 13 14	809	Pereira, S.G., 2013. Anatomia ossea, muscular e consideracoes adaptativas do membro toracico de
15 16	810	Tapirus terrestris (Perissodactyla, Tapiridae). Universidade Federal de Uberlandia.
17 18 19	811	Prothero, D.R., 2005. Postcranial Osteology. In The Evolution of North American Rhinoceroses.
20 21	812	Cambridge: Cambridge University Press, pp. 146–181.
22 23 24 25	813	R Core Development Team, 2008. R: a language and environment for statistical computing.
25 26 27	814	Radinsky, L.B., 1965. Evolution of the tapiroid skeleton from Heptodon to Tapirus. Bulletin of The
28 29	815	Museum of Comparative Zoology, 134, pp.69–106.
30 31 32	816	Rajkumar, H.S. & Klein, H., 2014. First perissodactyl footprints from Flysch deposits of the Barail
33 34	817	Group (Lower Oligocene) of Manipur, India. Proceedings of the Indian Academy of Sciences.
35 36 37	818	Earth and planetary sciences, 123(2), pp.413–420.
38 39	819	Reghem, E. et al., 2012. Hand posture in the grey mouse lemur during arboreal locomotion on
40 41 42	820	narrow branches. Journal of Zoology, 288(1), pp.76–81.
43 44	821	Rohlf, F.J. & Corti, M., 2000. Use of two-block partial least-squares to study covariation in shape.
45 46 47	822	Systematic Biology, 49(4), pp.740–53.
48 49	823	Rohlf, F.J. & Slice, D., 1990. Extensions of the Procrustes method for the optimal superimposition of
50 51 52	824	landmarks. Systematic Zoology, 39(1), pp.40–59.
53 54	825	Ruiz-García, M. et al., 2016. Mitogenomics of the mountain tapir (Tapirus pinchaque, Tapiridae,
55 56 57 58 59 60	826	Perissodactyla, Mammalia) in Colombia and Ecuador: phylogeography and insights into the

MacLaren 36

2
3
4
5
7
8
9
10
11
1Z 13
14
15
16
17
10
20
21
22
23
24
25 26
27
28
29
30
31
33
34
35
36
37
30 39
40
41
42
43
44 45
46
47
48
49
50 51
52
53
54
55
56 57
ว/ 58
59
60

1

827 origin and systematics of the South American tapirs. *Mammalian Biology - Zeitschrift für*828 *Säugetierkunde*, 81(2), pp.163–175.

Ruiz-García, M. et al., 2012. Phylogeography of the mountain tapir (*Tapirus pinchaque*) and the

830 Central American tapir (*Tapirus bairdii*) and the origins of the three Latin-American tapirs by

831 means of mtCyt-B sequences. In K. Anamthawat-Jnsson, ed. *Current Topics in Phylogenetics*

and Phylogeography of Terrestrial and Aquatic Systems. InTech, pp. 83–116.

Schoch, R.M., 1989. A review of the tapiroids. In D. R. Prothero & R. M. Schoch, eds. *The evolution of perissodactyls*. New York: Oxford University Press, pp. 299–320.

835 Simpson, G.G., 1945. Notes on Pleistocene and recent tapirs. *Bulletin of the American Museum of*836 *Natural History*, 86(2), pp.33–82.

837 Steiner, C.C. & Ryder, O.A., 2011. Molecular phylogeny and evolution of the Perissodactyla.

838 Zoological Journal of the Linnean Society, 163(4), pp.1289–1303.

de Thoisy, B. et al., 2014. *Tapir Veterinary Manual* 2nd ed. V. Quse & R. C. Fernandes-Santos, eds.,

840 IUCN/SSC Tapir Specialist Group.

841 Tougard, C. et al., 2001. Phylogenetic relationships of the five extant Rhinoceros species

842 (Rhinocerotidae, Perissodactyla) based on mitochondrial cytochrome b and 12S rRNA genes.

843 *Molecular phylogenetics and evolution*, 19(1), pp.34–44.

Wiley, D., 2006. Landmark Editor 3.0. *Institute for Data Analysis and Visualization, University of Oregon.*

846 Wood, A.R. et al., 2010. Postcranial functional morphology of *Hyracotherium* (Equidae,

Perissodactyla) and locomotion in the earliest horses. *Journal of Mammalian Evolution*, 18(1),
pp.1–32.

849 Yalden, D.W., 1971. The functional morphology of the carpus in ungulate mammals. *Cells Tissues*

1		
2 3 4	850	<i>Organs</i> , 78(4), pp.461–487.
5 6 7	851	Zelditch, M.L., Swiderski, D.L. & Sheets, H.D., 2012. Geometric Morphometrics for Biologists: A
8 9	852	Primer, New York: Academic Press.
9 10 11 23 45 67 89 01 23 22 24 25 67 89 01 23 34 56 78 90 41 23 44 56 78 90 12 34 56 78 90 41 23 45 67 89 00 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 45 67 89 00 123 34 56 78 90 123 45 67 89 00 123 45 67 89 00 123 45 67 89 00 123 45 67 89 00 123 34 56 78 90 123 34 56 78 90 123 45 67 89 00 123 45 67 89 00 123 45 89 00 123 45 67 89 00 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 34 56 78 90 123 55 56 7 56 7 56 7 56 7 56 7 56 7 56 7	853	to per perion





298x154mm (150 x 150 DPI)

John Wiley & Sons

Journal of Morphology

Graphical Abstract Text:

A simplified phylogeny demonstrating gross differences in the manus morphology of modern tapirs. Bones with lighter colours (higher specificity %) represent higher levels of morphological difference between species, darker colours represent bones that are misclassified more frequently. Each tapir species is represented by a silhouette diagram demonstrating generalised appearance.

- Table 1. Jack-knifed classification accuracy for autopodial specimen assignments from linear
- discriminant analysis. Specificity of classification for each bone are presented alongside % accuracy
- following jack-knifing the dataset.

	Autopodial Bone		% Accuracy			
	-	T. bairdii	T. indicus	T. pinchaque	T. terrestris	-
	Pisiform	1.00	1.00	1.00	0.80	94.7
	Cuneiform	0.60	1.00	0.75	0.60	75.0
	Lunate	1.00	1.00	1.00	0.80	95.0
	Scaphoid	1.00	1.00	1.00	1.00	100
	Trapezoid	1.00	1.00	1.00	1.00	100
	Magnum	1.00	1.00	0.67	1.00	95.2
	Unciform	1.00	1.00	1.00	1.00	100
	MC2	1.00	1.00	1.00	1.00	100
	MC3	1.00	1.00	1.00	1.00	100
	MC4	1.00	1.00	0.75	0.83	90.9
	MC5	1.00	1.00	1.00	0.83	95.5
5 6 7 8 9 0						

Table 2. Average centroid sizes per species for each bone in the autopodium. Mean average and

17 standard deviation are reported for each species. Centroids based on full adult specimens (excluding

sub-adults), with number of adult specimens for each bone also listed (n).

Bone		T. bairdii	T. indicus	T. pinchaque	T. terrestris
Digiform	mean	54.73 ± 5.6	64.46 ± 3.3	51.42 ± 5.6	53.55 ± 3.3
PISHOIIII	п	3	5	3	5
Cunaiform	mean	59.48 ± 1.5	64.81 ± 1.2	54.65 ± 2.4	55.59 ± 5.1
Cullenoini	п	3	6	3	5
Lunata	mean	74.09 ± 2.1	83.86 ± 3.0	69.67 ± 1.0	68.40 ± 7.1
Lunate	п	3	6	3	5
Sambaid	mean	71.43 ± 2.7	86.83 ± 3.5	68.13 ± 3.5	69.24 ± 6.8
Scaphold	п	3	7	T. pinchaque T. terr 3.3 51.42 ± 5.6 53.5 3 31.2 54.65 ± 2.4 55.5 3 31.2 54.65 ± 2.4 55.5 3 30 69.67 ± 1.0 68.4 3 3.5 68.13 ± 3.5 69.2 3 3.6 70.90 ± 1.3 74.1 3 3.0 70.90 ± 1.3 74.1 3 3.7 218.25 ± 9.5 214.3 3 3.7 218.25 ± 9.5 214.3 3.7 202.33 ± 7.3 202.0 3 $3.131.3$ $3.131.3$	5
Trapazaid	mean	37.52 ± 1.6	43.08 ± 2.1	32.28 ± 0.4	34.41 ± 2.4
Trapezoid	п	3	4	3	5
Magnum	mean	76.50 ± 0.7	87.01 ± 3.0	70.90 ± 1.3	74.15 ± 4.7
Magnum	п	3	7	3	6
Unaiform	mean	71.61 ± 2.0	79.44 ± 2.8	66.00 ± 0.5	69.52 ± 5.5
Unchorn	п	3	6	3	5
MC2	mean	214.74 ± 9.5	228.14 ± 3.7	218.25 ± 9.5	214.27 ± 9.2
WIC2	п	3	7	3	6
MC3	mean	256.57 ± 15.0	272.32 ± 7.4	263.21 ± 7.9	256.65 ± 11.5
MC3	п	3	7	64.46 ± 3.3 51.42 ± 5.6 5 5 3 64.81 ± 1.2 54.65 ± 2.4 5 6 3 83.86 ± 3.0 69.67 ± 1.0 6 6 3 86.83 ± 3.5 68.13 ± 3.5 6 7 3 3 43.08 ± 2.1 32.28 ± 0.4 3 4 3 7 3 7 3 7 3 70.1 ± 3.0 70.90 ± 1.3 7 7 3 7 3 72.32 ± 7.4 263.21 ± 7.9 25 7 3 202.33 ± 7.3 20 7 3 3 3 218.12 ± 7.0 202.33 ± 7.3 20 7 3 3 3	6
MC4	mean	202.86 ± 10.7	218.12 ± 7.0	36.83 ± 3.5 68.13 ± 3.5 69.24 ± 3.5 7 3 5 43.08 ± 2.1 32.28 ± 0.4 34.41 ± 3.5 4 3 5 87.01 ± 3.0 70.90 ± 1.3 74.15 ± 7 7 3 6 7 3 6 79.44 ± 2.8 66.00 ± 0.5 69.52 ± 6 6 3 5 228.14 ± 3.7 218.25 ± 9.5 214.27 7 3 6 272.32 ± 7.4 263.21 ± 7.9 256.65 ± 7 7 3 6 7 3 6 218.12 ± 7.0 202.33 ± 7.3 202.03 ± 7.3 7 3 6 7 3 6 7 3 6 7 3 6 7 3 6 7 3 6 7 3 6 7 3 6 7 3 6	
WIC4	п	5	7	3	6
MC5	mean	124.66 ± 5.3	153.21 ± 4.4	130.82 ± 4.3	131.20 ± 9.8
MC3	п	3	7	3	6

27 Table 3. Procrustes ANOVA significance test results for subset of landmarks describing

28 metacarpophalangeal facet of Mc2, Mc3, Mc4 and Mc5 across four tapir species. Bold values denote

29 significant differences.

Variable	Sum of Squares	Mean Squares	df	F	Parametric p-value
Species	0.0839	0.001645	51	3.38	<0.01
Metacarpal	0.9891	0.019394	51	39.81	<0.01

Table 4. Tukey-HSD significance test results from one-way ANOVA of hamate-magnum facet area

	Species						
	T. indicus	T. bairdii	T. pinchaque	T. terrestris			
T. indicus		0.731	0.014	0.668			
T. bairdii	0.731		0.110	1.000			
T. pinchaque	0.014	0.110		0.131			
T. terrestris	0.668	1.000	0.131				
4							
_							
5							
5							

33 ratios. Bold values denote significant differences.

36 Table 5. Tukey-HSD significance test results from one-way ANOVA of distal lunate facet area

37 ratios. Bold values denote significant differences.

	Species						
	T. indicus	T. bairdii	T. pinchaque	T. terrestris			
T. indicus		0.249	1.000	0.134			
T. bairdii	0.249		0.324	0.004			
T. pinchaque	1.000	0.324		0.199			
T. terrestris	0.134	0.004	0.199				

Page 43 of 53

Journal of Morphology

Table 6. Two-Block Partial Least Squares analysis results for pairwise comparisons between key
joint articulations. Number of covariance occurrences per combination tested (n), PLS axes
accounting for the greatest covariance are included with % accounted for, correlation coefficient (*r*)
and significance (*p*) for those axes are presented. Bold RV coefficient of integration and *p*-values
represent overall results for the covariation analysis. Species specific comparisons are presented for
trapezoid-magnum facet.

Facet combination (<i>n</i>)	PLS axis	% covar.	r	р	RV	<i>p</i> -value
Unciform-Magnum (20)	PLS1	100.0	0.363	0.117	0.132	0.117
Distal Lunate facets (20)	PLS1	100.0	0.738	< 0.001	0.545	<0.001
Transport Magnum (17)	PLS1	80.7	0.937	< 0.001	0.778	<0.001
Trapezoid-Magnum $(1/)$	PLS2	12.7	0.892	0.006		<0.001
Tapirus indicus (4)	PLS1	78.5	0.959	0.259	0.915	0.086
Tapirus baridii (5)	PLS1	71.6	0.978	0.205	0.667	0.446
Tapirus pinchaque (3)	PLS1	86.3	0.999	0.170	0.992	0.170
Tapirus terrestris (5)	PLS1	93.3	0.992	0.009	0.931	0.009
T_{1} : 10 1 : 1(15)	PLS1	60.6	0.903	0.005	0 41 5	0.089
Trapezoid-Scaphoid (13)	PLS2	19.5	0.727	0.205	0.415	







182x142mm (150 x 150 DPI)





Figure 2. Landmark placement on the seven bones of the tapir carpus. Proximal row (a) – (d) and distal row (e) – (g). Carpus position in the foot depicted within the grey outline (left). Position of the bone in the carpus relative to other elements demonstrated on each autopodium diagram (orange bone). Specific landmark denomination for each bone can be found in Supplementary Information 3. Representative bones from scans of MEO 2203a.

395x288mm (150 x 150 DPI)



Figure 3. Landmark placement on the four tapir metacarpals. (a) metacarpal II, (b) metacarpal III, (c) metacarpal IV and (d) metacarpal V. Metacarpal position in the foot depicted within the grey outline. Position of each metacarpal relative to other bones portrayed in each autopodium diagram (orange bone). Specific landmark denomination for each bone can be found in Supplementary Information 3. Representative bones from scans of MEO 2204b.

388x260mm (150 x 150 DPI)



Figure 4. Specificity of autopodial discrimination across four tapir species. From top left: (a) *Tapirus indicus;*(b) *T. bairdii;* (c) *T. pinchaque;* (d) *T. terrestris.* Bones of the autopodium shaded to represent the accuracy of classification from Linear Discriminant Analysis. Darker colours represent lower % specificity, with light colours representing high % accuracy of interspecific discrimination. *T. indicus* demonstrates 100% classification accuracy; the cuneiform represents the bone most frequently misclassified across neotropical taxa.

193x243mm (150 x 150 DPI)



Figure 5. Results of 2B-PLS regression analyses comparing (a) unciform and anterior magnum facet area, (b) distal lunate facet areas, (c) trapezoid and magnum articulation facets, and (d) trapezoid and scaphoid articulation facets. Symbols: *Tapirus indicus* (squares), *T. bairdii* (diamonds), *T. pinchaque* (triangles) and *T. terrestris* (circles); dotted line denotes line of best fit for all data-points. Statistical data for these plots can be found in Table 6.

191x183mm (150 x 150 DPI)





Figure 6. Ratio of areas for unciform (blue) and anterior magnum (yellow) facets of the lunate (intermediate carpal). Box plots represent intraspecific variation, with black bar highlighting the mean value; open circles represent outlying specimens. Representative bones (nearest to mean facet value) and ratios: *T. indicus* (RMNH 17923; 0.09), *T. bairdii* (MVZ 141173; 0.13), *T. pinchaque* (MNHN 1982-34; 0.21), *T. terrestris* (RMNH 12827; 0.15).

190x231mm (150 x 150 DPI)



Figure 7. Ratio of areas for distal facets of the lunate. Box plots represent intraspecific variation, with black bar highlighting the mean value. Representative bones warped mean landmark configurations applied to RMNH 43495. Facets highlighted on representative bones: anterior distal facet (to proximal unciform) (blue); posterior distal facet (to volar process of magnum) (green).

190x221mm (150 x 150 DPI)



Figure 8. Comparison of pisiform-ulna articular morphology in *Tapirus terrestris* (a) and *T. pinchaque* (b). Ulnae scaled to same size. Red shaded areas on both ulnae and pisiforms represent articular surface. Ulnae and pisiform depicted in posterior view (pisiform reflected from joint facet; pisiform with landmarks depicted from dorsolateral view. Approximate placement of *m. flexor carpi ulnaris* insertion (i) and carpal retinaculum (grey; connecting pisiform and ulna) is shown. piLm 4 represents landmark most heavily affecting classification along DF 1 for pisiform. Bones represented depict average morphology for *T. terrestris* and *T. pinchaque* applied to scans of MNHN 1982-34.

191x228mm (150 x 150 DPI)





Figure 9. Comparison of trapezoid (second carpal) facet morphology of the magnum (third carpal) in extant *Tapirus*. Medial view. From top left: (a) *T. bairdii*; (b) *T. indicus*; (c) *T. pinchaque*; (d) *T. terrestris*. Representative facet areas shaded and outlined with landmarks (white circles) used in morphometric analysis; maLm 15 (yellow circle) highly discriminatory along DF 1 for magnum. Concave dorsal margin of trapezoid facet marked in bold black.

187x181mm (150 x 150 DPI)



Figure 10. Comparison of the morphology of the palmar metacarpophalangeal joint facet in extant *Tapirus*. Medial and lateral metacarpals represented: (a) MC2; (b) MC4; (c) MC5. Shaded regions represent approximate facet surface for articulation with the proximal sesamoids either side of the palmar sagittal ridge: green = medial sesamoid; blue = lateral sesamoid. White circles = landmark placement on palmar metacarpophalangeal joint. Average landmark configurations warped onto metacarpals of MEO 2204e.

227x277mm (150 x 150 DPI)