

**This item is the archived peer-reviewed author-version of:**

Trophic ecology drives contaminant concentrations within a tropical seabird community

**Reference:**

Sebastiano Manrico, Bustamante Paco, Eulaers Igor, Govindan Malarvannan, Mendez-Fernandez Paula, Churlaud Carine, Blévin Pierre, Hauselmann Antoine, Covaci Adrian, Eens Marcel, ....- Trophic ecology drives contaminant concentrations within a tropical seabird community  
Environmental pollution - ISSN 0269-7491 - 227(2017), p. 183-193  
Full text (Publisher's DOI): <https://doi.org/10.1016/J.ENVPOL.2017.04.040>  
To cite this reference: <https://hdl.handle.net/10067/1427550151162165141>



## 25 **Abstract**

26 To support environmental management programs, there is an urgent need to know about the  
27 presence and understand the dynamics of major contaminants in seabird communities of key  
28 marine ecosystems. In this study, we therefore investigated the concentrations and  
29 trophodynamics of trace elements in six seabird species and persistent organic pollutants  
30 (POPs) in three seabird species breeding on Grand Connétable Island (French Guiana), an area  
31 where the increase in human population and mining activities has raised concerns in recent  
32 years. Red blood cell (RBC) Hg concentrations in adults were the highest in Magnificent  
33 frigatebirds *Fregata magnificens* (median: 5.6  $\mu\text{g g}^{-1}$  dw; range: 3.8-7.8  $\mu\text{g g}^{-1}$  dw) and lowest  
34 in Sooty terns *Onychoprion fuscatus* (median: 0.9  $\mu\text{g g}^{-1}$  dw; range: 0.6-1.1  $\mu\text{g g}^{-1}$  dw). Among  
35 POPs, dichlorodiphenyldichloroethylene (*p,p'*-DDE) was the most abundant compound in  
36 plasma of Cayenne terns *Thalasseus sandvicensis* (median: 1100  $\text{pg g}^{-1}$  ww; range: 160±5100  
37  $\text{pg g}^{-1}$  ww), while polychlorinated biphenyls (PCBs) were the most abundant compound class  
38 in plasma of Magnificent frigatebirds (median: 640  $\text{pg g}^{-1}$  ww; range 330±2700  $\text{pg g}^{-1}$  ww).  
39 While low intensity of POP exposure does not appear to pose a health threat to this seabird  
40 community, Hg concentration in several adults Laughing gulls *Leucophaeus atricilla* and Royal  
41 terns *Thalasseus maximus*, and in all Magnificent frigatebirds was similar or higher than that  
42 of high contaminated seabird populations. Furthermore, nestling plasma also contained Hg  
43 concentrations of concern, and further studies should investigate its potential health impact in  
44 this seabird community. Differences in adult trophic ecology of the six species explained to  
45 some extent interspecific variation in exposure to trace element and POPs, while nestling  
46 trophic ecology provides indications about the diverse feeding strategies adopted by the six  
47 species, with the consequent variation in exposure to contaminants.

48

## 49 **Capsule**

50 In the present study we found mercury (Hg) concentrations of concern in both adults and  
51 nestlings within a seabird community in a protected area of South America.

52

53 **Keywords:** trace elements; persistent organic pollutants; stable isotopes; French Guiana;  
54 mercury

55

## 56 **1. Introduction**

57 Exposure to persistent toxicants may have detrimental effects on reproductive success,  
58 immunity, regulation of oxidative balance, endocrine system and survival perspectives of  
59 wildlife, even years after these toxicants have been banned (Burger and Gochfeld, 2001;  
60 Costantini et al., 2014; Erikstad et al., 2013; Goutte et al., 2015; Tartu et al., 2015a; Tartu et al.,  
61 2015c). Persistent organic pollutants (POPs) are among the major contaminants currently  
62 detected in wildlife, of which polychlorinated biphenyls (PCBs) remain the most dominant  
63 chemical class despite that they have been banned more than 30 years ago (Tartu et al., 2015b).  
64 Among trace elements there has been growing interest in mercury (Hg), lead (Pb), and cadmium  
65 (Cd) because of their well-known detrimental effects on vertebrates (Beyer et al., 2011). For  
66 example, after Hg is deposited in aquatic ecosystems, it is rapidly transformed by  
67 microorganisms into methyl-Hg, a toxic form that bioaccumulates in organisms and  
68 biomagnifies in food webs (Fitzgerald et al., 2007). Because seabirds are apex long-lived  
69 predators, they are particularly exposed to these major environmental contaminants (Rowe,  
70 2008), and are therefore utilized as sentinel species for environmental monitoring (Furness and  
71 Camphuysen, 1997; Moreno et al., 2011).

72 While considerable attention has been paid to the occurrence and health effects of these  
73 contaminants in sub-polar and polar regions due to their potential to act as final sink (Blévin et  
74 al., 2016; Tartu et al., 2015b; Tartu et al., 2016), comparatively less attention has been given to  
75 wildlife from other geographical regions (Bastos et al., 2015; De Andres et al., 2016; Frery et  
76 al., 2001), especially in South America (De Andres et al., 2016; Guirlet et al., 2010; Sebastiano  
77 et al., 2016). This is surprising because local releases of contaminants (e.g. Hg) from major  
78 mining activities in the Amazon area may be considerable (Fujimura et al., 2012; Lodenius and  
79 Malm, 1998). Furthermore, although to the best of our knowledge there are no sources of  
80 organic pollutants, the long-range transport of these contaminants and the bioaccumulation and

81 biomagnification processes they undergo, might pose a threat to top predators. The Grand  
82 Connétable Island, a small rocky island located off the coast of French Guiana and close to the  
83 Brazilian border, with its strategic position for wildlife and the presence of six breeding seabird  
84 species, offers a unique opportunity to assess the presence and quantify the concentrations of  
85 both organic and inorganic pollutants in a tropical seabird community. Moreover, given the  
86 expected high variation in trophic ecology of the seabirds breeding on the Grand Connétable  
87 Island, this also enabled us to assess the importance of feeding ecology in driving inter- and  
88 intraspecific variation in exposure to contaminants.

89         To this end, we quantified the concentrations of POPs and trace elements in red blood  
90 cells and plasma, respectively, in the seabird community that breed on Grand Connétable Island.  
91 We also measured the stable nitrogen and carbon isotope composition of red blood cells to test  
92 whether trophodynamics explain among and within species variation in contaminant burden.  
93 Of the trace elements, we focused particularly on Hg given the growing concern about the  
94 impact of this element on the health of South-American ecosystems.

95

## 96 **2. Materials and methods**

### 97 **2.1 Sample collection**

98 In 2013 we performed sample collection on the Grand Connétable Nature Reserve, a small  
99 island located 18 km off Cayenne (French Guiana, 4°49'30 N; 51°56'00 W). The seabird  
100 community of Grand Connétable Island typically includes six species: the Laughing gull  
101 (*Leucophaeus atricilla*), the Brown noddy (*Anous stolidus*), the Royal tern (*Thalasseus*  
102 *maximus*), the Cayenne tern (*Thalasseus sandvicensis*), the Magnificent frigatebird (*Fregata*  
103 *magnificens*; hereafter Frigatebird), and the Sooty tern (*Onychoprion fuscatus*) (Dujardin and  
104 Tostain, 1990). These six seabird species differ in both feeding style (pelagic versus benthic)  
105 and foraging area (inshore versus offshore). For instance, Frigatebirds feed on both pelagic and

106 benthic fish by surface dipping, kleptoparasitism and opportunistic feeding (mostly on shrimp  
107 trawler discards), and although most foraging occurs in coastal waters, some foraging trips can  
108 exceed 200 km away from the breeding colony (Weimerskirch et al., 2003). Finally, they are  
109 also seen to follow tuna school formations because tuna and other marine predators push other  
110 small fish toward the surface, making them accessible to frigatebirds. The Laughing gull feeds  
111 on coastal pelagic fish, marine invertebrates and fishery discards, while the Brown noddy feeds  
112 on fish and squid in offshore waters by dipping the surface, and may show kleptoparasitism.  
113 Terns diet consists predominantly of small fish, squids and crustaceans, obtained by dipping  
114 the surface and occasionally diving (del Hoyo et al., 1996; Dujardin and Tostain, 1990).

115         Adult seabirds were sampled during the incubation or early chick rearing (27<sup>th</sup> to 30<sup>th</sup> of  
116 May) while nestlings were sampled within a few weeks after adult sampling (24<sup>th</sup> to 26<sup>th</sup> of  
117 June). A total of 101 adults and 102 nestlings were captured. Since all nestlings were captured  
118 by hand on their nests while adults were captured by mist nets (or, in case of frigatebirds, were  
119 captured with a noose attached to a fishing rod), adults and nestlings are likely unrelated to each  
120 other. For the Laughing gull, Royal tern, Cayenne tern, Sooty tern and Brown noddy, the egg  
121 laying period usually begins around mid-April and ends around the end of April (Dujardin and  
122 Tostain, 1990), and the incubation period lasts 25 to 30 days (except for the Brown noddy that  
123 can take a few more days; Dujardin and Tostain, 1990). Therefore, nestlings of these species  
124 had approximately the same age. Frigatebird, instead, were a few weeks older than the nestlings  
125 of the other species, with an approximate age of three to four months (the age of nestling  
126 frigatebirds was incorrectly reported in Sebastiano et al. 2016). Blood samples (around 2 mL)  
127 were collected from the brachial vein using a heparinized syringe (25 G needle) within a few  
128 minutes after capture, and samples were immediately put on ice. Blood was then centrifuged  
129 within one hour to separate plasma and red blood cells. Both fractions were kept at -20 °C until  
130 laboratory analyses.

131

## 132 **2.2 Stable isotope analysis**

133 The analysis of the carbon and nitrogen stable isotopes is considered an important tool for the  
134 interpretation of both the foraging area and the trophic level of the species. In marine  
135 ecosystems, higher nitrogen values are associated with higher trophic level prey, e.g. bigger  
136 prey (Overman and Parrish, 2001), while the carbon stable isotope can decrease with decreasing  
137 latitudes (Kelly, 2000) and seem to decrease from the coast to the open sea in the Southern  
138 Indian Ocean (Cherel and Hobson, 2007), even if proofs of such stratification in the Southern  
139 Atlantic Ocean are not available. In this study, for instance, the stable carbon and nitrogen  
140 values were measured in RBCs, therefore providing trophic information integrated over a few  
141 weeks prior to sampling (Hobson and Clark, 1993; Newsome et al., 2007). The composition of  
142 the carbon and nitrogen isotopes of the species, which provides information on the isotopic  
143 niches of the birds, was used as a proxy of their ecological niche (Jackson et al., 2011). Analyses  
144 were carried out following a previous protocol (Sebastiano et al., 2016), and results are  
145 expressed as  $\delta$  (‰) for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ , respectively, calibrated against the international isotopic  
146 references (atmospheric nitrogen for  $\delta^{15}\text{N}$  and Pee Dee Belemnite for  $\delta^{13}\text{C}$ ). The experimental  
147 imprecision, based on secondary isotopic reference material, did not exceed  $\pm 0.15$  and  $\pm 0.20$   
148 ‰ for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ , respectively.

149

## 150 **2.3 Contaminant analysis**

151 Trace element concentration analyses (Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se,  
152 V, and Zn) were performed by the Littoral Environnement et Sociétés (LIENSs) laboratory on  
153 lyophilized RBCs as previously described (Sebastiano et al., 2016). In brief, an Altec Advanced  
154 Mercury Analyzer AMA 254 spectrophotometer was used for the quantification of total Hg,  
155 while the other trace elements were quantified using a Varian Vista-Pro ICP-OES or a Series II



156 Thermo Fisher Scientific ICP-MS (aliquots mass: 3-8 mg for AMA and 50–200 mg dw for  
157 ICP). Analyses on blanks and Certified Reference Materials (CRM) from NRCC (dogfish liver  
158 DOLT-4 and lobster hepatopancreas TORT-2) were carried out as for the samples. Since results  
159 showed low standard deviations and were in good agreement with the certified values, the  
160 methodology showed good repeatability. Quantification limits and mean recovery rates were,  
161 respectively, equal to 0.1  $\mu\text{g L}^{-1}$  and 79 % for Ag, 1  $\mu\text{g L}^{-1}$  and 94 % for As, 0.1  $\mu\text{g L}^{-1}$  and 99  
162 % for Cd, 0.1  $\mu\text{g L}^{-1}$  and 97 % for Co, 0.1  $\mu\text{g L}^{-1}$  and 95 % for Cr, 0.5  $\mu\text{g L}^{-1}$  and 96 % for Cu,  
163 20  $\mu\text{g L}^{-1}$  and 92 % for Fe, 0.5  $\mu\text{g L}^{-1}$  and 94 % for Mn, 0.2  $\mu\text{g L}^{-1}$  and 99 % for Ni, 0.1  $\mu\text{g L}^{-1}$   
164 and 89 % for Pb, 0.5  $\mu\text{g L}^{-1}$  and 118 % for Se, 2  $\mu\text{g L}^{-1}$  and 98 % for V, and 20  $\mu\text{g L}^{-1}$  and 105  
165 % for Zn. All trace element concentrations are expressed as  $\mu\text{g g}^{-1}$  dry weight (dw).

166 Persistent organic pollutant analyses were carried out at the University of Antwerp  
167 (Toxicological Centre), following a previous protocol (Sebastiano et al., 2016). The protocol  
168 allowed the detection of 26 PCB congeners (CB 28, 49, 52, 74, 99, 101, 105, 118, 128, 138,  
169 146, 153, 156, 170, 171, 174, 177, 180, 183, 187, 194, 196, 199, 203, 206, and 209),  
170 organochlorine pesticides (OCPs), amongst which dichlorodiphenyltrichloroethane (*p,p'*-DDT)  
171 and its metabolite dichlorodiphenyldichloroethylene (*p,p'*-DDE), hexachlorobenzene (HCB),  
172  $\alpha$ ,  $\beta$ - and  $\gamma$ -hexachlorocyclohexanes (HCHs), the Chlordanes (CHLs) *cis*-nonachlor  
173 (CN), *trans*-nonachlor (TN), and oxychlordane (OxC), and 7 polybrominated diphenyl ethers  
174 (PBDEs: BDE 28, 47, 99, 100, 153, 154, and 183). Plasma samples (around 1mL) were analysed  
175 for POPs. Solid-phase extraction (SPE) on OASIS HLB cartridges was used followed by  
176 fractionation on SPE cartridges topped with 1.5 g of acidified silica (44%  $\text{H}_2\text{SO}_4$ , w/w) and  
177 eluted with 10 mL hexane: dichloromethane (1:1). The cleaned extract was evaporated to  
178 incipient dryness and re-dissolved in 100  $\mu\text{L}$  iso-octane. Quantification of POPs were further  
179 analysed by gas chromatography coupled to mass spectrometry operated either in electron  
180 capture negative chemical ionization (GC-ECNI-MS) or electron ionization (GC-EI-MS)

181 depending on the analyses' sensitivity. All plasma POP concentrations are expressed as  
182  $\text{pg g}^{-1}$  wet weight (ww). POPs were performed in the Cayenne tern, Frigatebird, and Brown  
183 noddy. These species were chosen because they differ in the feeding strategies and they are  
184 representative of the region.

185

#### 186 **2.4. Quality assurance/quality control**

187 The extraction, clean-up, and fractionation steps were evaluated following a previous protocol  
188 (Dimitriadou et al., 2016). Mean  $\pm$  SD recoveries of the internal standards PCB 143,  $\epsilon$ -HCH  
189 and BDE 77 were  $86 \pm 6\%$ ,  $98 \pm 8\%$  and  $93 \pm 10\%$ , respectively. The quality control was  
190 performed by regular analyses of procedural blanks, sample replicates, by random injection of  
191 standards, spiked samples and solvent blanks. The quality control scheme was also assessed  
192 through regular participation to inter-laboratory comparison exercises (POPs in serum)  
193 organized three times per year by the arctic monitoring and assessment program (AMAP, 2015).  
194 The obtained values were deviating with less than 20% from the consensus values.

195

#### 196 **2.5 Statistical analysis**

197 Concentrations of trace elements and POPs below the limit of quantification (LOQ) were  
198 replaced with a value equal to  $\frac{1}{2} \cdot \text{LOQ}$  when the detection frequency was greater than 50%. For  
199 Ag, Co, Cr, and V, concentrations were below the LOQ for all individuals (both adults and  
200 nestlings), and were therefore excluded from the statistical analyses. Furthermore, since the  
201 concentration of Se is important in detoxifying Hg (Ralston and Raymond, 2010), the molar  
202 ratio Hg:Se was also calculated using the formula “[concentration Hg ( $\mu\text{g g}^{-1}$ ) / atomic weight  
203 Hg ( $\text{g mol}^{-1}$ )] / [concentration Se ( $\mu\text{g g}^{-1}$ ) / atomic weight Se ( $\text{g mol}^{-1}$ )]”

204 Linear models were used to test trace element and POP concentration variations across  
205 the six species and both age classes. When the interaction between *species* and *age* was

206 significant, statistical testing was performed for adults and nestlings separately in order to  
207 increase the statistical power by excluding biologically meaningless post-hoc comparisons (e.g.  
208 adults of one species versus nestlings of another species). Then, a principal component analysis  
209 (PCA) based on the correlation matrix was used to reduce contaminant data to fewer  
210 uncorrelated variables. This approach (which has been used only on trace elements since POP  
211 concentrations were very low in all species), enabled us to investigate if the contaminant pattern  
212 was similar between age classes. The Kaiser-Mayer-Olkin measure of sampling adequacy  
213 (KMO=0.58 in nestlings and KMO=0.67 in adults) and the Bartlett's test of sphericity ( $P<0.01$   
214 in both groups), confirmed the appropriate use of the PCA. After the scree plot was examined,  
215 components with an eigenvalue  $> 1$  were selected. Parametric correlations were used to test  
216 associations between stable isotope values in the different species and age classes, which is  
217 important to describe the association between the trophic level of the species and the location  
218 of the food source. Any data transformation to achieve normality or any violation of models  
219 assumptions is reported in the manuscript when necessary. The graphic representation of the  
220 isotopic niche of the different species has been performed using Stable Isotope Bayesian  
221 Ellipses in R (SIBER) package (Jackson et al., 2011). In order to graphically compare individual  
222 groups within the community with each other, we used the Standard Ellipse Area (SEA) method  
223 (Jackson et al., 2011). In this package, the stable carbon and nitrogen isotopes are used to  
224 calculate the isotopic niche of each species in the community, allowing a comparison among  
225 species. Furthermore, the calculation of the ellipse area of each species is not influenced by the  
226 sample size, allowing a graphical and statistical comparison among different species or diverse  
227 studies with different samples sizes (Jackson et al., 2011). Furthermore, linear models  
228 estimating Hg or *log*-transformed POP class concentrations based on stable isotope values were  
229 also carried out on both the pooled data and separately for each species. For pooled data (overall  
230 models), linear mixed models using species as a random effect were applied in order to control

231 for the non-independence of data points. All statistical analyses were performed using R (3.1.1  
232 version).

233

### 234 **3. Results**

#### 235 **3.1 Stable isotopes**

236 The overall correlation between  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  was significant both in adults ( $n=101$ ;  $r=0.54$ ;  
237  $P<0.01$ ; Figure 1, Table 1) and nestlings ( $n=102$ ;  $r=0.88$ ;  $P<0.01$ ; Figure 1, Table 1), and  
238 density plots of the Standard Ellipse Area (SEA) representing the niche of the six species are  
239 shown in Figure 2. Species-specific correlations between isotopes, which explain the  
240 relationship between the trophic level and the foraging area of each species, are shown in Table  
241 1.  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  levels significantly differed among species and also within the same species  
242 between adults and nestlings ( $\delta^{13}\text{C}$ :  $F=36.40$ ,  $P<0.01$  and  $\delta^{15}\text{N}$ :  $F=21.50$ ,  $P<0.01$ , respectively),  
243 and interspecific differences in the isotope values between adults and nestlings are shown in  
244 Figure S1 and Table S1.

245

#### 246 **3.2 Trace elements**

247 Cd was detected only in the Brown noddy (both adults and nestlings) and in Sooty tern adults,  
248 while Ni was detected in Laughing gull and Brown noddy nestlings only. Linear models for Cu  
249 and Pb did not show a significant interaction between species and age, and post-hoc analysis  
250 was not carried out. For the other trace elements (As, Fe, Hg, Mn, Se, and Zn), the interaction  
251 between species and age was significant ( $F>5.60$ ;  $P<0.01$ ), and post-hoc comparison of  
252 contaminants within age classes are shown in Figure 3. Average trace elements, concentrations,  
253 median along with standard deviation (SD) values and their range are shown in Table 2.

254 PCA on adults reduced the variation in the concentrations of eight trace elements to two  
255 principal components (PCs), explaining 55 % of the total variance (Figure 4a). The first axis

256 indicated that adults with high PC1 scores are associated with high levels of As (loading =  
257 0.497) and Se (0.441), and low levels of Cu (-0.396), Mn (-0.383), Pb (-0.371), Zn (-0.237),  
258 and Hg (-0.232), while the second axis indicated that adults with high PC2 scores are associated  
259 with low levels of Hg (-0.618) and high levels of Zn (0.448), Cu (0.397), Se (0.346), Pb (0.280),  
260 and As (0.223), as showed in Figure 4a. The factor species significantly explained the variation  
261 of trace elements represented by both PC1 ( $F=75.70$ ;  $P<0.01$ ) and PC2 ( $F=47.40$ ;  $P<0.01$ ).  
262 Moreover, both PC1 and PC2 were significantly negatively correlated to  $\delta^{13}\text{C}$  (PC1:  $r=-0.61$ ;  
263  $P<0.01$ ; PC2:  $r=-0.57$ ;  $P<0.01$ ) and  $\delta^{15}\text{N}$  (PC1:  $r=-0.31$ ;  $P<0.01$  and PC2:  $r=-0.63$ ;  $P<0.01$ ).

264 PCA performed on nestlings also reduced the variation of concentrations for the eight  
265 trace elements to two PCs, explaining 62 % of the total variance (Figure 4b). The first axis  
266 indicated that nestlings with high PC1 scores are associated with high levels of Se (loading =  
267 0.458) and As (0.323) and low levels of Mn (-0.444), Fe (-0.380), Hg (-0.366), and Zn (-0.353),  
268 while the second axis indicated that adults with high PC2 scores are associated with high levels  
269 of Cu (0.629), Pb (0.438), and As (0.339), and low levels of Hg (-0.432), as showed in Figure  
270 4b. The factor species significantly explained the variation of trace elements represented by  
271 both PC1 ( $F=132.50$ ;  $P<0.01$ ) and PC2 ( $F=104.60$ ;  $P<0.01$ ). Moreover, PC1 was significantly  
272 negatively correlated to  $\delta^{13}\text{C}$  ( $r=-0.66$ ;  $P<0.01$ ) and  $\delta^{15}\text{N}$  ( $r=-0.75$ ;  $P<0.01$ ), while PC2 was not  
273 correlated with  $\delta^{13}\text{C}$  nor  $\delta^{15}\text{N}$ .

274

### 275 **3.3 Mercury**

276 Hg concentrations widely varied among species, ranging from a minimum in a Brown noddy  
277 nestling ( $0.1 \mu\text{g g}^{-1}$  dw, Table 2) and a maximum value in an adult Frigatebird ( $7.8 \mu\text{g g}^{-1}$  dw,  
278 Table 2). Concentrations in adults were highest in Frigatebirds (median:  $5.6 \mu\text{g g}^{-1}$  dw; range:  
279  $3.8\text{-}7.8 \mu\text{g g}^{-1}$  dw) and lowest in Sooty terns (median:  $0.9 \mu\text{g g}^{-1}$  dw; range:  $0.6\text{-}1.1 \mu\text{g g}^{-1}$  dw).  
280 In nestlings, Hg concentrations were highest in Frigatebirds (median:  $1.0 \mu\text{g g}^{-1}$  dw; range:  $0.7\text{-}$

281 1.7  $\mu\text{g g}^{-1}$  dw) and lowest in Brown noddy (median: 0.1  $\mu\text{g g}^{-1}$  dw; range: 0.1-0.2  $\mu\text{g g}^{-1}$  dw).  
282 Linear models showed a significant and positive association between Hg concentrations and  
283  $\delta^{15}\text{N}$  values in nestlings (Table 3, Figure 5), and between Hg concentrations and  $\delta^{13}\text{C}$  values  
284 both in adults and nestlings (Table 3).

285 In adult individuals, Hg increased with  $\delta^{15}\text{N}$  in Frigatebirds, and Hg also increased with  
286  $\delta^{13}\text{C}$  in all species (Table 3). In nestlings, Hg increased with  $\delta^{15}\text{N}$  in all species, while there was  
287 not association between Hg and  $\delta^{13}\text{C}$  in any species (Table 3). Finally, the molar ratio Hg:Se  
288 did not exceed 1 in any sampled individual (both adults and nestlings). The Hg:Se ratio showed  
289 an average value of  $0.053 \pm 0.085$  (median: 0.015), ranging from 0.001 in a Sooty tern adult to  
290 0.407 in a Frigatebird adult.

291

### 292 **3.4 POPs**

293 POP measurements were only performed in the Cayenne tern, Frigatebird, and Brown noddy,  
294 and their concentrations widely varied across these three species in a compound-specific way.  
295 Since *p,p'*-DDT was not detected in more than half of the individuals per species, comparisons  
296 were made for *p,p'*-DDE only. Furthermore, statistics were not performed for POP compounds  
297 below the LOQ (Table 4). Average, median, ranges along with standard deviation (SD) values  
298 of POPs are shown in Table 4.

299 Among *DDTs*, *p,p'*-DDE was the most abundant POP in adult Cayenne tern (median:  
300 1,100  $\text{pg g}^{-1}$  ww; range: 160-5100  $\text{pg g}^{-1}$  ww), while PCBs were the most abundant POP class  
301 in adult Frigatebird (median: 640  $\text{pg g}^{-1}$  ww; range: 330-2700  $\text{pg g}^{-1}$  ww). Adult Brown noddy  
302 had very low levels of POPs in comparison with the other two species, while *p,p'*-DDE was the  
303 most abundant compound in this species (median: 200  $\text{pg g}^{-1}$  ww; range: 5-600  $\text{pg g}^{-1}$  ww).  
304 Further statistical analyses could be performed only for  $\sum\text{PCBs}$  and *p,p'*-DDE because all the  
305 other compounds were not detectable in all three species. Log-transformed  $\sum\text{PCBs}$  values

306 showed significant differences among the three species ( $F=52.14$ ;  $P<0.01$ ), with Frigatebirds  
307 having higher PCB concentrations than both the Cayenne tern ( $t=2.78$ ;  $P=0.02$ ) and the Brown  
308 noddy ( $t=9.90$ ;  $P<0.01$ ), while the latter had lower PCB concentrations than Cayenne tern  
309 ( $t=7.12$ ;  $P<0.01$ ). Log-transformed  $P,p'$ -DDE values also showed significant differences  
310 among the three species ( $F=21.68$ ;  $P<0.01$ ), and concentrations in the Cayenne tern were higher  
311 than those in Frigatebirds ( $t=-3.09$ ;  $P<0.01$ ) and Brown noddy ( $t=6.58$ ;  $P<0.01$ ), while the latter  
312 had lower  $p,p'$ -DDE concentrations than Frigatebirds ( $t=3.49$ ;  $P<0.01$ ). In adults,  $\Sigma$ PCBs did  
313 not increase with  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  (Table 3), while there was an association between  $p,p'$ -DDE  
314 and  $\delta^{13}\text{C}$  (Table 3).

315

#### 316 **4. Discussion**

317 The present study provides unique information on the occurrence and trophodynamics of trace  
318 elements and POPs in a seabird community from French Guiana on the east coast of South  
319 America. Our data showed the presence of high levels of Hg and strong associations with stable  
320 carbon and nitrogen isotopes, indicating that feeding is a driver of inter- and intra-specific  
321 variation in RBC levels of Hg. Furthermore, the strong interspecific variation in some trace  
322 elements was also explained by both the stable carbon and the nitrogen isotopes. Finally, our  
323 results confirmed that POPs in French Guiana marine seabirds occur at very low concentrations.

324

##### 325 *4.1 Foraging ecology and trophic niche*

326 Our study has found an overlap in the trophic niches of adults and nestlings for some species  
327 only (Figure 1). Laughing gulls had a wide range of both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values, and was the  
328 only species to show a negative correlation between  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ , likely due to their wide  
329 foraging area and opportunistic feeding (Figure 1 and Supplementary Figure S1). Furthermore,  
330 the Brown noddy and Sooty tern seem to adopt opposite feeding strategies for their nestlings.

331 Specifically, Sooty tern adults showed significantly depleted  $\delta^{15}\text{N}$  values compared to their  
332 nestlings (Supplementary Figure S1), which are likely fed with higher trophic level food, a  
333 strategy to optimize foraging during the chick rearing period (Bugge et al., 2011). Brown noddy  
334 adults, instead, had much enriched  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values compared to nestlings (Supplementary  
335 Figure S1). This does not mean, however, that Brown noddy were feeding their nestlings with  
336 low quality diet. For instance, a previous study has found no support for the hypothesis that  
337 high quality diet contains a greater proportion of upper trophic level prey (Morrison et al.,  
338 2014). Conversely, this study found nestlings with a diet biased for lower trophic level prey to  
339 be in a better body condition than nestlings fed with higher trophic level prey (Morrison et al.,  
340 2014). In Frigatebirds, we would have expected depleted  $\delta^{13}\text{C}$  values in comparison to other  
341 species, given their attitude to feed far from the breeding colony and possibly in southern  
342 latitudes (Sebastiano et al., 2016), but they were not, possibly because Frigatebirds also feed on  
343 the shrimp fishery discards, a strategy commonly used in this colony (Martinet and Blanchard,  
344 2009).

345 Standard ellipse areas also confirmed a difference in the foraging ecology of the six  
346 species and provided some insight on their feeding habits. Figure 2 clearly shows how  
347 Frigatebird, Brown noddy and Cayenne tern adults are specialized foragers, while the Royal  
348 tern, the Sooty tern and especially the Laughing gull adults are generalists. This scenario is,  
349 however, different for their nestlings. Most species (except Sooty tern nestlings for which the  
350 sample size is small) seem to have a more generalist foraging strategy, especially for Brown  
351 noddy, and Frigatebirds, which might indicate, for instance, that during the reproductive period  
352 adults may catch prey items that are not generally present in their diet. However, this pattern  
353 was not found in Laughing gull nestlings, which showed a consistently smaller SEA than adults.  
354 This might indicate, for instance, that although the general diet of gulls includes a huge variety  
355 of prey items and might also include discards from the shrimp fishery and, in some populations,



356 human waste products, adults might restrict the variety of food that is given to their nestlings  
357 during their development.

358

#### 359 *4.2 Trace element exposure*

360 In addition to providing information on the foraging ecology of the entire community,  $\delta^{15}\text{N}$  and  
361  $\delta^{13}\text{C}$  values explained interspecific variation in the exposure to trace elements. Indeed, trace  
362 element concentrations were significantly different among the investigated species, both in  
363 adults and in nestlings, especially for As, Hg, and Se (Figure 3). Adult Sooty tern, Brown noddy,  
364 and partially the Cayenne tern were more associated with the first PC, while both Frigatebird  
365 and Sooty tern were more associated with the PC2. Both PC1 and PC2 values were significantly  
366 correlated to  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values, indicating that inter-individual differences in trace element  
367 concentrations in adults can be explained by both the trophic level and the foraging areas of the  
368 species. This means, for instance, that the accumulation pattern of some trace elements might  
369 be more related to the foraging area of the species, while it might be more dependent on the  
370 trophic level of the species for other trace elements, and further studies are needed clarify this  
371 aspect.

372         Regarding the nestlings, a similar pattern was found (Figure 4b). Most species, except  
373 for the Sooty tern and the Royal tern, were associated with the first PC. Frigatebird, Laughing  
374 gull and Royal tern, instead, were more associated with the PC2. The variation in PC1 values  
375 was correlated to  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values, while PC2 values were not.

376 To the best of our knowledge, the present study is the first to show this pattern between the  
377 levels of As, Hg, and Se in a seabird community. Species with high levels of Hg have also low  
378 As and Se. Studies have shown that As might be an essential trace element (Nielsen, 1998;  
379 Uthus, 2003), and a more recent study has highlighted how the main function of As might be

380 to maintain optimal levels of S-adenosylmethionine (Uthus, 2003), a common co-substrate  
381 which is involved in diverse essential metabolic pathways (Lu, 2000).

382 Selenium, first known as a toxic element, is an essential element and the main  
383 component of both selenoproteins, a wide group of proteins with the role of antioxidant  
384 enzymes (e.g. glutathione peroxidase), and enzymes required to maintain an optimal thyroid  
385 functioning, as the iodothyronine deiodinases, and thioredoxin reductase (Tapiero et al., 2003).  
386 Over the past years, given its involvement in some crucial physiological functions (Rayman,  
387 2012), the knowledge on the relationship between Se bioavailability and health status and on  
388 the important role that Se supplementation plays in diseases has radically increased in humans  
389 (Rayman, 2012; Tinggi, 2008). One of the aspects that have been well studied is the “protective  
390 effect” of Se against Hg toxicity (Ackerman et al., 2016; Polak-Juszczak and Robak, 2015;  
391 Sørmo et al., 2011). Indeed, because of the high affinity between these two elements, Hg binds  
392 to Se to produce insoluble tiemannite in the liver of many mammals and certain birds (Ikemoto  
393 et al., 2004). Despite this mechanism is essential for Hg detoxification, the formation of this  
394 insoluble compound compromises Se biological functions and availability, which might pose a  
395 threat when Se availability is reduced (Ralston and Raymond, 2010).

396 Hg varied widely within this seabird community, with the highest concentrations in  
397 Frigatebirds, Royal tern, and Laughing gull, while Brown noddy, Cayenne tern, and Sooty tern  
398 had lower concentrations (Table 2). Such a difference among species is related to their foraging  
399 ecology. The overall positive association between Hg and  $\delta^{15}\text{N}$  in nestlings (Figure 5) and the  
400 significant positive association between Hg and  $\delta^{15}\text{N}$  in adult Frigatebirds (Table 3, Figure 5)  
401 suggest that Hg is effectively bioaccumulated and biomagnified in this seabird community.  
402 Furthermore, the positive association between Hg and  $\delta^{13}\text{C}$  suggests that variation in Hg  
403 concentrations within this community is also related to the foraging location, as it has been  
404 shown in previous studies (Bearhop et al., 2000; Blévin et al., 2013). However, the association

405 between Hg concentrations and stable isotope values showed a different pattern among species.  
406 In adults, Hg is effectively bioaccumulated only in Frigatebirds, and Hg concentrations  
407 increased with  $\delta^{13}\text{C}$  in all species. The relationship with the carbon stable isotope indicates that  
408 Hg exposure in oceanic habitats is lower than the exposure in costal habitats, likely due to gold-  
409 mining activities. In nestlings, Hg is effectively bioaccumulated in all species, while there was  
410 not association with  $\delta^{13}\text{C}$ .

411 Finally, adult seabirds showed higher Hg levels than nestlings did (Table 2). Since seabirds are  
412 able to excrete Hg in feathers (Dauwe et al., 2003), blood Hg in adults reflects the short-term  
413 Hg contamination of individuals (Fort et al., 2015), and therefore provides information for the  
414 sampling period, usually the breeding season in seabirds (Goodale et al., 2008). In nestlings,  
415 concentrations usually reflect the Hg exposure since hatching and maternal transfer of Hg  
416 (Ackerman et al., 2016), even if the latter can be also excreted in the down.

417

#### 418 *4.3 Persistent organic pollutant exposure*

419 Among environmental contaminants, POPs may act as disruptors of endocrine function and  
420 may stimulate or inhibit the secretion of both reproductive and pituitary hormones (Tartu et al.,  
421 2015a; Verboven et al., 2010; Verreault et al., 2008). POPs are of concern because they are  
422 known to bioaccumulate and biomagnify (Bustnes et al., 2013; Elliott et al., 2015; Mello et al.,  
423 2016). Generally, these concentrations are in agreement with the few studies that have been  
424 recently carried out in this region on sea turtles (De Andres et al., 2016; Guirlet et al., 2010;  
425 Sebastiano et al., 2016).

426

#### 427 *4.4 Trace elements and POPs toxicity*

428 Adult and nestling of the 6 seabird species from French Guiana did not contain As  
429 concentrations exceeding  $50 \mu\text{g g}^{-1} \text{ dw}$ , a commonly used threshold for direct As toxicity in

430 seabirds (Neff, 1997) and should therefore not represent a threat. Furthermore, despite we found  
431 an association between low concentrations of Se and high concentrations of Hg, the molar ratio  
432 Hg:Se was much below 1 in all individuals, indicating that Se is in excess compared to Hg and  
433 thus is not a limiting factor for the detoxification of Hg (Sørmo et al., 2011). Hg concentrations  
434 within the Grand Connétable avian community widely vary among species, and were high as  
435 compared to literature values for tropical regions. However, they are similar to high-  
436 contaminated seabirds in other areas whether from temperate, subpolar, or polar regions  
437 (Supplementary Figure S2). Our study shows, for instance, that average Hg concentration was  
438 much higher in French Guiana compared to another tropical seabird community in Seychelles  
439 archipelago (Catry et al., 2008). Further, despite in this study Hg analyses were performed using  
440 whole blood, which reflects relatively similar concentrations than that of RBCs, Hg  
441 concentrations in the Brown noddy were much lower than those of French Guiana (Catry et al.,  
442 2008). Similar Hg concentrations have been previously associated with deleterious effects  
443 (Costantini et al., 2014; Tartu et al., 2016) with consequences at the population level with  
444 reduction of breeding, hatching and fledging success, and thus lead to population decline  
445 (Goutte et al., 2014a; Goutte et al., 2014b). Finally, as compared to other seabirds exposed to  
446 POPs, our results revealed concentrations of thousand times lower than those associated with  
447 deleterious effects (Erikstad et al., 2013; Goutte et al., 2015), which very likely do not pose a  
448 health threat to this seabird community.

449

## 450 **Conclusions**

451 Our study confirms that POPs are present in very low concentrations in French Guiana seabird  
452 species and likely represent a minor concern. However, it confirms the presence of high levels  
453 of Hg, showing a possible health concern for some species. Further efforts should be made to  
454 investigate the health impact of Hg exposure on these species and the community as a whole.

455 Possibly, the analysis of Hg in other tissues and with other non-destructive tools (e.g. feathers),  
456 might also provide information on Hg exposure at different stages of the breeding period. These  
457 investigations should also clarify whether the negative relationship between high Hg and low  
458 As and Se concentrations might lead to side effects due to their potential deficiency, as it has  
459 been previously shown in captive animals (Fischer et al., 2008; Wang et al., 2009).

460

#### 461 **Acknowledgments**

462 We are grateful to DEAL Guyane, CNRS-funded project *SENTINEL* (O.Chastel), the  
463 University of La Rochelle, the University of Antwerp and FWO (Fonds Wetenschappelijk  
464 Onderzoek), who funded field operations, and contaminant and stable isotope analyses.  
465 Moreover, we are grateful to the GEPOG (Groupe d'Etude et de Protection des Oiseaux en  
466 Guyane) and to the ONCFS (Office national de la Chasse et de la Faune Sauvage), who manage  
467 the Grand Connétable Nature reserve since 2008, for their logistic support and access to the  
468 Grand Connétable Nature Reserve. We are grateful to Grand Connétable Nature staff (A.  
469 Alcide and L. Bétremieux) for their great help in the field and we are also grateful to M. Brault-  
470 Favrou from the platform 'Analyses Élémentaires' (University of La Rochelle - LIENSs) for  
471 her assistance with laboratory work and to G. Guillou and B. Lebreton from the platform  
472 'Analyses Isotopiques' (University of La Rochelle - LIENSs) for their assistance in stable  
473 isotope analysis. The Contrat de Projet Etat Région (CPER 13) is also acknowledged for  
474 funding the AMA and IR-MS. We also thank four anonymous reviewers who provided valuable  
475 comments that helped us to improve the presentation of the work.

476

477 **References**

- 478 Ackerman, J.T., Eagles-Smith, C.A., Herzog, M.P., Hartman, C.A., 2016. Maternal transfer of  
 479 contaminants in birds: Mercury and selenium concentrations in parents and their eggs. *Environ*  
 480 *Pollut* 210, 145-154.
- 481 AMAP, 2015. (The Arctic Monitoring and Assessment Program) Ring Test. Centre de  
 482 Toxicologie, Institute National de Sauté Publique, Québec. Report 2015.
- 483 Bastos, W.R., Dorea, J.G., Bernardi, J.V., Lauthartte, L.C., Mussu, M.H., Lacerda, L.D., Malm,  
 484 O., 2015. Mercury in fish of the Madeira river (temporal and spatial assessment), Brazilian  
 485 Amazon. *Environ Res* 140, 191-197.
- 486 Bearhop, S., Phillips, R.A., Thompson, D.R., Waldron, S., Furness, R.W., 2000. Variability in  
 487 mercury concentrations of great skuas *Catharacta skua*: the influence of colony, diet and trophic  
 488 status inferred from stable isotope signatures. *Marine Ecology Progress Series* 195, 261-268.
- 489 Beyer, N.W., Heinz, G.H., Redmon-Norwood, A.W., 2011. Environmental contaminants in  
 490 wildlife: interpreting tissue concentrations. Second edition. Lewis publisher.
- 491 Blévin, P., Angelier, F., Tartu, S., Ruault, S., Bustamante, P., Herzke, D., Moe, B., Bech, C.,  
 492 Gabrielsen, G.W., Bustnes, J.O., Chastel, O., 2016. Exposure to oxychlordan is associated  
 493 with shorter telomeres in arctic breeding kittiwakes. *Science of The Total Environment* 563–  
 494 564, 125-130.
- 495 Blévin, P., Carravieri, A., Jaeger, A., Chastel, O., Bustamante, P., Cherel, Y., 2013. Wide Range  
 496 of Mercury Contamination in Chicks of Southern Ocean Seabirds. *Plos One* 8, e54508.
- 497 Bugge, J., Barrett, R.T., Pedersen, T., 2011. Optimal foraging in chick-raising Common  
 498 Guillemots (*Uria aalge*). *Journal of Ornithology* 152, 253-259.
- 499 Burger, J., Gochfeld, M., 2001. Metal levels in feathers of cormorants, flamingos and gulls from  
 500 the Coast of Namibia in Southern Africa. *Environmental Monitoring and Assessment* 69, 195-  
 501 203.
- 502 Bustnes, J.O., Bardsen, B.J., Herzke, D., Johnsen, T.V., Eulaers, I., Ballesteros, M., Hanssen,  
 503 S.A., Covaci, A., Jaspers, V.L., Eens, M., Sonne, C., Halley, D., Moum, T., Nost, T.H.,  
 504 Erikstad, K.E., Ims, R.A., 2013. Plasma concentrations of organohalogenated pollutants in  
 505 predatory bird nestlings: associations to growth rate and dietary tracers. *Environ Toxicol Chem*  
 506 32, 2520-2527.
- 507 Catry, T., Ramos, J.A., Le Corre, M., Kojadinovic, J., Bustamante, P., 2008. The role of stable  
 508 isotopes and mercury concentrations to describe seabird foraging ecology in tropical  
 509 environments. *Marine Biology* 155, 637-647.
- 510 Cherel, Y. and Hobson, K.A. 2007. Geographical variation in carbon stable isotope signatures  
 511 of marine predators: a tool to investigate their foraging areas in the Southern Ocean. *Mar Ecol*  
 512 *Prog Ser* 329, 281-287.
- 513 Costantini, D., Meillere, A., Carravieri, A., Lecomte, V., Sorci, G., Faivre, B., Weimerskirch,  
 514 H., Bustamante, P., Labadie, P., Budzinski, H., Chastel, O., 2014. Oxidative stress in relation  
 515 to reproduction, contaminants, gender and age in a long-lived seabird. *Oecologia* 175, 1107-  
 516 1116.
- 517 Dauwe, T., Bervoets, L., Pinxten, R., Blust, R., Eens, M., 2003. Variation of heavy metals  
 518 within and among feathers of birds of prey: effects of molt and external contamination.  
 519 *Environmental Pollution* 124, 429-436.
- 520 De Andres, E., Gomara, B., Gonzalez-Paredes, D., Ruiz-Martin, J., Marco, A., 2016. Persistent  
 521 organic pollutant levels in eggs of leatherback turtles (*Dermochelys coriacea*) point to a  
 522 decrease in hatching success. *Chemosphere* 146, 354-361.
- 523 del Hoyo, J., Collar, N.J., Christie, D.A., Elliott, A., Fishpool, L.D.C., 1996. Handbook of the  
 524 Birds of the World, vol. 3: Hoatzin to Auks. Lynx Edicions, Barcelona, Spain.

525 Dimitriadou, L., Malarvannan, G., Covaci, A., Iossifidou, E., Tzafettas, J., Zournatzi-Koiou,  
526 V., Kalantzi, O.-I., 2016. Levels and profiles of brominated and chlorinated contaminants in  
527 human breast milk from Thessaloniki, Greece. *Science of The Total Environment* 539, 350-  
528 358.

529 Dujardin, J.L., Tostain, O., 1990. Les oiseaux de mer nicheurs de Guyane Francaise. *Alauda*  
530 58, 107-134.

531 Elliott, J.E., Brogan, J., Lee, S.L., Drouillard, K.G., Elliott, K.H., 2015. PBDEs and other POPs  
532 in urban birds of prey partly explained by trophic level and carbon source. *Sci Total Environ*  
533 524-525, 157-165.

534 Erikstad, K.E., Sandvik, H., Reiertsen, T.K., Bustnes, J.O., Strøm, H., 2013. Persistent organic  
535 pollution in a high-Arctic top predator: sex-dependent thresholds in adult survival. *Proceedings*  
536 *of the Royal Society B: Biological Sciences* 280, DOI: 10.1098/rspb.2013.1483.

537 Fischer, J., Bosse, A., Pallauf, J., 2008. Effect of selenium deficiency on the antioxidative status  
538 and muscle damage in growing turkeys. *Arch Anim Nutr* 62, 485-497.

539 Fitzgerald, W.F., Lamborg, C.H., Hammerschmidt, C.R., 2007. Marine Biogeochemical  
540 Cycling of Mercury. *Chemical Reviews* 107, 641-662.

541 Fort, J., Lacoue-Labarthe, T., Nguyen, H.L., Boué, A., Spitz, J., Bustamante, P., 2015. Mercury  
542 in wintering seabirds, an aggravating factor to winter wrecks? *Science of The Total*  
543 *Environment* 527-528, 448-454.

544 Frery, N., Maury-Brachet, R., Maillot, E., Deheeger, M., de Merona, B., Boudou, A., 2001.  
545 Gold-mining activities and mercury contamination of native amerindian communities in French  
546 Guiana: key role of fish in dietary uptake. *Environ Health Perspect* 109, 449-456.

547 Fujimura, M., Matsuyama, A., Harvard, J.P., Bourdineaud, J.P., Nakamura, K., 2012. Mercury  
548 contamination in humans in Upper Maroni, French Guiana between 2004 and 2009. *Bull*  
549 *Environ Contam Toxicol* 88, 135-139.

550 Furness, R.W., Camphuysen, K., 1997. Seabirds as monitors of the marine environment. *ICES*  
551 *Journal of Marine Science: Journal du Conseil* 54, 726-737.

552 Goodale, M.W., Evers, D.C., Mierzykowski, S.E., Bond, A.L., Burgess, N.M., Otorowski, C.I.,  
553 Welch, L.J., Hall, C.S., Ellis, J.C., Allen, R.B., Diamond, A.W., Kress, S.W., Taylor, R.J., 2008.  
554 Marine foraging birds as bioindicators of mercury in the Gulf of Maine. *Ecohealth* 5, 409-425.

555 Goutte, A., Barbraud, C., Herzke, D., Bustamante, P., Angelier, F., Tartu, S., Clement-Chastel,  
556 C., Moe, B., Bech, C., Gabrielsen, G.W., Bustnes, J.O., Chastel, O., 2015. Survival rate and  
557 breeding outputs in a high Arctic seabird exposed to legacy persistent organic pollutants and  
558 mercury. *Environmental Pollution* 200, 1-9.

559 Goutte, A., Barbraud, C., Meillere, A., Carravieri, A., Bustamante, P., Labadie, P., Budzinski,  
560 H., Delord, K., Cherel, Y., Weimerskirch, H., Chastel, O., 2014a. Demographic consequences  
561 of heavy metals and persistent organic pollutants in a vulnerable long-lived bird, the wandering  
562 albatross. *Proceedings of the Royal Society B-Biological Sciences* 281, DOI:  
563 10.1098/rspb.2013.3313.

564 Goutte, A., Bustamante, P., Barbraud, C., Delord, K., Weimerskirch, H., Chastel, O., 2014b.  
565 Demographic responses to mercury exposure in two closely related Antarctic top predators.  
566 *Ecology* 95, 1075-1086.

567 Guirlet, E., Das, K., Thome, J.P., Girondot, M., 2010. Maternal transfer of chlorinated  
568 contaminants in the leatherback turtles, *Dermochelys coriacea*, nesting in French Guiana.  
569 *Chemosphere* 79, 720-726.

570 Hobson, K.A., Clark, R.G., 1993. Turnover of <sup>13</sup>C in cellular and plasma fractions of blood:  
571 implications for nondestructive sampling in avian dietary studies. *The Auk* 110, 638-641.

572 Ikemoto, T., Kunito, T., Tanaka, H., Baba, N., Miyazaki, N., Tanabe, S., 2004. Detoxification  
573 mechanism of heavy metals in marine mammals and seabirds: interaction of selenium with

574 mercury, silver, copper, zinc, and cadmium in liver. *Arch Environ Contam Toxicol* 47, 402-  
575 413.

576 Jackson, A.L., Inger, R., Parnell, A.C., Bearhop, S., 2011. Comparing isotopic niche widths  
577 among and within communities: SIBER - Stable Isotope Bayesian Ellipses in R. *J Anim Ecol*  
578 80, 595-602.

579 Kelly, J.F. 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian  
580 trophic ecology. *Can. J. Zool.* 78, 1-27.

581 Lodenius, M., Malm, O., 1998. Mercury in the Amazon. *Rev Environ Contam Toxicol* 157, 25-  
582 52.

583 Lu, S.C., 2000. S-Adenosylmethionine. *Int J Biochem Cell Biol* 32, 391-395.

584 Martinet, V., Blanchard, F., 2009. Fishery externalities and biodiversity: Trade-offs between  
585 the viability of shrimp trawling and the conservation of Frigatebirds in French Guiana.  
586 *Ecological Economics* 68, 2960-2968.

587 Mello, F.V., Roscales, J.L., Guida, Y.S., Menezes, J.F., Vicente, A., Costa, E.S., Jimenez, B.,  
588 Torres, J.P., 2016. Relationship between legacy and emerging organic pollutants in Antarctic  
589 seabirds and their foraging ecology as shown by  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ . *Sci Total Environ* 573,  
590 1380-1389.

591 Moreno, R., Jover, L., Diez, C., Sanpera, C., 2011. Seabird feathers as monitors of the levels  
592 and persistence of heavy metal pollution after the Prestige oil spill. *Environ Pollut* 159, 2454-  
593 2460.

594 Morrison, K.W., Bury, S.J., Thompson, D.R., 2014. Higher trophic level prey does not represent  
595 a higher quality diet in a threatened seabird: implications for relating population dynamics to  
596 diet shifts inferred from stable isotopes. *Marine Biology* 161, 2243-2255.

597 Neff, J.M., 1997. Ecotoxicology of arsenic in the marine environment. *Environmental*  
598 *Toxicology and Chemistry* 16, 917-927.

599 Newsome, S.D., Martinez del Rio, C., Bearhop, S., Phillips, D.L., 2007. A niche for isotopic  
600 ecology. *Frontiers in Ecology and the Environment* 5, 429-436.

601 Nielsen, F.H., 1998. Ultratrace elements in nutrition: Current knowledge and speculation. *The*  
602 *Journal of Trace Elements in Experimental Medicine* 11, 251-274.

603 Overman, N.C. and Parrish, D.L. 2001. Stable isotope composition of walleye:  $\delta^{15}\text{N}$   
604 accumulation with age and area specific differences in  $\delta^{13}\text{C}$ . *Can. J. Fish. Aquat. Sci.* 58,  
605 1253-1260.

606 Polak-Juszczak, L., Robak, S., 2015. Mercury toxicity and the protective role of selenium in  
607 eel, *Anguilla anguilla*. *Environ Sci Pollut Res Int* 22, 679-688.

608 Ralston, N.V.C., Raymond, L.J., 2010. Dietary selenium's protective effects against  
609 methylmercury toxicity. *Toxicology* 278, 112-123.

610 Rayman, M.P., 2012. Selenium and human health. *The Lancet* 379, 1256-1268.

611 Rowe, C.L., 2008. "The Calamity of So Long Life": Life Histories, Contaminants, and Potential  
612 Emerging Threats to Long-lived Vertebrates. *BioScience* 58, 623-631.

613 Sebastiano, M., Bustamante, P., Costantini, D., Eulaers, I., Malarvannan, G., Mendez-  
614 Fernandez, P., Churlaud, C., Blevin, P., Hauselmann, A., Dell'Omo, G., Covaci, A., Eens, M.,  
615 Chastel, O., 2016. High levels of mercury and low levels of persistent organic pollutants in a  
616 tropical seabird in French Guiana, the Magnificent frigatebird, *Fregata magnificens*. *Environ*  
617 *Pollut* 214, 384-393.

618 Sørmo, E.G., Ciesielski, T.M., Øverjordet, I.B., Lierhagen, S., Eggen, G.S., Berg, T., Jenssen,  
619 B.M., 2011. Selenium Moderates Mercury Toxicity in Free-Ranging Freshwater Fish. *Environ*  
620 *Sci Technol* 45, 6561-6566.

621 Tapiero, H., Townsend, D.M., Tew, K.D., 2003. The antioxidant role of selenium and seleno-  
622 compounds. *Biomed Pharmacother* 57, 134-144.



623 Tartu, S., Angelier, F., Bustnes, J.O., Moe, B., Hanssen, S.A., Herzke, D., Gabrielsen, G.W.,  
624 Verboven, N., Verreault, J., Labadie, P., Budzinski, H., Wingfield, J.C., Chastel, O., 2015b.  
625 Polychlorinated biphenyl exposure and corticosterone levels in seven polar seabird species.  
626 *Environ Pollut* 197, 173-180.

627 Tartu, S., Angelier, F., Wingfield, J.C., Bustamante, P., Labadie, P., Budzinski, H.,  
628 Weimerskirch, H., Bustnes, J.O., Chastel, O., 2015a. Corticosterone, prolactin and egg neglect  
629 behavior in relation to mercury and legacy POPs in a long-lived Antarctic bird. *Science of The*  
630 *Total Environment* 505, 180-188.

631 Tartu, S., Lendvai, A.Z., Blevin, P., Herzke, D., Bustamante, P., Moe, B., Gabrielsen, G.W.,  
632 Bustnes, J.O., Chastel, O., 2015c. Increased adrenal responsiveness and delayed hatching date  
633 in relation to polychlorinated biphenyl exposure in Arctic-breeding black-legged kittiwakes  
634 (*Rissa tridactyla*). *General and Comparative Endocrinology* 219, 165-172.

635 Tartu, S., Bustamante, P., Angelier, F., Lendvai, Á.Z., Moe, B., Blévin, P., Bech, C.,  
636 Gabrielsen, G.W., Bustnes, J.O., Chastel, O., Hőrak, P., 2016. Mercury exposure, stress and  
637 prolactin secretion in an Arctic seabird: an experimental study. *Functional Ecology* 30, 596-  
638 604.

639 Tinggi, U., 2008. Selenium: its role as antioxidant in human health. *Environmental Health and*  
640 *Preventive Medicine* 13, 102-108.

641 Uthus, E.O., 2003. Arsenic essentiality: A role affecting methionine metabolism. *The Journal*  
642 *of Trace Elements in Experimental Medicine* 16, 345-355.

643 Verboven, N., Verreault, J., Letcher, R.J., Gabrielsen, G.W., Evans, N.P., 2010. Adrenocortical  
644 function of Arctic-breeding glaucous gulls in relation to persistent organic pollutants. *Gen*  
645 *Comp Endocrinol* 166, 25-32.

646 Verreault, J., Verboven, N., Gabrielsen, G.W., Letcher, R.J., Chastel, O., 2008. Changes in  
647 prolactin in a highly organohalogen contaminated Arctic top predator seabird, the glaucous gull.  
648 *Gen Comp Endocrinol* 156, 569-576.


649 Wang, C., Wang, H., Luo, J., Hu, Y., Wei, L., Duan, M., He, H., 2009. Selenium deficiency  
650 impairs host innate immune response and induces susceptibility to *Listeria monocytogenes*  
651 infection. *BMC Immunology* 10, 1-12.

652 Weimerskirch, H., Chastel, O., Barbraud, C., Tostain, O., 2003. Flight performance:  
653 Frigatebirds ride high on thermals. *Nature* 421, 333-334.

654

655 **Figures**

656




657

658 **Figure 1:** Individual  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values (‰) in adults (left) and nestlings (right) of the six

659 seabird species from the Grand Connétable Island, French Guiana.

660




661

662


663 **Figure 2:** Density plots of the SEA based on  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values (‰) in adults (left) and  
 664 nestlings (right) of the six seabird species from the Grand Connétable Island, French Guiana.  
 665 Black dots represent their mode, and the shaded boxes representing 50, 75 and 95% credible  
 666 intervals from dark to light grey, respectively.

667




668

669 **Figure 3:** Interspecific comparison for adult and nestling trace element concentrations ( $\mu\text{g g}^{-1}$   
670 dw) in red blood cells of the six seabird species from the Grand Connétable Island, French  
671 Guiana. Values are mean  $\pm$  SD. Species (arranged along increasing adult Hg levels) sharing the  
672 same letter have burdens that are not significantly different (Tukey HSD,  $P > 0.05$ ). Nestlings'  
673 values of Hg and Se, and adults' values of As, Hg, Mn, Se, and Zn were  $\log_{10}$ -transformed to  
674 achieve the requested normality for statistical tests. Species abbreviation: CT = Cayenne tern;  
675 LG = Laughing gull; BN = Brown noddy; FB = Magnificent frigatebird; RT = Royal tern; ST  
676 = Sooty tern.



677  
 678 **Figure 4:** Principal component analysis of trace element exposure in adult (a) and nestling (b)  
 679 of the six seabird species from the Grand Connétable Island, French Guiana. The graphs on the  
 680 left show species-specific PC values. On the right, each arrow represents a contaminant with  
 681 the direction representing where the contaminants load in the principal component space. The  
 682 length of the arrow represents the magnitude of the loading.  
 683

684



685

686 **Figure 5:** Relationship between Hg concentrations ( $\mu\text{g g}^{-1} \text{dw}$ ) and  $\delta^{15}\text{N}$  (‰) in adults and  
687 nestlings of the six seabird species from the Grand Connétable Island, French Guiana.

688

689 **Table 1.** Pearson's correlations between  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  in adults and nestlings of the six seabird  
690 species from the Grand Connétable Island, French Guiana. Significant correlation values are  
691 bolded. ND refers to the inability to carry out statistical testing due to a low sample size.

692

Species	Adults			Nestlings		
	<i>n</i>	<i>r</i>	<i>P</i>	<i>n</i>	<i>r</i>	<i>P</i>
Cayenne tern	20	0.63	<b>&lt;0.01</b>	20	0.75	<b>&lt;0.01</b>
Laughing gull	20	-0.52	<b>0.02</b>	20	-0.15	0.54
Brown noddy	20	0.43	0.06	20	0.71	<b>&lt;0.01</b>
Frigatebirds	20	0.48	<b>0.03</b>	20	0.43	0.06
Royal tern	15	0.09	0.74	20	0.01	0.96
Sooty tern	6	0.96	<b>&lt;0.01</b>	2	ND	ND
Overall	101	0.54	<b>&lt;0.01</b>	102	0.88	<b>&lt;0.01</b>

693

694

695 **Table 2:** Red blood cell trace element concentrations ( $\mu\text{g g}^{-1}$  dw) in adults and nestlings of the  
696 six seabird species from the Grand Connétable Island, French Guiana. For each age class, first  
697 row values are mean  $\pm$  SD and second row values represent the range (median). Detection  
698 frequencies were lower than 100% in the Ni content in Laughing gull nestlings (65%) and  
699 Brown noddy nestlings (55%). ND refers to non-detects. Frigatebirds data are earlier reported  
700 by (Sebastiano et al., 2016).

		<b>Cayenne tern</b>	<b>Laughing gull</b>	<b>Brown noddy</b>	<b>Frigatebirds</b>	<b>Royal tern</b>	<b>Sooty tern</b>
<b>As</b>	adults	8.7 $\pm$ 4.4	1.5 $\pm$ 1.0	8.4 $\pm$ 3.9	2.4 $\pm$ 1.4	3.9 $\pm$ 2.6	10.7 $\pm$ 4.9
		2.0-20.8 (9.1)	0.5-4.3 (1.1)	2.5-16.7 (7.8)	0.6-7.3 (2.1)	0.7-9.1 (3.0)	4.6-19.0 (9.2)
	nestlings	4.6 $\pm$ 1.2	2.7 $\pm$ 0.8	1.9 $\pm$ 0.5	1.6 $\pm$ 0.7	3.6 $\pm$ 1.0	2.3 $\pm$ 0.6
		2.2-6.2 (4.9)	1.3-3.9 (2.7)	1.1-2.8 (1.9)	0.7-3.6 (1.5)	2.0-6.3 (3.5)	1.9-2.7 (2.3)
<b>Cd</b>	adults	ND	ND	0.051 $\pm$ 0.021	ND	ND	0.026 $\pm$ 0.014
				0.034-0.118 (0.044)			0.015-0.050
	nestlings	ND	ND	0.015 $\pm$ 0.004	ND	ND	ND
				0.010-0.029 (0.015)			
<b>Cu</b>	adults	0.9 $\pm$ 0.1	1.3 $\pm$ 0.2	0.8 $\pm$ 0.1	0.8 $\pm$ 0.1	1.1 $\pm$ 0.1	1.1 $\pm$ 0.1
		0.8-1.0 (0.9)	1.0-1.9 (1.3)	0.6-0.9 (0.8)	0.7-0.9 (0.8)	0.9-1.3 (1.1)	1.0-1.1 (1.1)
	nestlings	0.8 $\pm$ 0.1	1.3 $\pm$ 0.1	0.8 $\pm$ 0.1	0.7 $\pm$ 0.1	1.1 $\pm$ 0.1	1.1 $\pm$ <0.1
		0.7-1.0 (0.8)	1.1-1.5 (1.3)	0.7-1.2 (0.8)	0.6-0.9 (0.7)	0.9-1.3 (1.1)	1.0-1.1 (1.1)
<b>Fe</b>	adults	2368 $\pm$ 123	2348 $\pm$ 65	2382 $\pm$ 54	2413 $\pm$ 68	2388 $\pm$ 71	2444 $\pm$ 78
		2000-2528 (2397)	2197-2432 (2363)	2269-2485 (2385)	2235-2503 (2411)	2226-2475 (2410)	2348-2582 (2441)
	nestlings	2207 $\pm$ 58	2337 $\pm$ 64	2171 $\pm$ 78	2330 $\pm$ 80	2133 $\pm$ 56	2151 $\pm$ 34
		2113-2326 (2205)	2178-2423 (2339)	2029-2302 (2164)	2146-2477 (2337)	2050-2286 (2123)	2127-2175 (2151)
<b>Hg</b>	adults	1.1 $\pm$ 0.3	2.5 $\pm$ 1.4	1.1 $\pm$ 0.1	5.8 $\pm$ 1.3	2.6 $\pm$ 0.7	0.9 $\pm$ 0.2
		0.6-1.7 (1.0)	0.5-5.8 (2.3)	0.9-1.4 (1.1)	3.8-7.8 (5.6)	1.5-3.8 (2.8)	0.6-1.1 (0.9)
	nestlings	<0.1 $\pm$ 0.1	0.4 $\pm$ 0.1	0.2 $\pm$ <0.1	1.0 $\pm$ 0.2	0.3 $\pm$ 0.1	0.3 $\pm$ <0.1
		0.1-0.3 (0.2)	0.2-0.6 (0.3)	0.1-0.2 (0.1)	0.7-1.7 (1.0)	0.2-0.4 (0.3)	0.2-0.2 (0.2)
<b>Mn</b>	adults	0.1 $\pm$ <0.1	0.1 $\pm$ <0.1	0.1 $\pm$ <0.1	0.1 $\pm$ <0.1	0.2 $\pm$ 0.1	0.1 $\pm$ <0.1
		0.1-0.2 (0.1)	0.1-0.2 (0.1)	0.1-0.1 (0.1)	0.1-0.2 (0.1)	0.1-0.3 (0.2)	0.2-0.2 (0.1)
	nestlings	0.1 $\pm$ <0.1	0.3 $\pm$ 0.1	0.1 $\pm$ <0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	0.3 $\pm$ <0.1
		0.1-0.2 (0.1)	0.2-0.4 (0.3)	0.1-0.1 (0.1)	0.1-0.3 (0.2)	0.1-0.4 (0.2)	0.3-0.3 (0.3)
<b>Ni</b>	adults	ND	ND	ND	ND	ND	ND
	nestlings	ND	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	ND	ND	ND
			<0.1-0.2 (0.1)	<0.1-0.3 (0.1)			
<b>Pb</b>	adults	0.02 $\pm$ <0.01	0.06 $\pm$ 0.04	0.02 $\pm$ 0.01	0.02 $\pm$ 0.01	0.03 $\pm$ 0.03	0.02 $\pm$ 0.00
		0.01-0.06 (0.02)	0.02-0.20 (0.04)	0.01-0.05 (0.01)	0.02-0.04 (0.02)	0.02-0.11 (0.02)	0.01-0.03 (0.01)
	nestlings	0.02 $\pm$ <0.01	0.04 $\pm$ 0.03	0.02 $\pm$ 0.00	0.02 $\pm$ 0.00	0.02 $\pm$ 0.01	0.02 $\pm$ 0.01
		0.01-0.06 (0.02)	0.02-0.11 (0.03)	0.01-0.03 (0.02)	0.01-0.03 (0.02)	0.01-0.05 (0.02)	0.01-0.02 (0.02)
<b>Se</b>	adults	41.6 $\pm$ 7.0	10.7 $\pm$ 3.8	72.7 $\pm$ 18.6	9.1 $\pm$ 1.9	28.8 $\pm$ 13.2	160.1 $\pm$ 54.1
		26.3-52.7 (41.7)	4.3-17.9 (10.3)	42.5-103.0 (71.7)	6.7-13.1 (8.7)	9.1-62.9 (27.9)	62.2-215.6 (173.1)
	nestlings	30.3 $\pm$ 4.6	7.4 $\pm$ 2.7	19.6 $\pm$ 3.9	5.8 $\pm$ 0.6	8.9 $\pm$ 2.9	8.4 $\pm$ 0.2
		22.2-39.0 (29.5)	5.0-15.6 (6.8)	12.4-27.7 (19.4)	4.6-6.6 (5.8)	4.8-16.8 (8.4)	8.3-8.5 (8.4)
<b>Zn</b>	adults	21.0 $\pm$ 4.3	22.4 $\pm$ 1.6	20.7 $\pm$ 1.1	19.4 $\pm$ 0.9	20.7 $\pm$ 1.7	20.0 $\pm$ 1.8
		17.0-37.4 (20.0)	18.5-24.4 (22.7)	18.9-22.8 (20.6)	18.3-22.1 (19.4)	19.0-24.8 (20.2)	18.5-23.1 (19.3)
	nestlings	22.4 $\pm$ 1.7	26.0 $\pm$ 1.4	25.3 $\pm$ 1.7	26.9 $\pm$ 3.0	23.5 $\pm$ 2.5	21.2 $\pm$ 0.4
		19.5-26.5 (22.1)	22.4-27.8 (26.1)	22.2-28.1 (25.1)	22.5-32.6 (26.8)	20.4-27.5 (22.9)	20.9-21.5 (21.2)



702 **Table 3:** Linear models and linear mixed models (for overall comparison only) explaining the  
 703 association between mercury concentration and stable isotopes and between persistent organic  
 704 pollutants and stable isotopes.

705



			Adults				Nestlings			
			Slope (SE)	<i>t</i> -value	<i>df</i>	<i>P</i>	Slope (SE)	<i>t</i> -value	<i>df</i>	<i>P</i>
<b>Hg &amp; <math>\delta^{15}\text{N}</math></b>	<b>Overall</b>	(intercept)	-0.77 (0.85)	-0.90	86.6	0.37	-4.78 (0.86)	-5.57	94.1	<0.01
		$\delta^{15}\text{N}$	0.11 (0.07)	1.65	98.8	0.10	0.29 (0.07)	4.27	100	<0.01
	<b>Among species</b>	(intercept)	-0.54 (0.82)	0.82	94	0.51	-4.39 (0.85)	-5.14	95	<0.01
		Cayenne tern	0.05 (0.07)	0.70	94	0.49	0.23 (0.07)	3.21	95	<0.01
		Laughing gull	0.11 (0.07)	1.58	94	0.12	0.26 (0.07)	3.90	95	<0.01
		Brown noddy	0.05 (0.07)	0.80	94	0.43	0.22 (0.07)	2.89	95	<0.01
		Frigatebird	0.17 (0.06)	2.78	94	<0.01	0.33 (0.06)	5.11	95	<0.01
		Royal tern	0.12 (0.06)	1.80	94	0.08	0.25 (0.07)	3.68	95	<0.01
		Sooty tern	0.03 (0.08)	0.44	94	0.66	0.26 (0.07)	3.46	95	<0.01
<b>Hg &amp; <math>\delta^{13}\text{C}</math></b>	<b>Overall</b>	(intercept)	8.79 (1.27)	6.91	90.3	<0.01	1.79 (1.49)	1.21	88.6	0.23
		$\delta^{13}\text{C}$	0.52 (0.08)	6.52	94.2	<0.01	0.19 (0.09)	2.05	93.5	<0.05
	<b>Among species</b>	(intercept)	8.13 (1.29)	6.29	94	<0.01	0.52 (1.60)	0.33	95	0.75
		Cayenne tern	0.50 (0.08)	6.26	94	<0.01	0.13 (0.10)	1.35	95	0.13
		Laughing gull	0.48 (0.08)	5.70	94	<0.01	0.10 (0.10)	0.99	95	0.18
		Brown noddy	0.50 (0.08)	6.19	94	<0.01	0.14 (0.09)	1.53	95	0.73
		Frigatebird	0.43 (0.09)	4.94	94	<0.01	0.04 (0.11)	0.34	95	0.32
		Royal tern	0.47 (0.08)	5.56	94	<0.01	0.11 (0.10)	1.10	95	0.27
		Sooty tern	0.49 (0.08)	6.41	94	<0.01	0.12 (0.10)	1.19	95	0.24
$\Sigma\text{PCBs & } \delta^{15}\text{N}$	<b>Overall</b>	(intercept)	2.11 (4.04)	0.52	50.1	0.60	ND	ND	ND	ND
		$\delta^{15}\text{N}$	0.28 (0.32)	0.88	54.4	0.39				
$\Sigma\text{PCBs & } \delta^{13}\text{C}$	<b>Overall</b>	(intercept)	12.62 (10.67)	1.2	19.9	0.25	ND	ND	ND	ND
		$\delta^{13}\text{C}$	0.45 (0.68)	0.66	20.2	0.52				
<i>p,p'</i> -DDE $\delta^{15}\text{N}$	<b>Overall</b>	(intercept)	3.77 (6.92)	0.54	39.6	0.59	ND	ND	ND	ND
		$\delta^{15}\text{N}$	0.14 (0.55)	0.26	42.3	0.80				
<i>p,p'</i> -DDE $\delta^{13}\text{C}$	<b>Overall</b>	(intercept)	-34.38 (18.2)	-1.89	18.0	0.08	ND	ND	ND	ND
		$\delta^{13}\text{C}$	-2.54 (1.16)	-2.20	18.35	<0.05				



706

707 **Table 4:** Values of plasma levels of POPs across species and age classes of the three seabird  
708 species from the Grand Connétable Island, French Guiana. For each age class, first row values  
709 are mean  $\pm$  SD and second row values represent the range (median). Concentrations are  
710 expressed as  $\text{pg g}^{-1}$  wet weight. ND refers to non-detects. Frigatebirds data are earlier reported  
711 by (Sebastiano et al., 2016).  
712







	Adults			Nestlings		
	Cayenne tern	Brown noddy	Frigatebirds	Cayenne tern	Brown noddy	Frigatebirds
$\Sigma$ PCBs	510 $\pm$ 470	91 $\pm$ 68	920 $\pm$ 640	30 $\pm$ 30	ND	44 $\pm$ 39
	67-2200 (370)	5-280 (79)	330-2700 (640)	3-110 (15)		14-200 (35)
$\Sigma$ CHLs	11 $\pm$ 9	ND	10 $\pm$ 3	ND	ND	4 $\pm$ 7
	4-37 (8)		5-16 (11)			2-32 (3)
HCB	150 $\pm$ 140	ND	12 $\pm$ 11	ND	ND	11 $\pm$ 6
	3-480 (100)		2-41 (7)			2-33 (11)
<i>p,p'</i> -DDE	1300 $\pm$ 1100	200 $\pm$ 200	430 $\pm$ 560	ND	ND	40 $\pm$ 45
	160-5100 (1100)	5-600 (200)	75-2300 (220)			13-210 (25)
HCHs	34 $\pm$ 27	13 $\pm$ 13	ND	20 $\pm$ 14	11 $\pm$ 8	11 $\pm$ 7
	5-120 (29)	3-45 (8)		3-55 (17)	3-26 (9)	2-20 (12)
$\Sigma$ PBDEs	ND	ND	ND	ND	ND	ND

713













### Adults





### Nestlings





### a) Adults



### b) Nestlings





# **Trophic ecology drives contaminant concentrations within a tropical seabird community**

**Manrico Sebastiano<sup>1\*</sup>, Paco Bustamante<sup>2</sup>, Igor Eulaers<sup>3</sup>, Govindan Malarvannan<sup>4</sup>, Paula Mendez-Fernandez<sup>2</sup>, Carine Churlaud<sup>2</sup>, Pierre Blévin<sup>5</sup>, Antoine Hauselmann<sup>6</sup>, Adrian Covaci<sup>4</sup>, Marcel Eens<sup>1</sup>, David Costantini<sup>1,7</sup>, Olivier Chastel<sup>5</sup>**

1. Behavioural Ecology & Ecophysiology group, Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
2. Littoral Environnement et Sociétés, UMR 7266 CNRS-Université La Rochelle, 2 rue Olympe de Gouges, 17000 La Rochelle, France
3. Department of Bioscience, Aarhus University, Frederiksborgvej 399, PO Box 358, 4000, Roskilde, Denmark
4. Toxicological Centre, Department of Pharmaceutical Sciences, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
5. Centre d'Etudes Biologiques de Chizé (CEBC), UMR7372, CNRS-Université de La Rochelle, F-79360, France
6. Association GEPOG, 15 Av Louis Pasteur, 97300 Cayenne, French Guiana
7. UMR 7221, Muséum National d'Histoire Naturelle, 7 rue Cuvier 75231 Paris cedex 05, France

\* Corresponding author: [Manrico.Sebastiano@uantwerpen.be](mailto:Manrico.Sebastiano@uantwerpen.be)



## **Highlights**

- POPs and trace elements were studied in a tropical seabird community.
- Trace element concentrations widely vary among the six investigated species.
- Hg concentrations were strongly associated with carbon and nitrogen isotopes.
- POPs were generally low in all three species analysed.
- Hg impact on these species and the community as a whole should be investigated.

## Supplementary Information

### **Trophic ecology drives contaminant concentrations within a tropical seabird community**

**Manrico Sebastiano<sup>1\*</sup>, Paco Bustamante<sup>2</sup>, Igor Eulaers<sup>3</sup>, Govindan Malarvannan<sup>4</sup>, Paula Mendez-Fernandez<sup>2</sup>, Carine Churlaud<sup>2</sup>, Pierre Blévin<sup>5</sup>, Antoine Hauselmann<sup>6</sup>, Adrian Covaci<sup>4</sup>, Marcel Eens<sup>1</sup>, David Costantini<sup>1,7</sup>, Olivier Chastel<sup>5</sup>**


1. Behavioural Ecology & Ecophysiology group, Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
2. Littoral Environnement et Sociétés, UMR 7266 CNRS-Université La Rochelle, 2 rue Olympe de Gouges, 17000 La Rochelle, France
3. Department of Bioscience, Aarhus University, Frederiksborgvej 399, PO Box 358, 4000, Roskilde, Denmark
4. Toxicological Centre, Department of Pharmaceutical Sciences, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
5. Centre d'Etudes Biologiques de Chizé (CEBC), UMR7372, CNRS-Université de La Rochelle, F-79360, France
6. Association GEPOG, 15 Av Louis Pasteur, 97300 Cayenne, French Guiana
7. UMR 7221, Muséum National d'Histoire Naturelle, 7 rue Cuvier 75231 Paris cedex 05, France

\* Corresponding author: [Manrico.Sebastiano@uantwerpen.be](mailto:Manrico.Sebastiano@uantwerpen.be)


**Supplementary Table S1:**  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values (‰) in both adults and nestlings of the six seabird species from the Grand Connétable Island, French Guiana.

<b>Species</b>	<b>Mean <math>\pm</math> SD</b>	<b>Median</b>	<b>Range</b>	<b>Mean <math>\pm</math> SD</b>	<b>Median</b>	<b>Range</b>
	<b>Adults: <math>\delta^{13}\text{C}</math></b>			<b>Nestlings: <math>\delta^{13}\text{C}</math></b>		
Cayenne tern	-16.26 $\pm$ 0.15	-16.21	-16.53, -16.07	-16.62 $\pm$ 0.14	-16.64	-16.82, -16.36
Laughing gull	-15.30 $\pm$ 0.67	-15.38	-16.57, -14.01	-15.75 $\pm$ 0.25	-15.70	-16.43, -15.29
Brown noddy	-15.94 $\pm$ 0.12	-15.95	-16.16, -15.69	-17.59 $\pm$ 0.40	-17.61	-18.53, -16.90
Frigatebird	-15.01 $\pm$ 0.11	-14.99	-15.30, -14.82	-15.19 $\pm$ 0.09	-15.20	-15.34, -15.00
Royal tern	-15.36 $\pm$ 0.35	-15.41	-15.88, -14.65	-15.84 $\pm$ 0.21	-15.87	-16.12, -15.48
Sooty tern	-17.04 $\pm$ 0.31	-17.12	-17.38, -16.48	-16.29 $\pm$ 0.09	-16.29	-16.36, -16.23
	<b>Adults: <math>\delta^{15}\text{N}</math></b>			<b>Nestlings: <math>\delta^{15}\text{N}</math></b>		
Cayenne tern	11.98 $\pm$ 0.45	12.05	11.02, 12.66	11.83 $\pm$ 0.23	11.75	11.53, 12.32
Laughing gull	12.26 $\pm$ 0.86	12.18	11.56, 13.52	12.96 $\pm$ 0.41	13.01	12.05, 13.53
Brown noddy	12.30 $\pm$ 0.19	12.32	11.95, 12.62	11.43 $\pm$ 0.32	11.42	10.70, 11.91
Frigatebird	13.37 $\pm$ 0.20	13.37	13.04, 13.77	13.41 $\pm$ 0.28	13.33	13.07, 14.17
Royal tern	12.71 $\pm$ 0.26	12.78	12.23, 13.03	12.50 $\pm$ 0.31	12.60	11.74, 12.87
Sooty tern	10.71 $\pm$ 0.86	10.63	9.61, 12.27	12.68 $\pm$ 0.21	12.68	12.53, 12.83

**Figure S1:** Box plot of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values (‰) in both adults (white plots) and nestlings (blue plots) of the six seabird species from the Grand Connétable Island, French Guiana. Within age classes, species with non-significant differences in carbon or nitrogen isotope values share the same letter. Significant differences between adults and nestlings are indicated by \* next to the name of the species.



**Figure S2:** Comparison of blood Hg concentrations and SD expressed as  $\mu\text{g g}^{-1}$  dw in different seabird species. Black and grey histograms refer to adults and nestlings, respectively. Species abbreviations: *AS*=*Anous stolidus*; *AT*= *Anous tenuirostris*; *DE*=*Diomedea exulans*; *FM*=*Fregata magnificens*; *GA*= *Gygis alba*; *HC*=*Hydroprogne caspia*; *HL*=*Halobaena caerulea*; *HM*=*Himantopus mexicanus*; *LA*=*Leucophaeus atricilla*; *MG*=*Macronectes giganteus*; *MH*=*Macronectes halli*; *OF*=*Onychoprion fuscatus*; *PA*=*Procellaria aequinoctialis*; *PD*=*Pachyptila desolata*; *PG*=*Pelacanoides georgicus*; *PL*=*Puffinus lherminieri*; *PLE*= *Phaethon lepturus*; *PN*=*Pagodroma nivea*; *PU*=*Pelacanoides urinatrix*; *RA*=*Recurvirostra americana*; *SA*=*Stercorarius antarcticus*; *SF*=*Sterna forsteri*; *SM*=*Stercorarius maccormicki*; *TC*=*Thalassarche chrysostoma*; *TM*=*Thalassarche melanophrys*; *TS*=*Thalasseus sandvicensis*; *TX*=*Thalasseus maximus*. References: San Francisco Bay (Eagles-Smith et al., 2008); Grand Connétable (this study); Seychelles Archipelago (Catry et al., 2008); Kerguelen Archipelago (Fromant et al., 2016; Goutte et al., 2014); Bird Island (Anderson et al., 2009); Adélie Land (Goutte et al., 2014; Tartu et al., 2014). Measurements were carried out in whole blood for all species in San Francisco Bay, Seychelles, Bird Island, and in the Antarctic prion in Kerguelen Archipelago. All other measurements were made in red blood cells. Hg concentration in Seychelles archipelago were extrapolated from graphs (Catry et al., 2008).



## References

- Anderson, O.R.J., Phillips, R.A., McDonald, R.A., Shore, R.F., McGill, R.A.R., Bearhop, S., 2009. Influence of trophic position and foraging range on mercury levels within a seabird community. *Marine Ecology Progress Series* 375, 277-288.
- Catry, T., Ramos, J.A., Le Corre, M., Kojadinovic, J., Bustamante, P., 2008. The role of stable isotopes and mercury concentrations to describe seabird foraging ecology in tropical environments. *Marine Biology* 155, 637-647.
- Eagles-Smith, C.A., Ackerman, J.T., Adelsbach, T.L., Takekawa, J.Y., Miles, A.K., Keister, R.A., 2008. Mercury correlations among six tissues for four waterbird species breeding in San Francisco Bay, California, USA. *Environmental Toxicology and Chemistry* 27, 2136-2153.
- Fromant, A., Carravieri, A., Bustamante, P., Labadie, P., Budzinski, H., Peluhet, L., Churlaud, C., Chastel, O., Cherel, Y., 2016. Wide range of metallic and organic contaminants in various tissues of the Antarctic prion, a planktonophagous seabird from the Southern Ocean. *Sci Total Environ* 544, 754-764.
- Goutte, A., Bustamante, P., Barbraud, C., Delord, K., Weimerskirch, H., Chastel, O., 2014. Demographic responses to mercury exposure in two closely related Antarctic top predators. *Ecology* 95, 1075-1086.
- Tartu, S., Bustamante, P., Goutte, A., Cherel, Y., Weimerskirch, H., Bustnes, J.O., Chastel, O., 2014. Age-Related Mercury Contamination and Relationship with Luteinizing Hormone in a Long-Lived Antarctic Bird. *Plos One* 9, doi: 10.1371/journal.pone.0103642.