

This item is the archived peer-reviewed author-version of:

The specificity of emotional switching in borderline personality disorder in comparison to other clinical groups

Reference:

Houben Marlies, Bohus Martin, Santangelo Philip S., Ebner-Priemer Ulrich, Trull Timothy J., Kuppens Peter.- The specificity of emotional switching in borderline personality disorder in comparison to other clinical groups
Personality disorders - ISSN 1949-2715 - 7:2(2016), p. 198-204
Full text (Publisher's DOI): <https://doi.org/10.1037/PER0000172>
To cite this reference: <https://hdl.handle.net/10067/1381920151162165141>



Published in final edited form as:

Personal Disord. 2016 April ; 7(2): 198–204. doi:10.1037/per0000172.

The Specificity of Emotional Switching in Borderline Personality Disorder in Comparison to Other Clinical Groups

Marlies Houben¹, Martin Bohus², Philip Santangelo³, Ulrich Ebner-Priemer^{2,3}, Timothy Trull⁴, and Peter Kuppens¹

¹KU Leuven

²Central Institute of Mental Health Mannheim, Germany

³Karlsruhe Institute of Technology, Germany

⁴University of Missouri, Columbia, MO, USA

Abstract

In an attempt to better understand the nature of emotion dysregulation in the daily lives of persons with a borderline personality disorder (BPD), Houben, Vansteelandt et al. (2015) recently identified emotional switching, which refers to the tendency to make large changes between positive and negative emotional states over time, as a possible defining characteristic of the emotion dynamics observed in BPD. The goal of this study was to examine the specificity of these previous findings in two samples, by comparing BPD patients ($N = 43$ in sample 1; $N = 81$ in sample 2) to patients with bulimia nervosa ($N = 20$) or posttraumatic stress disorder ($N = 28$) or healthy controls ($N = 28$) in sample 1, and to patients with depressive disorder ($N = 50$) in sample 2, with respect to measures of emotional switching. Analyses of these two experience sampling datasets revealed that contrary to expectations, BPD patients did not differ from the clinical groups regarding their mere tendency to switch between positive and negative emotional states on consecutive moments over time, and regarding the magnitude of such changes between positive and negative emotional states over time. However, all clinical groups did differ from healthy controls regarding all switch measures in dataset 1. These results indicate that emotional switching, similar to other more traditional indicators of overall changes in emotional intensity in daily life, might reflect a feature of emotional responding characterising a range of disorders with mood disturbances.

Emotion dysregulation is generally considered a core problem of borderline personality disorder (BPD; Carpenter, & Trull, 2013; Linehan, 1993). Therefore, over the past few years, studies have increasingly investigated the emotional world of people suffering from BPD in daily life (Santangelo, Bohus, & Ebner-Priemer, 2014), in order to examine the exact nature of emotional dysfunction in BPD in an ecologically valid way. Such daily life studies typically use experience sampling methods (ESM), in which emotions and symptoms are tracked over time in daily life using for example smartphones (Ebner-Priemer, & Trull, 2009), that enable the investigation of patterns of emotional change that are characteristic of

BPD. Understanding the true nature of emotional instability in daily life is crucial to understanding the core of BPD and to inform its treatment. In this paper, we examine whether emotional switching, a specific type of emotional instability that may be particularly typical for BPD, is specific to BPD compared to other clinically relevant comparison groups.

Emotion Dynamics in Borderline Personality Disorder

Experience sampling studies investigating patterns of emotional change in BPD have shown that BPD is mainly characterized by large changes in emotional intensity over time. For example, a recent meta-analysis linking different patterns of emotional change to different indicators of psychopathology including BPD (Houben, Van Den Noortgate, & Kuppens, 2015), concluded that BPD was consistently linked to negative emotions that are more variable over time (meaning they show larger deviations from mean emotional levels over time, measured by the within-person variance of repeated emotions ratings; see also Ebner-Priemer et al., 2015), and positive or negative emotions that show larger changes from one moment to the next (measured using for instance the mean square successive difference or MSSD; see also Ebner-Priemer et al., 2007; Trull et al., 2008). Moreover, findings based on other operationalizations of emotional change (i.e. acute change, Trull et al., 2008; aggregated point-to-point-changes, Ebner-Priemer et al., 2007) also support the idea that BPD patients show larger changes in the intensity of their emotions over time in daily live.

However, recently, a study by Houben, Vansteelandt et al. (2015) approached emotional instability more from a qualitative perspective and showed that what distinguishes BPD patients from healthy controls is not just the occurrence of overall larger changes in emotional intensity, as suggested by previous studies, but specifically larger changes in which a person switches from a positive to a negative emotional state or from a negative to a positive emotional state from one moment to the next, labelled emotional switching. They found that although BPD patients did not differ from healthy controls regarding their overall tendency to switch from a positive to a negative emotional state or vice versa (i.e. switch propensity), the size of such changes between positive and negative emotional states (i.e. switch distances) was significantly larger in BPD patients. In contrast, the magnitude of emotional changes within the negative or positive emotional range separately (e.g. if they change from one unpleasant to another unpleasant state) was similar for both groups. Thus by taking the occurrence of sudden changes in valence of the emotional states into account, they found that not overall changes in intensity, but specifically changes in which one switches between a positive and a negative emotional state are specially larger in BPD when compared to healthy people.

A crucial next step is to establish to what extent emotional switching is specific to BPD, or rather reflects a transdiagnostic feature of several mental disorders. On the one hand, emotional switching may be specific to BPD, reflecting a type of emotional dysfunction that differentiates BPD from other mental disorders and specifically underlies BPD. The notion of emotional switching resonates with the concept of dichotomous or black-white thinking (see Houben, Vansteelandt et al., 2015), which has been linked to BPD and involves evaluations of people, self, and feelings in terms of all-good or all-bad, with the possibility

of abrupt switches between the two extremes (Beck et al, 1990). This suggests that emotional switching might reflect a kind of dysfunction specifically linked to BPD.

On the other hand, it could be that emotional switching reflects a feature of emotional responding characterizing people with a range of mental disorders. This idea is in line with findings from previous studies in which no difference between BPD and other clinical disorders were found for several features of emotional responding in daily life. In a meta-analysis linking BPD to more variable emotions and emotions that showed larger changes from one point to the next (Houben, Van Den Noortgate et al, 2015), results revealed that several indicators of psychological ill-being, such as major depressive disorder or mere neuroticism were linked to the same emotion dynamical patterns. Moreover, a recent study by Santangelo, Reinhard et al. (2014) has shown that in comparison to persons with a diagnosis of bulimia nervosa or posttraumatic stress disorder, no difference was found for BPD patients concerning various indicators of overall changes in emotional intensity, such as the MSSD, probability of acute change, and aggregated point-by-point changes. On the other hand, a study by Trull et al. (2008) found that BPD patients showed higher variability in average negative affect and fear over time, but also larger changes from one point to the next (as measured by MSSD) in hostility, fear, and sadness specifically when compared to patients with a depressive disorder. Moreover, BPD patients were also more likely to make acute changes in hostility in comparison to patients with a depressive disorder. Moreover, when taking comorbidity into account, Scheiderer, Wang, Tomko, Wood, and Trull (2015) indicated that persons with BPD that had a comorbid diagnosis of posttraumatic stress disorder showed larger changes from one point to the next in the intensity of fear and sadness specifically than persons with only a diagnosis of BPD, while for persons with a depressive disorder the reverse pattern was found. In sum, results are mixed regarding the specificity of several indicators of change in the mere intensity of emotions, and it is not clear to what degree measures of switching between positive and negative emotional states do or do not reflect a feature that is specific to BPD patients.

The Current Study

The current study builds on the previous study by Houben, Vansteelandt et al. (2015) in which the concept of emotional switching in BPD was introduced and presents a reanalysis of the data reported in Santangelo, Reinhard et al. (2014) and in Scheiderer et al. (2015), in which so far only measures of changes in overall emotional intensity or negative emotional intensity specifically (i.e. quantitative changes) were examined.

The goal of this study is to investigate whether emotional switching between positive and negative emotions states, a measure that captures more qualitative emotional changes, can differentiate between persons with BPD and with other psychiatric diagnoses in two datasets. Using data from persons with BPD, bulimia nervosa or posttraumatic stress disorder and healthy controls as reported in Santangelo, Reinhard et al. (2014) and data of persons with BPD and with a depressive disorder (major depressive disorder or dysthymic disorder) as reported in Scheiderer et al. (2015) and Trull et al. (2008), we compared the different groups in terms of their propensity to switch (i.e. the probability that someone switches between positive and negative emotions states, irrespective of the size of such

changes), and their average switch distance from a positive to a negative emotional state or vice versa (i.e. reflecting how large/extreme such changes are) from one assessment occasion to the next. As the idea of emotional switching is based on clinical observations of BPD and theoretical concepts linked to BPD, we hypothesized that the occurrence of larger switch distances, as found in a previous study (Houben, Vansteelandt et al, 2015), is a distinctive feature of BPD, when compared to other clinical groups.

Method

Participants

Sample one—The first sample consisted of 43 persons with BPD, 28 persons with a posttraumatic stress disorder, 20 persons with bulimia nervosa, and 28 healthy controls. Participants were all female with an age between 18 and 48 ($M = 26.72$, $SD = 7.07$ for BPD subsample; $M = 35.25$, $SD = 7.53$ for posttraumatic stress subsample; $M = 23.70$, $SD = 5.97$ for bulimia nervosa subsample; $M = 28.82$, $SD = 7.47$ for healthy controls). The clinical groups consisted of inpatients and outpatients recruited at German mental health institutions. Healthy controls had no current or past Axis I or Axis II disorder diagnoses, and were recruited using advertisements in local new papers or were randomly selected from the national resident register of the City of Mannheim. For detailed diagnostic information, and for information regarding comorbidity, exclusion criteria and diagnostic procedure, see Santangelo, Reinhard et al. (2014).

Sample two—Our second sample consisted of 81 persons with BPD and 50 persons with a depressive disorder (major depressive disorder or dysthymic disorder). Most participants were female (93% of the BPD subsample; 78% of the depressive disorder subsample). Mean age was 31.94 ($SD = 11.68$) for the BPD group and 34.76 ($SD = 11.95$) for the depressive disorder group. Recruitment of all patients took place in four local outpatients psychiatric clinics. For more information about (comorbid) diagnoses, diagnostic procedure and exclusion criteria, see Scheiderer et al. (2015) and Trull et al. (2008).

Experience Sampling Procedure

Sample one—After being carefully trained, participants carried a palmtop (Tungsten E) with them that generated a questionnaire every 15 minutes (± 1 minute) during 24 hours, however excluding sleeping hours during which the palmtop was shut down. Each questionnaire was announced with a beeping signal. At each assessment moment, participants were asked to identify the emotion they were experiencing from a list of possible discrete emotional states, including happy, anxious, angry, shame, disgust, sad, guilt, interest, envy/jealousy, emotion but cannot name it, and no emotion. Next they were asked to rate the intensity of that emotion using a scale ranging from 1 to 11. These intensity scores were converted into a bipolar valence score for each moment by only multiplying the intensity of a current negative emotion by -1 , yielding a negative valence score if they reported a negative emotion, and retaining a positive valence score if participants reported a positive emotion, ranging from -11 to 11 . Average compliance rates to the ESM protocol was very good (BPD patients completed 94% of ESM questionnaires, patients with posttraumatic stress disorder 93%, and persons with bulimia nervosa and healthy controls

94%). For more detailed information about the data collection procedure, see Santangelo, Reinhard et al. (2014).

Sample two—Participants received training and next carried a palmtop (Palm Zire 31) with them for approximately 28 days. During these days, the palmtop generated a questionnaire 6 times a day, equally spread over the day using stratified interval schemes. Each questionnaire was signaled by a beeping sound. Next, among other questions, participants were asked to indicate to what degree they had experienced certain emotions since the last beep on a scale of 1 (not at all/very slightly) to 5 (extremely) using the emotion items from the Positive and Negative Affect Schedule (PANAS; Watson & Clark, 1999). The 10 positive items and the 10 negative items were averaged to form a composite positive affect scale (PA) and negative affect scale (NA), respectively. Reliability estimates were obtained following suggestions by Nezlek (2012), using three-level models with items at Level 1, nested within beeps at Level 2, nested within persons at Level 3 and showed excellent reliability for PA (estimate = .83 at the level of the measurement occasion; estimate = .99 at the person level) and for NA (estimate = .81 at the level of the measurement occasion; estimate = .99 at the person level). Next, a bipolar valence variable was created by subtracting NA from PA, yielding positive valence scores if PA was higher than NA, and negative valence scores if NA was higher than PA, ranging from -4 to 4. BPD patients and patients with a depressive disorder showed good compliance to the ESM protocol and completed an average of 85% and 87% of ESM questionnaires respectively. For more detailed information about the data collection procedure, see Scheiderer et al. (2015) and Trull et al. (2008).

In both datasets, a switch was defined as a change from an emotional state with positive valence to an emotional state at the next time point within the same day that was characterized by negative valence, or the other way around. Changes within the positive emotional range or within the negative emotional range within the same day were defined as non-switches.

Statistical analysis

In both datasets, repeated measurements are nested within persons. To deal with this nested structure, we applied two-level analyses to analyze the data, using HLM 7 (Raudenbush, Bryk, Cheong, Congdon, & du Toit, 2011). For detailed information about the modeling approach, see Houben, Vansteelandt et al. (2015).

Differences in switch propensity (i.e. the probability for a switch to occur) were modeled using two-level logistic regression models, using a random intercept at the moment-to-moment level (i.e. level 1) of the model, and using diagnostic group dummies (leaving out the intercept) at the person-level (i.e. level 2) of the model. Differences between the slopes of the different dummy variables were statistically tested using General Linear hypothesis testing. Unit-specific models were used, although population-average models led to similar conclusions.

Differences in switch (and non-switch) distances were modeled using two-level linear regression models in which absolute changes in valence between two consecutive time points were modeled as a function of a switch and a non-switch dummy at the moment-to-moment level of the model, leaving out the intercept, and different diagnostic group dummies at the person-level of the model, again leaving out the intercepts. Differences in slopes were tested using General Linear hypothesis testing.

Results

Switch Propensity

Table 1 shows the estimated switch propensity for each group and the difference between the groups in each dataset. For dataset 1, results showed that all clinical groups (including persons with BPD, bulimia nervosa and posttraumatic stress) had a higher propensity to switch, in comparison to healthy controls. However, no differences were found among the clinical groups. In dataset 2, similarly, no difference was found between patients with BPD and with a depressive disorder for the propensity to switch.

Switch Distance

In Table 2, the estimated switch and non-switch distances for all groups and the estimated differences between groups are shown for both samples. In Table 3, similar results are shown, however distinguishing between switching from positive to negative and from negative to positive emotional states.

In dataset 1, BPD patients, but also persons with bulimia nervosa and with posttraumatic stress disorder showed significantly larger switch distances and larger non-switch distances in comparison to healthy controls. These results hold for switching in both directions. However, no differences were found between BPD patients and any of the clinical control groups (i.e. patients with bulimia nervosa or posttraumatic stress disorder), for both switch and non-switch distances. To check the robustness of findings, we repeated the analysis estimating switch and non-switch distances, however correcting for individual differences in the extremity of the starting points of calculated distances at level 1 of the model (i.e. the absolute value of the valence rating at time $t-1$). This was done because for more extreme valence ratings that deviate more from the neutral valence point, a larger change has to occur in order to result in a switch. For switch distance, similar results were obtained as described above. However for non-switch distances, differences between all clinical groups and healthy controls were no longer significant after correction for starting points ($ps > .50$).

For dataset 2, we found that BPD patients showed switch distances that were significantly larger than those of persons with a depressive disorder. No difference was found for non-switch distances. Distinguishing between the two directions of switching, results indicated that BPD individuals displayed significantly larger switch distances from positive to negative emotions states specifically. However, if we modeled switch and non-switch distances, additionally corrected for the starting point of the calculated distances, the difference between BPD patients and patients with a depressive disorder was non-significant ($p = .11$).

Discussion

The goal of this study was to examine the specificity of emotional switching between positive and negative emotional states for BPD in comparison to other clinical disorders. We used two datasets in which previously it was shown that BPD patients tend to show large changes in overall emotional intensity over time in daily life (Santangelo, Reinhard et al., 2014 and Scheiderer et al., 2015). However, these emotional intensity results were not always specific to BPD after comparison with other clinical groups. The current study extended on these studies, by investigating the specificity of not just changes in intensity of emotions in general but specifically of changes between positive and negative emotional changes (i.e. emotional switching) over time.

Our findings indicated that in comparison to healthy controls, all clinical groups (including BPD, bulimia nervosa or posttraumatic stress disorder) were more likely to make switches between positive and negative emotions states over time. Moreover, if they switched, the size of the emotional changes from positive to negative, or from negative to positive emotional states were also larger. However, regarding the specificity of these findings, in the first dataset, no differences were observed between persons with BPD and the two clinical control groups, or between the two different clinical control groups. In dataset 2, some indications were found that patients with BPD show larger switch distances than patients with a depressive disorder, however results were not robust when taking into account individual differences in the extremity of start points of calculated distances.

Overall, these results indicate that switching might not be a phenomenon specific to BPD, but most likely reflects a feature of emotional responding common among several disorders (at least, those studied here). These findings are in line with the conclusion formulated by Houben, Van Den Noortgate et al. (2015), in which was shown that several indicators of emotional change were not specifically linked to BPD, as other clinical groups or other indicators of psychological ill-being were characterized by similar patterns. Moreover, the results show that next to limited specificity for indicators of changes in the overall intensity of emotions as concluded in Santangelo, Reinhard et al. (2014), also indicators of emotional switching between positive and negative emotions are not very specific to BPD after comparison with other clinical groups.

From one perspective, the lack of specificity of emotional switching for BPD is a rather surprising finding. Although emotional instability is an important and characterizing diagnostic criterion of BPD, this is not explicitly the case for major depressive disorder or dysthymia, posttraumatic stress disorder, or bulimia nervosa. Moreover, emotion lability is even assumed to play an etiological role for BPD (Linehan, 1993). Therefore, the finding that other clinical groups showed similar levels of emotional instability in daily life (as indicated by emotional switching) compared to BPD raises questions about the completeness of the current diagnostic criteria. However, differences between clinical groups may be found in different aspects of emotional responding. For example, although groups might not differ in the resulting patterns of emotional change, the nature of triggers of such changes could distinguish between the groups.

From a different perspective, however, emotional instability, as indicated by larger emotional switching between positive and negative emotional states might reflect a feature of emotional responding that underlies groups of mental disorders. This finding would fit with current calls for a transdiagnostic approach to the investigation and classification of psychopathology, with a focus on mechanisms or core disturbances (Insel et al., 2010). For example, in the research domain criteria (RDoC), developed by the National Institute of Mental Health a call for a focus on the investigation of underlying mechanisms and measures that cut across current clinical groups was formulated. Related to these propositions, emotional instability could reflect a transdiagnostic dimension underlying psychopathology. Future research should provide more insight into the mechanisms underlying emotional instability. Speculatively, emotional instability could signify a lack of emotional control, which could act as a risk factor for the development of psychopathological conditions in general. On the other hand, emotional instability could also result from negative life events that are often experienced by people with psychopathology in general and that tend to underlie mental disorders, such as traumatic life events. Alternatively, emotional instability could also reflect a possible consequence of dysfunctional behavior exhibited by people with psychopathological problems.

Several limitations should be mentioned. First, our measures of emotional switching are based on self-reported emotions, which has its shortcomings (e.g. possible influence of response biases, it is based on introspective abilities etc.). However, subjective self-report is most commonly used to assess the experiential aspect of affective states, and using ESM limits the influence of memory bias and single assessments as much as possible. Second, the two datasets that were reported differ in two important ways: how valence of the emotion was determined, and the sampling rates used in the studies. In both samples, valence was operationalized in different ways, and therefore may not be directly comparable. Concerning sample rates, in dataset 1 very short time intervals were used that capture consecutive changes in emotions on a small time scale, which is claimed to be most appropriate for the investigation of emotional instability in BPD (Ebner-Priemer, & Sawitzki, 2007). However, assessments only stretch over a time period of 24 hours, in which only a limited amount of variation in events and experiences can be captured. In contrast, in dataset 2 only six assessments per day were recorded, resulting in larger time intervals between consecutive measurements. However, patients were sampled over a period of around 28 days which more likely captured a diverse set of events during the study. Due to these differences it is not clear whether we are measuring the same process across different time scales. Nonetheless, the different studies led to similar conclusions about the specificity of emotional switching. Third, although sample sizes were not very small, we cannot fully exclude the possibility that we lacked sufficient statistical power to detect changes between clinical groups, likely more of a concern in dataset 1. However, subsamples in dataset 2 were larger and resulted in similar conclusions.

Conclusions

Based on two independent samples, we found that several clinical groups, including BPD, bulimia nervosa, and posttraumatic stress disorder all displayed a higher tendency to switch between positive and negative emotional states on consecutive moments over time.

Moreover, if they switched, the size of the emotional changes from positive to negative, or from negative to positive emotional states were also larger in comparison to healthy controls. However, limited specificity of emotional switching was found for BPD, in comparison to patients with bulimia nervosa, posttraumatic stress disorder or a depressive disorder. These findings imply that emotional switching might reflect a feature of emotional responding that characterizes several psychopathological conditions.

References

- Beck, AT.; Freeman, A., et al. Cognitive therapy of personality disorders. New York: Guilford Press; 1990.
- Carpenter RW, Trull TJ. Components of emotion dysregulation in borderline personality disorder: a review. *Current psychiatry reports*. 2013; 15:1–8.
- Ebner-Priemer UW, Houben M, Santangelo P, Kleindienst N, Tuerlinckx F, Oravecz Z, Kuppens P. Unraveling affective dysregulation in borderline personality disorder: A theoretical model and empirical evidence. *Journal of Abnormal Psychology*. 2015; 124:186–198. [PubMed: 25603359]
- Ebner-Priemer UW, Kuo J, Kleindienst N, Welch SS, Reisch T, Reinhard I, Bohus M. State affective instability in borderline personality disorder assessed by ambulatory monitoring. *Psychological Medicine*. 2007; 37:961–970. [PubMed: 17202005]
- Ebner-Priemer UW, Sawitzki G. Ambulatory assessment of affective instability in borderline personality disorder: The effect of the sampling frequency. *European Journal of Psychological Assessment*. 2007; 23:238–247.
- Ebner-Priemer UW, Trull TJ. Ambulatory assessment: An innovative and promising approach for clinical psychology. *European Psychologist*. 2009; 14:109.
- Houben M, Van Den Noortgate W, Kuppens P. The relation between short-term emotion dynamics and psychological well-being: A meta-analysis. *Psychological Bulletin*. 2015; 141:901–930. [PubMed: 25822133]
- Houben M, Vansteelandt K, Claes L, Sienaert P, Berens A, Sleuwaegen E, Kuppens P. Emotional switching in borderline personality disorder: A daily life study. *Personality Disorders: Theory, Research, and Treatment*. 2015 Advance online publication.
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, Wang P. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *The American journal of psychiatry*. 2010; 167:748–751. [PubMed: 20595427]
- Linehan, M. Cognitive behavioral treatment of borderline personality disorder. Guilford Press; 1993.
- Raudenbush, SW.; Bryk, AS.; Cheong, YF.; Congdon, RT.; du Toit, M. HLM 7: Hierarchical linear and nonlinear modeling. Chicago, IL: Scientific Software International; 2011.
- Santangelo P, Bohus M, Ebner-Priemer UW. Ecological momentary assessment in borderline personality disorder: A review on recent findings and methodological challenges. *Journal of Personality Disorders*. 2014; 28:555–576. [PubMed: 22984853]
- Santangelo P, Reinhard I, Mussgay L, Steil R, Sawitzki G, Klein C, Ebner-Priemer UW. Specificity of affective instability in patients with borderline personality disorder compared to posttraumatic stress disorder, bulimia nervosa, and healthy controls. *Journal of abnormal psychology*. 2014; 123:258–272. [PubMed: 24661176]
- Scheiderer EM, Wang T, Tomko RL, Wood PK, Trull TJ. Negative affect instability among individuals with comorbid borderline personality disorder and posttraumatic stress disorder. *Clinical Psychological Science*. 2015 Advance online publication.
- Nezlek, JB. Multilevel modeling analyses of diary-style data. In: Mehl, MR.; Conner, TS., editors. *Handbook of research methods for studying daily life*. New York, NY: Guilford Press; 2012. p. 357–383.
- Trull TJ, Solhan MB, Tragesser SL, Jahng S, Wood PK, Piasecki TM, Watson D. Affective instability: Measuring a core feature of borderline personality disorder with ecological momentary assessment. *Journal of Abnormal Psychology*. 2008; 117:647–661. [PubMed: 18729616]

Table 1
Results from Two-level Logistic Regression Models Predicting Switch Propensity by Diagnostic Status

SAMPLE 1		Test of difference with BPD						
Predicted event	Level 1	Level 2	Estimate	SE	t (df)	p value	χ^2 (df)	p-value
Switch	Intercept, β_{0j}	BPD dummy, γ_{01}	-1.87 _a	0.14	-13.37 (115)	< .01		
		PTSD dummy, γ_{02}	-2.17 _a	0.20	-10.74 (115)	< .01	1.44 (1)	.23
		BN dummy, γ_{03}	-1.76 _a	0.22	-7.92 (115)	< .01	0.19 (1)	> .50
		HC dummy, γ_{04}	-3.01 _b	0.22	-13.44 (115)	< .01	18.59 (1)	< .01
SAMPLE 2								
Switch	Intercept, β_{0j}	BPD dummy, γ_{01}	-2.03 _a	0.10	-20.26 (129)	< .01		
		DEP dummy, γ_{02}	-1.95 _a	0.14	-13.60 (129)	< .01	0.21 (1)	> .50

Note. BPD = borderline personality disorder; PTSD = posttraumatic stress disorder; BN = bulimia nervosa; HC = healthy controls; DEP = depressive disorder. For each sample, coefficients at level 2 having a common subscript are not statistically different at $\alpha = .05$.

Table 2

Results from Two-level Linear Regression Models estimating Switch Distances and Non-Switch Distances as a Function of Diagnostic Status

SAMPLE 1		Test of difference with BPD				
Level 1	Level 2	Estimate	SE	t (df)	p value	p-value
Switches slope, β_{ij}	BPD dummy, γ_{11}	2.50 _a	0.04	59.67 (115)	< .01	
	PTSD dummy, γ_{12}	2.54 _a	0.06	42.67 (115)	< .01	0.32 (1)
	BN dummy, γ_{13}	2.50 _a	0.06	41.18 (115)	< .01	.00 (1)
	HC dummy, γ_{14}	2.11 _b	0.07	29.59 (115)	< .01	21.33 (1)
Non-switches slope, β_{2j}	BPD dummy, γ_{21}	0.96 _a	0.03	33.84 (155)	< .01	
	PTSD dummy, γ_{22}	0.94 _a	0.05	19.09 (115)	< .01	.09 (1)
	BN dummy, γ_{23}	0.92 _a	0.05	18.77 (155)	< .01	0.43 (1)
	HC dummy, γ_{24}	0.74 _b	0.06	12.60 (115)	< .01	11.23 (1)
SAMPLE 2						
Switches slope, β_{ij}	BPD dummy, γ_{11}	0.81 _a	0.03	26.40 (129)	< .01	
	DEP dummy, γ_{12}	0.72 _b	0.03	21.98 (129)	< .01	3.99 (1)
Non-switches slope, β_{2j}	BPD dummy, γ_{21}	0.44 _a	0.02	29.27 (129)	< .01	
	DEP dummy, γ_{22}	0.43 _a	0.02	25.84 (129)	< .01	0.57 (1)

Note. BPD = borderline personality disorder; PTSD = posttraumatic stress disorder; BN = bulimia nervosa; HC = healthy controls; DEP = depressive disorder. For each slope at level 1 separately, corresponding coefficients at level 2 having a common subscript are not statistically different at $\alpha = .05$.

¹Note that the differences between clinical groups and HC become insignificant after correction for starting points (ps > .50)

²Note that this difference between BPD and DEP becomes insignificant after correction for starting points (p = .11).

Table 3

Results from Two-level Linear Regression Models estimating Switch Distances To Different Directions and Non-Switch Distances as a Function of Diagnostic Status

SAMPLE 1		Test of difference with BPD				
Level 1	Level 2	Estimate	SE	t (df)	p value	χ^2 (df)
Switches to positive, β_{1j}	BPD dummy, γ_{11}	2.50 _a	0.04	58.50 (115)	<.01	
	PTSD dummy, γ_{12}	2.54 _a	0.06	41.51 (115)	<.01	0.32 (1)
	BN dummy, γ_{13}	2.51 _a	0.06	39.29 (115)	<.01	0.03 (1)
	HC dummy, γ_{14}	2.11 _b	0.07	29.49 (115)	<.01	21.43 (1)
Switches to negative, β_{2j}	BPD dummy, γ_{21}	2.50 _a	0.04	57.57 (115)	<.01	
	PTSD dummy, γ_{22}	2.54 _a	0.06	42.12 (115)	<.01	0.30 (1)
	BN dummy, γ_{23}	2.49 _a	0.06	39.76 (115)	<.01	0.01 (1)
	HC dummy, γ_{24}	2.12 _b	0.09	24.08 (115)	<.01	14.95 (1)
Non-switches, β_{3j}	BPD dummy, γ_{31}	0.96 _a	0.03	33.84 (115)	<.01	
	PTSD dummy, γ_{32}	0.94 _a	0.05	19.09 (115)	<.01	0.09 (1)
	BN dummy, γ_{33}	0.92 _a	0.05	18.76 (115)	<.01	0.43 (1)
	HC dummy, γ_{34}	0.74 _b	0.06	12.60 (115)	<.01	11.23 (1)
SAMPLE 2						
Switches to positive, β_{1j}	BPD dummy, γ_{11}	0.78 _a	0.03	24.47 (129)	<.01	
	DEP dummy, γ_{12}	0.72 _a	0.04	19.83 (129)	<.01	1.71 (1)
Switches to negative, β_{2j}	BPD dummy, γ_{21}	0.84 _a	0.03	26.03 (129)	<.01	
	DEP dummy, γ_{22}	0.73 _b	0.03	20.95 (129)	<.01	6.05 (1)
Non-switches, β_{3j}	BPD dummy, γ_{31}	0.44 _a	0.02	29.27 (129)	<.01	
	DEP dummy, γ_{32}	0.43 _a	0.02	25.84 (129)	<.01	0.57 (1)

Note. BPD = borderline personality disorder; PTSD = posttraumatic stress disorder; BN = bulimia nervosa; HC = healthy controls; DEP = depressive disorder. For each slope at level 1 separately, corresponding coefficients at level 2 sharing a common subscript are not statistically different at $\alpha = .05$.