

This item is the archived peer-reviewed author-version of:

Cannabis sativa allergy : looking through the fog

Reference:

Decuyper Ine, Van Gasse A.L., Cop Nathalie, Sabato Vito, Faber Margaretha, Mertens Christel, Bridts Christiaan, Hagendorens Margo, De Clerck Luc S., Rihs H.P.,- Cannabis sativa allergy : looking through the fog
Allergy: European journal of allergy and clinical immunology - ISSN 0105-4538 - 72:2(2017), p. 201-206
Full text (Publisher's DOI): <http://dx.doi.org/doi:10.1111/ALL.13043>
To cite this reference: <http://hdl.handle.net/10067/1450870151162165141>

Cannabis sativa allergy: looking through the fog

Ine I Decuyper, MD^{1,2}, Athina L Van Gasse, MD^{1,2}, Nathalie Cop MLT¹, Vito Sabato, MD, PhD¹, Margaretha Faber MD, PhD¹, Christel Mertens MLT¹, Chris H Bridts MLT¹, Margo M Hagendorens, MD, PhD^{1,2}, Luc De Clerck, MD, PhD¹, H.P. Rihs, PhD³, Didier G Ebo, MD, PhD^{1, *}

¹ Faculty of Medicine and Health Sciences, Department of Immunology – Allergology – Rheumatology, University of Antwerp, Antwerp University Hospital, Belgium

² Faculty of Medicine and Health Sciences, Department of Pediatrics, University of Antwerp, Antwerp University Hospital, Belgium

³ Ruhr-University Bochum, IPA - Institute for Prevention and Occupational Medicine, German Social Accident Insurance, Bochum, Germany

Word count abstract: 194

Word count manuscript: 2375

***Correspondence to:**

Didier Ebo

University of Antwerp

Faculty of Medicine and Health Sciences

Department of Immunology, Allergology, Rheumatology

Campus Drie Eiken T5.95

Universiteitsplein 1

2610 Antwerpen, Belgium.

Email: didier.ebo@uantwerpen.be

Institutional email: immuno@uantwerpen.be

Abstract

IgE-mediated *Cannabis* (*C. sativa*, marijuana) allergy seems to be on the rise. Both active and passive exposure to cannabis allergens may trigger a *C. sativa* sensitization and/or allergy. The clinical presentation of a *C. sativa* allergy varies from mild to life-threatening reactions and often seems to depend on the route of exposure. In addition, sensitization to cannabis allergens can result in various cross-allergies, mostly for plant foods. This clinical entity, designated as the “cannabis-fruit/vegetable syndrome” might also imply cross-reactivity with tobacco, natural latex and plant-food derived alcoholic beverages. Hitherto, these cross-allergies are predominantly reported in Europe and appear mainly to rely upon cross-reactivity between non-specific lipid transfer proteins (ns-LTPs) or thaumatin-like proteins (TLPs) present in *C. sativa* and their homologues, ubiquitously distributed throughout plant kingdom. At present, diagnosis of cannabis-related allergies predominantly rests upon a thorough history completed with skin testing using native extracts from crushed buds and leaves. However, quantification of specific IgE (sIgE) antibodies and basophil activation tests (BAT) can also be helpful to establish correct diagnosis. In the absence of a cure, treatment comprises absolute avoidance measures. Whether avoidance of further use will halt the extension of related cross-allergies remains uncertain.

Key words: basophil activation, cannabis sativa, cross-reactivity, diagnosis, IgE.

List of abbreviations

Act d	<i>Actinidia deliciosa</i> (kiwi fruit)
Ara h	<i>Arachis hypogaea</i> (peanut)
Art v 3	<i>Artemisia vulgaris</i> (Mugwort)
ATP	adenosine triphosphate
BAT	basophil activation test
Bet v	<i>Betula verrucosa</i> (birch)
Can s	<i>Cannabis sativa</i>
CCD	cross-reactive carbohydrate determinants
Cit s	<i>Citrus sinensis</i> (tangerine)
Cor a	<i>Corylus avellana</i> (hazelnut)
CRD	component resolved diagnosis
Cup a	<i>Cupressus arizonica</i> (cypress native to the southwest of north America)
Hev b	<i>Hevea brasiliensis</i> (natural rubber tree)
Jug r 3	<i>Juglans regia</i> (walnut)
Lyc e	<i>Lycopersicon esculentum</i> (tomato)
Mal d	<i>Malus domestica</i> (apple)
Mus a	<i>Musa acuminata</i> (banana)
Ns-LTP	non-specific lipid transfer protein
NSAIDs	nonsteroidal anti-inflammatory drugs
OAS	oral allergy syndrome
Phl p	<i>Phleum pratense</i> (Timothy grass)
PRP	pathogenesis-related protein
Pru av	<i>Prunus avium</i> (cherry)

Pru p	<i>Prunus persica</i> (peach)
RuBisCo	ribulose-1,5-biphosphate carboxylase/oxygenase
slgE	specific immunoglobulin E
THC	tetrahydrocannabinol
TLP	thaumatin-like protein
Tri a	<i>Triticum aestivum</i> (wheat)
Vit v	<i>Vitis vinifera</i> (grape)

Introduction

Cannabis sativa (*C. sativa*) is an annual, dioecious and anemophilous flowering plant (order: Rosales, family *Cannabaceae*) that is native to central and southern Asia and the Caucasian region. Different preparations [dried flowering tops, hashish, hashish oil] are obtained from cannabis varieties containing elevated levels of cannabinoids, especially delta 9-tetrahydrocannabinol (THC); several of them being more or less potent psychoactive substances. Although, today, cannabis use is still illegal in most countries, it is widespread for its relaxing and euphoric effects. Furthermore, Worldwide the illegal status of the drug has gained more and more resistance recently, resulting in legalisation of both sale and possession of marijuana in Colorado, Alaska, Oregon and Washington for example, both for medicinal and recreational use (1). Different European countries are debating the legalisation of cannabis as well. While consumption of cannabis has been legal under certain conditions for a while in the Netherlands, both Germany's and Belgium's minister of Health plan to legalise certain medicinal cannabis preparations in the near future (2, 3)_When used, it can be consumed by smoking, vaporizing or gastro-intestinal ingestion. In addition to hempseed and hempseed oil, derivatives of dried flowers or resinous extract can be incorporated in food and ingested (4-6). *C. sativa* in the form of industrial hemp containing lower amounts of THC is commercially used for fibre, cosmetics, and clothing. Apart from these various (ab)uses unfortunately, cannabis may elicit some undesirable effects amongst which allergy and the consequential allergic symptoms seem to be an increasing problem. This review aims to summarize the current knowledge on cannabis-related allergy, with a particular focus on clinical manifestations, allergenic components and management of the patients.

Prevalence

Few true IgE-dependent allergic reactions in illicit abusers have been published (7-10) making it hard to determine the true prevalence of this allergy. Nevertheless, multiple reports on cutaneous and respiratory allergies to different members of the *Cannabaceae* family like industrial hemp and hop (*Humulus lupulus*) have been published (11-15). This under-reporting probably results from the illegal status of cannabis use, which makes the patients reluctant to admit their abuse. Liskow *et al.*, in 1971, described a 29-year-old house wife who suffered an allergic reaction upon smoking a marijuana cigarette. Diagnosis of cannabis allergy was documented by a positive scratch test and passive transfer studies (16). Today, no information is available about the prevalence of IgE-mediated *C. sativa* allergy but it is likely that cannabis allergy will be an increasing problem in the future, partly in the settings of its evolving legal status (6).

Routes of exposure and sensitization

When cannabis is used for its psychoactive effects, drug(ab)users may become sensitized by inhalation of cannabis allergens through active smoking and/or vaporizing the drug. Cutaneous contact through handling of the drug is another possible route of sensitization. This last route of exposure is not only relevant in drug (ab) users but could be important in cannabis growers and police men seizing illegal cannabis (plants). Herzinger *et al.* (15) also reported cannabis allergy in laboratory workers handling cannabis for professional purposes. Cannabis derivatives may be used in food preparations which can cause allergic sensitization by chewing or ingestion. Intravenous use can also cause both sensitization and elicitation of allergic symptoms. An alternative route of sensitization could be exposure by proxy when allergens become airborne or are transferred via indirect cutaneous contact. Such a second-hand exposure might be particularly relevant in children or bystanders who may become sensitized to cannabis allergens through smoking by relatives or friends (8).

In contrast to a birch pollen sensitization that is known to cause a pollen–food syndrome with secondary food allergies, cannabis seems not always the primary allergic source and allergic symptoms on exposure to cannabis might also originate from cross-reactivity with allergenic compounds such as non-specific lipid transfer proteins (ns-LTPs) or thaumatin-like proteins (TLPs) present in other plants from closely or more distantly related origin (10). Finally, *C. sativa* produces wind-borne pollen easily transported over long distances (17-22). For example, in Nebraska, where industrial *C. sativa* is cultivated, *C. sativa* pollen account for 36% of the total pollen count during mid- to late-August (18). Similar observations were made in Italy (19) and Spain (21). Multiple reports were published in which allergic symptoms are presumed to result from cannabis pollen sensitization (23-25). Interestingly, the link with ns-LTPs from plant-foods has only been described in relation to cannabis sensitization in drug (ab)users no studies have been published relating ns-LTPs or TLPs to cannabis pollen. As only female (non-pollinating) plants are cultivated for illicit use, it is less likely for abusers of cannabis, who grow their own plants, to become sensitized to marijuana through pollen exposure. Furthermore, a correlation between ns-LTP sensitization more specifically Can s 3 and cannabis pollen has not been described up till now.

Allergenic components

At present, the allergenic composition of *C. sativa* remains largely unknown. Larramendi *et al.* (10) described 6 different bands with a molecular weight varying between 10 and 60-kDa in a *C. sativa* leaf extract that were recognized by the individual patients' sera. The 10-kDa IgE binding band was already described in other reports (10, 26, 27) and corresponds to Can s 3, the non-specific lipid protein (ns-LTP) of *C. sativa* (10) that belongs to the pathogenesis-related proteins (PR)-14 group (28). Sensitization to the ns-LTP Can s 3 was observed in 124 out of 130 patients (95.3%) with a primary cannabis allergy in a Spanish study (29). In our own case-control

study sensitization to ns-LTP was demonstrable in 10 out of 12 (83%) patients with cannabis-related food allergies (8). In comparison, in our regions sIgE reactivity towards ns-LTP in patients with pollen and/or plant-derived food allergies is observed in about 25% of the cases (30). Anyhow, it appears that next to peach (26) and mugwort (31) *C. Sativa* might be another source that dominates sensitization to ns-LTP and triggers ns-LTP-related allergies that extend far beyond vegetables and fruits. The 38-kDa band corresponds with a TLP, which belongs to the PR-5 family (10). Although in the study of Larramendi *et al.* (10) no homology was found between the 14-kDa band and any known allergen, it was speculated that this band could be a profilin (32).

In a study of Nayak *et al.* (33), multiple IgE binding proteins were observed. The 23-kDa band was identified as 'oxygen-evolving enhancer protein 2', an enzyme involved in the photosynthesis. The 50-kDa band corresponds with the heavy chain subunit of ribulose-1,5-biphosphate carboxylase/oxygenase (RuBisCo). This is a highly abundant protein in nature that catalyses a reaction that is rate-limiting for photosynthesis. Other putative allergens identified by Nayak *et al.* (33) are glyceraldehyde-3-phosphate and adenosine triphosphate (ATP) synthase. Finally, the authors observed that ubiquitously distributed cross-reactive carbohydrate determinants (CCDs) might also be the cause of some IgE reactivity. Unlike the European studies, in this American/Canadian proteomics study no IgE binding sequences of the pan-allergen ns-LTP were observed, even though IgE reactivity at approximately 10-kDa was observed in two patients. Moreover, in contrast to the European series, most of the Canadian patients apparently did not suffer from a cannabis-related cross-reactivity syndrome as is described below. Whether this indicates cannabis allergy to display geographically different sensitization profiles with distinct clinical phenotypes remains elusive. The most relevant putative cannabis allergens are displayed in table 1.

Table 1. (Putative) <i>cannabis sativa</i> allergens				
Molecular weight (kDa)	Allergen	Function	Homologues (not exhaustive)	References
9	Can s 3	Ns-LTP (PR-14)	Pru p 3, Mal d 3, Cor a 8, Hev b 12, Ara h 9, Tri a 14, Jug r 3, Art v 3	(8, 10, 26, 29, 31, 32, 34-36)
14	Profilin (?)	Cytoskeleton	Bet v 2, Phl p 12	(32)
23	Oxygen-evolving enhancer protein	Photosynthesis		(33)
38	TLP (thaumatin-like)	PR-5	Act d 2, Mal d 2, Mus a 4, Pru av 2, Cup a 3	(10)
50	Can s RuBisCo	Photosynthesis		(33)
Table updated from Decuyper <i>et al.</i> (37).				
Ns-LTP: non-specific lipid protein, PR: pathogenesis-related.				
Except the study published by Larramendi <i>et al.</i> (10) allergens described in the table have been suggested without deep investigation and warrant further studies.				


Clinical manifestations

The clinical presentation of an IgE-mediated cannabis allergy can vary considerably from mild to life-threatening reactions and seems to relate to the route of exposure. First, respiratory

reactions like rhinitis, conjunctivitis, asthma and palpebral angioedema have been described. These reactions predominantly occur when cannabis is consumed by smoking or vaporizing (for review: (6, 37)) but can also arise from passive second-hand exposure to cannabis smoke, indirect cutaneous transmission (8, 26) or inhalation of *C. sativa* pollen (6, 17-22).

Direct handling of *C. sativa* plants may lead to contact urticaria (13, 35, 38) and contact dermatitis (13). Finally, anaphylaxis can result from ingestion of hempseed (5), drinking marijuana tea (39) and smoking (40).

Patients with IgE-mediated cannabis allergy can display distinct sensitization profiles such as sensitization to the ns-LTP of *C. sativa*, *i.e.* Can s 3. Non-specific lipid transfer proteins are pan-allergens ubiquitously present throughout the plant kingdom including fruits and vegetables (41). Consequently, sensitization to Can s 3 might be an explanation for the high variety of secondary plant-derived food allergies seen in European patients with a cannabis allergy. This, sometimes extensive, cross-reactivity between cannabis and plant-derived food has been described in multiple studies (6, 8, 37, 40) and was recently designated as the “cannabis-fruit/vegetable syndrome” (8, 42) The food most commonly implicated in this allergy syndrome are peach, apple, nuts, tomato and occasionally citrus fruit as orange and grapefruit. Figure 1, adapted from (37) gives a non-exhaustive overview of this “cannabis-fruit/vegetable syndrome”.



Legend to Figure 1. Non-specific lipid transfer proteins (ns-LTPs) are ubiquitously present in the plant kingdom. Consequently, sensitization to Can s 3, the ns-LTP from *Cannabis sativa*, could lead to a broad variety of cross-reactions. Cross-reactive substances displayed in the figure: cherry (*Prunus avium*), tangerine (*Citrus reticulata*), orange (*Citrus sinensis*), peach (*Prunus persica*), apple (*Malus domestica*), tomato (*Solanum Lycopersicum*), hazelnut (*Corylus avellana*), walnut (*Juglans regia*), banana (*Musa acuminata*), wheat (*Triticum aestivum*), latex (*Hevea brasiliensis*), tobacco (*Nicotiana tabacum*) and alcoholic beverages such as wine (grapes: *Vitis vinifera*) and beer (common hop: *Humulus lupulus*). Percentages represent sequence homology. ND: no data (43).

It is important to note that the allergic reactions to these plant foods are often triggered or exacerbated by cofactors such as exercise or nonsteroidal anti-inflammatory drugs (NSAIDs)

and are therefore variable which can make history taking more complex (44). Furthermore, these reactions are frequently more severe than the classic oral allergy syndrome (OAS) expected in food allergy related to sensitization to Bet v 1, the major allergen from birch pollen (45). This could be explained by the fact that ns-LTPs, unlike Bet v 1 homologues, resist to gastroduodenal proteolysis (46, 47) and thermal processing (48-50). However, it should also be kept in mind that in addition to plant food allergies, sensitization to Can s 3 might also explain cross-reactions to ns-LTP present in various sources such as *Hevea* latex (51-53), alcoholic beverages such as beer and wine (54, 55) and finally also tobacco (*Nicotinia tabaccum*) (9, 40, 56).

As described in table 1, thaumatin-like proteins (TLP), belonging to the PR-5 family, constitute another important group of components that might explain extensive cross-reactivity between cannabis and plant-derived food in European patients (10). These TLPs can be found in pollen and different foods such as NP24 from tomato (*Solanum lycopersicon*) (57), Cup a 3 from cypress (*Cupressus arizonica*) (58), Act d 2 kiwi fruit (*Actinidia deliciosa*) (59) and Mal d 2 from apple (*Malus domestica*) (60). For the discussion about the clinical significance of the panallergen profilin the reader is referred elsewhere (61, 62).

Diagnosis

Although a thorough history is absolutely mandatory for correct diagnosis, it appears that it is frequently pieced together from inadequate and incomplete descriptions and recall by patients reluctant to admit or simply denying illicit drug abuse. History might also be misleading because of misinterpretation or misconception of symptoms related to active or passive exposure. Presently, sensitization and allergy to cannabis is almost exclusively studied or documented by prick-prick skin testing. Prick-prick skin tests use a broad variety of raw materials such as macerated *C. sativa* leaves, buds and flowers (5, 8, 10, 14, 17, 18, 22, 26, 29, 32-34, 39, 63).

Needless to say that this approach is virtually impossible to standardize, mainly because of unpredictable variations in composition and potential contaminations with other allergens of the raw material.

A second method that can be applied to document cannabis allergy is quantification of serum specific IgE (sIgE) antibodies towards industrial hemp, an assay that can be obtained from Thermo Fisher Scientific (Uppsala, Sweden) but has not been thoroughly clinically validated. In our case-control study, a positive industrial hemp sIgE test result was observed in all 12 cannabis allergic patients but unfortunately also in 3 out of 8 pollen allergic patients without overt cannabis allergy (8). Using a whole protein extract, Larramendi *et al.* (10) found a positive sIgE result to a native cannabis extract in 21 out of 32 individuals who had their cannabis sensitization documented by a positive cannabis skin test. Herzinger *et al.* (63) found positive sIgE results to marihuana and/or hashish in 2 patients with professional exposure to cannabis. During the last two decades significant advances in biochemistry and molecular biology enabled the characterization, cloning and recombinant synthesis of relevant allergenic components and epitope-emulating peptides enabling quantification of serum sIgE antibodies to these components or sequential epitopes; a method known as component resolved diagnosis (CRD). In contrast to traditional sIgE tests, CRD does not rely upon whole extract preparations but upon single native or recombinant components (e.g. proteins or peptide components) (64, 65). CRD involves unique marker components to study the sensitization of patients towards a particular allergen and the presence of sIgE antibodies to cross-reactive components (e.g. profilins and CCD) that point to cross-reactivity. CRD not only allows one to discriminate between genuine allergy and cross-reactivity, but also enables to establish individual sensitization profiles which can be highly relevant in food allergy. In a study by Armentia *et al.* (29), sIgE antibodies against purified cannabis ns-LTP were demonstrable in over 95% of the

patients with a primary cannabis allergy. Recently, Rihs *et al.* (36) succeeded to clone Cannabis 3 from *C. sativa L ssp sativa cv Kompolti* and to study its IgE binding properties. Other *in vitro* tests that have been employed to document sensitization to cannabis are histamine release tests (63) and basophil activation tests (BAT) (8). In a preliminary study, we found BAT with a cannabis extract rich in ns-LTP to be absolutely discriminative between food allergic patients with and without cannabis allergy, larger studies are mandatory to confirm this finding. In healthy control individuals no basophil responses to this extract were demonstrable (8). Although cannabis challenge tests might add to the diagnosis of cannabis allergy (9, 14), it is highly unlikely this technique to enter mainstream use for obvious practical and ethical reasons.

Treatment

For the time being there is no cure for IgE-mediated *C. sativa* allergy nor for the cannabis-fruit/vegetable syndrome. Therefore, strict avoidance measures remain of utmost importance. These measures comprise a complete stop of further abuse of the drug, and avoidance of exposures to allergens implicated in the individual cross-reactivity syndrome. As reviewed by Ocampo *et al.* (6) hemp/cannabis desensitization has been described. However, these cases remain anecdotal and long-term follow-up data are lacking.

Natural history

The natural history of a cannabis allergy is currently unknown. Nevertheless, in our practice we observed several patients in who cannabis-related allergies were still evolving despite any further use of the drug. Longitudinal studies with larger numbers of participants are mandatory to establish whether cannabis allergy might have lighted an eternal flame.

Acknowledgements

This work was supported by the Agency for Innovation by Science and Technology (IWT) [grant number 140185]; ALVG is a Fellow of the FWO [1113617N].

The authors thank B. Van Camp, head of the Unit 'drug production', Central Drug Department, Directorate of organised crime, Belgian Federal Judicial police for providing cannabis sativa plants.

DGE is a Senior Clinical Researcher of the FWO (1800614N).

Conflict of interest: The authors report no conflict of interest with respect to this manuscript.

Author contributions:

Decuyper I.I.; interpretation of data, literature search, data analysis, writing

Van Gasse A.L.; proof reading final text

Cop Nathalie: proof reading final text

Sabato V.; proof reading final text

Faber M.A.; literature search, proof reading final text

Mertens Christel: interpretation diagnostic data, proof reading final text,

Bridts C.H.; data interpretation, proof reading final text

Hagendorens M.M.; proof reading final text

De Clerck L.; proof reading final text

Rihs H.P.; discussion diagnostic techniques, proof reading final text

Ebo D.G.; literature search, data analysis, writing, proof reading final text

References

1. Ferner M. Legal Marijuana Is The Fastest-Growing Industry In The U.S.: Report. Huffington post. Jan 26, 2015.
2. Berlinger J. Germany to legalize medicinal marijuana by 2017. CNN. May 4, 2016.
3. Furniere A. Medical cannabis plantation offers patients new perspectives. Flanderstoday. Aug 5, 2015.
4. Greydanus DE, Hawver EK, Greydanus MM, Merrick J. Marijuana: current concepts(dagger). *Front Public Health*. 2013;1:42. doi: 10.3389/fpubh.2013.00042. PubMed PMID: 24350211; PubMed Central PMCID: PMC3859982.
5. Stadtmauer G, Beyer K, Bardina L, Sicherer SH. Anaphylaxis to ingestion of hempseed (*Cannabis sativa*). *J Allergy Clin Immunol*. 2003;112(1):216-7.
6. Ocampo TL, Rans TS. Cannabis sativa: the unconventional "weed" allergen. *Ann Allergy Asthma Immunol*. 2015;114(3):187-92. doi: 10.1016/j.anai.2015.01.004. PubMed PMID: 25744904.
7. Gamboa P, Sanchez-Monge R, Sanz ML, Palacin A, Salcedo G, Diaz-Perales A. Sensitization to Cannabis sativa caused by a novel allergenic lipid transfer protein, Can s 3. *J Allergy Clin Immunol*. 2007;120(6):1459-60. doi: 10.1016/j.jaci.2007.07.052. PubMed PMID: 17881043.
8. Ebo DG, Swerts S, Sabato V, Hagendorens MM, Bridts CH, Jorens PG, et al. New food allergies in a European non-Mediterranean region: is Cannabis sativa to blame? *Int Arch Allergy Immunol*. 2013;161(3):220-8. doi: 10.1159/000346721. PubMed PMID: 23549061.
9. Armentia A, Castrodeza J, Ruiz-Munoz P, Martinez-Quesada J, Postigo I, Herrero M, et al. Allergic hypersensitivity to cannabis in patients with allergy and illicit drug users. *Allergol Immunopathol (Madr)*. 2011;39(5):271-9.
10. Larramendi CH, Lopez-Matas MA, Ferrer A, Huertas AJ, Pagan JA, Navarro LA, et al. Prevalence of sensitization to Cannabis sativa. Lipid-transfer and thaumatin-like proteins are relevant allergens. *Int Arch Allergy Immunol*. 2013;162(2):115-22. doi: 10.1159/000351068. PubMed PMID: 23921252.
11. Spiewak R, Gora A, Dutkiewicz J. Work-related skin symptoms and type I allergy among eastern-Polish farmers growing hops and other crops. *Ann Agric Environ Med*. 2001;8(1):51-6.
12. Kanceljak-Macan B, Zuskin E, Macan J. Organic aerosols and the development of allergic disorders. *Arh Hig Rada Toksikol*. 2004;55(2-3):213-20. PubMed PMID: 15285471.
13. Williams C, Thompstone J, Wilkinson M. Work-related contact urticaria to Cannabis sativa. *Contact Dermatitis*. 2008;58(1):62-3. doi: 10.1111/j.1600-0536.2007.01169.x. PubMed PMID: 18154568.
14. Vidal C, Fuente R, Iglesias A, Saez A. Bronchial asthma due to Cannabis sativa seed. *Allergy*. 1991;46(8):647-9. PubMed PMID: 1789408.
15. Herzinger T, Schopf P, Przybilla B, Rueff F. IgE-mediated hypersensitivity reactions to cannabis in laboratory personnel. *Int Arch Allergy Immunol*. 2011;156(4):423-6. doi: 10.1159/000324444. PubMed PMID: 21832832.
16. Liskow B, Liss JL, Parker CW. Allergy to marihuana. *Ann Intern Med*. 1971;75(4):571-3.
17. Freeman GL. Allergic skin test reactivity to marijuana in the Southwest. *West J Med*. 1983;138(6):829-31.
18. Stokes JR, Hartel R, Ford LB, Casale TB. Cannabis (hemp) positive skin tests and respiratory symptoms. *Ann Allergy Asthma Immunol*. 2000;85(3):238-40.
19. Torre FD, Limonta A, Molinari A, Masala E, Vercelloni S, Torre ED. Cannabaceae pollen in the atmosphere of Brianza, Northern Italy. *Eur Ann Allergy Clin Immunol*. 2007;39(1):9-11.
20. Singh AB, Shahi S. Aeroallergens in clinical practice of allergy in India- ARIA Asia Pacific Workshop report. *Asian Pac J Allergy Immunol*. 2008;26(4):245-56.
21. Mayoral M, Calderon H, Cano R, Lombardero M. Allergic rhinoconjunctivitis caused by Cannabis sativa pollen. *J Investig Allergol Clin Immunol*. 2008;18(1):73-4.

22. Prasad R, Verma SK, Dua R, Kant S, Kushwaha RA, Agarwal SP. A study of skin sensitivity to various allergens by skin prick test in patients of nasobronchial allergy. *Lung India*. 2009;26(3):70-3.
23. Mayoral M, Calderon H, Cano R, Lombardero M. Allergic rhinoconjunctivitis caused by Cannabis sativa pollen. *J Investig Allergol Clin Immunol*. 2008;18(1):73-4. PubMed PMID: 18361109.
24. Stokes JR, Hartel R, Ford LB, Casale TB. Cannabis (hemp) positive skin tests and respiratory symptoms. *Ann Allergy Asthma Immunol*. 2000;85(3):238-40. doi: 10.1016/S1081-1206(10)62473-8. PubMed PMID: 11030280.
25. Maloney ES, Brodkey, M.H. Hemp pollen sensitivity in Omaha. *Nebr Med J*. 1940;Volume 25:Pages 190-1.
26. Gamboa PM, Caceres O, Antepara I, Sanchez-Monge R, Ahrazem O, Salcedo G, et al. Two different profiles of peach allergy in the north of Spain. *Allergy*. 2007;62(4):408-14.
27. Tanaka H, Degawa M, Kawata E, Hayashi J, Shoyama Y. Identification of Cannabis pollens using an allergic patient's immunoglobulin E and purification and characterization of allergens in Cannabis pollens. *Forensic Sci Int*. 1998;97(2-3):139-53.
28. Van Loon LC. Occurrence and properties of plant pathogenesis-related proteins. *Pathogenesis-related proteins in plants* Eds SK Datta, S Muthukrishnan, CRC Press LLC, Boca Raton. 1999:1-19.
29. Armentia A, Herrero M, Martin-Armentia B, Rihs HP, Postigo I, Martinez-Quesada J. Molecular diagnosis in cannabis allergy. *J Allergy Clin Immunol Pract*. 2014;2(3):351-2. doi: 10.1016/j.jaip.2014.01.015. PubMed PMID: 24811031.
30. Faber MA VGA, Decuyper II, Uyttendaele A, Sabato V, Hagendorens MM, Bridts CH, De Clerck LS, Fernandez-Rivas M, Pascal M, Diaz-Perales A, Ebo DG. IgE-reactivity profiles to non-specific Lipid Transfer Proteins (ns-LTP) in a north-western European country. *Journal of Allergy and Clinical Immunology*. 2016. Epub July 14 2016
. doi: <http://dx.doi.org/10.1016/j.jaci.2016.06.016>.
31. Gao ZS, Yang ZW, Wu SD, Wang HY, Liu ML, Mao WL, et al. Peach allergy in China: a dominant role for mugwort pollen lipid transfer protein as a primary sensitizer. *J Allergy Clin Immunol*. 2013;131(1):224-6 e1-3. doi: 10.1016/j.jaci.2012.07.015. PubMed PMID: 22939759.
32. de Larramendi CH, Carnes J, Garcia-Abujeta JL, Garcia-Endrino A, Munoz-Palomino E, Huertas AJ, et al. Sensitization and allergy to Cannabis sativa leaves in a population of tomato (*Lycopersicon esculentum*)-sensitized patients. *Int Arch Allergy Immunol*. 2008;146(3):195-202. doi: 10.1159/000115887. PubMed PMID: 18268387.
33. Nayak AP, Green BJ, Sussman G, Berlin N, Lata H, Chandra S, et al. Characterization of Cannabis sativa allergens. *Ann Allergy Asthma Immunol*. 2013;111(1):32-7. doi: 10.1016/j.anai.2013.04.018. PubMed PMID: 23806457; PubMed Central PMCID: PMC3726218.
34. Metz-Favre C, Pauli G, Bessot JC, De BF. Molecular allergology in practice: an unusual case of LTP allergy. *Eur Ann Allergy Clin Immunol*. 2011;43(6):193-5.
35. Rojas Perez-Ezquerria P, Sanchez-Morillas L, Davila-Ferandez G, Ruiz-Hornillos FJ, Carrasco Garcia I, Herranz Manas M, et al. Contact urticaria to Cannabis sativa due to a lipid transfer protein (LTP). *Allergol Immunopathol (Madr)*. 2015;43(2):231-3. doi: 10.1016/j.aller.2013.10.002. PubMed PMID: 24657136.
36. Rihs HP, Armentia A, Sander I, Bruning T, Raulf M, Varga R. IgE-binding properties of a recombinant lipid transfer protein from Cannabis sativa. *Ann Allergy Asthma Immunol*. 2014;113(2):233-4. doi: 10.1016/j.anai.2014.05.022. PubMed PMID: 24954374.
37. Decuyper I, Ryckebosch H, Van Gasse AL, Sabato V, Faber M, Bridts CH, et al. Cannabis Allergy: What do We Know Anno 2015. *Arch Immunol Ther Exp (Warsz)*. 2015;63(5):327-32. doi: 10.1007/s00005-015-0352-z. PubMed PMID: 26178655.
38. Majmudar V, Azam NA, Finch T. Contact urticaria to Cannabis sativa. *Contact Dermatitis*. 2006;54(2):127.
39. Tessmer A, Berlin N, Sussman G, Leader N, Chung EC, Beezhold D. Hypersensitivity reactions to marijuana. *Ann Allergy Asthma Immunol*. 2012;108(4):282-4.

40. Faber M, Van Gasse A, Sabato V, Hagendorens MM, Bridts CH, De Clerck LS, et al. Marijuana allergy: beyond the joint. *J Invest Allergol Clin Immunol*. 2015;25(1):70-2. PubMed PMID: 25898704.
41. Egger M, Hauser M, Mari A, Ferreira F, Gadermaier G. The role of lipid transfer proteins in allergic diseases. *Curr Allergy Asthma Rep*. 2010;10(5):326-35.
42. Van Gasse AL, et al. L'allergie au cannabis: bien plus qu'un voyage stupéfiant. *Rev Fr Allergol*, <http://dxdoiorg/101016/jreval201401022>. 2014.
43. Boratyn GM, Schaffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. Domain enhanced lookup time accelerated BLAST. *Biol Direct*. 2012;7:12. doi: 10.1186/1745-6150-7-12. PubMed PMID: 22510480; PubMed Central PMCID: PMC3438057.
44. Asero R, Pravettoni V. Anaphylaxis to plant-foods and pollen allergens in patients with lipid transfer protein syndrome. *Current opinion in allergy and clinical immunology*. 2013;13(4):379-85. Epub 2013/02/22. doi: 10.1097/ACI.0b013e32835f5b07. PubMed PMID: 23426007.
45. Ebo DG, Stevens WJ. IgE-mediated food allergy--extensive review of the literature. *Acta Clin Belg*. 2001;56(4):234-47. doi: 10.1179/acb.2001.035. PubMed PMID: 11603253.
46. Cavatorta V, Sforza S, Aquino G, Galaverna G, Dossena A, Pastorello EA, et al. In vitro gastrointestinal digestion of the major peach allergen Pru p 3, a lipid transfer protein: molecular characterization of the products and assessment of their IgE binding abilities. *Mol Nutr Food Res*. 2010;54(10):1452-7.
47. Wijesinha-Bettoni R, Alexeev Y, Johnson P, Marsh J, Sancho AI, Abdullah SU, et al. The structural characteristics of nonspecific lipid transfer proteins explain their resistance to gastroduodenal proteolysis. *Biochemistry*. 2010;49(10):2130-9.
48. Scheurer S, Lauer I, Foetisch K, San Miguel MM, Retzek M, Hartz C, et al. Strong allergenicity of Pru av 3, the lipid transfer protein from cherry, is related to high stability against thermal processing and digestion. *J Allergy Clin Immunol*. 2004;114(4):900-7.
49. Sancho AI, Rigby NM, Zuidmeer L, Asero R, Mistrello G, Amato S, et al. The effect of thermal processing on the IgE reactivity of the non-specific lipid transfer protein from apple, Mal d 3. *Allergy*. 2005;60(10):1262-8.
50. Matejkova M, Zidkova J, Zidek L, Wimmerova M, Chmelik J, Sklenar V. Investigation of thermal denaturation of barley nonspecific lipid transfer protein 1 (ns-LTP1b) by nuclear magnetic resonance and differential scanning calorimetry. *J Agric Food Chem*. 2009;57(18):8444-52.
51. Beezhold DH, Hickey VL, Kostyal DA, Puhl H, Zuidmeer L, van Ree R, et al. Lipid transfer protein from *Hevea brasiliensis* (Hev b 12), a cross-reactive latex protein. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology*. 2003;90(4):439-45. Epub 2003/05/02. doi: 10.1016/s1081-1206(10)61830-3. PubMed PMID: 12722968.
52. Rihs HP, Rueff F, Lundberg M, Rozynek P, Barber D, Scheurer S, et al. Relevance of the recombinant lipid transfer protein of *Hevea brasiliensis*: IgE-binding reactivity in fruit-allergic adults. *Ann Allergy Asthma Immunol*. 2006;97(5):643-9.
53. Faber MA, Sabato V, Bridts CH, Nayak A, Beezhold DH, Ebo DG. Clinical relevance of the *Hevea brasiliensis* lipid transfer protein Hev b 12. *J Allergy Clin Immunol*. 2015;135(6):1645-8. doi: 10.1016/j.jaci.2014.12.1919. PubMed PMID: 25649078.
54. Jegou S, Douliez JP, Molle D, Boivin P, Marion D. Purification and structural characterization of LTP1 polypeptides from beer. *J Agric Food Chem*. 2000;48(10):5023-9. PubMed PMID: 11052772.
55. Asero R, Mistrello G, Roncarolo D, Casarini M, Falagiani P. Allergy to nonspecific lipid transfer proteins in Rosaceae: a comparative study of different in vivo diagnostic methods. *Ann Allergy Asthma Immunol*. 2001;87(1):68-71.
56. Stockli SS, Bircher AJ. Generalized pruritus in a patient sensitized to tobacco and cannabis. *J Dtsch Dermatol Ges*. 2007;5(4):303-4.
57. Sharma P, Singh AK, Singh BP, Gaur SN, Arora N. Allergenicity assessment of osmotin, a pathogenesis-related protein, used for transgenic crops. *J Agric Food Chem*. 2011;59(18):9990-5. doi: 10.1021/jf202265d. PubMed PMID: 21838306.

58. Breiteneder H. Thaumatin-like proteins -- a new family of pollen and fruit allergens. *Allergy*. 2004;59(5):479-81. doi: 10.1046/j.1398-9995.2003.00421.x. PubMed PMID: 15080826.
59. Bublin M, Mari A, Ebner C, Knulst A, Scheiner O, Hoffmann-Sommergruber K, et al. IgE sensitization profiles toward green and gold kiwifruits differ among patients allergic to kiwifruit from 3 European countries. *J Allergy Clin Immunol*. 2004;114(5):1169-75. doi: 10.1016/j.jaci.2004.07.016. PubMed PMID: 15536427.
60. Hsieh LS, Moos M, Jr., Lin Y. Characterization of apple 18 and 31 kd allergens by microsequencing and evaluation of their content during storage and ripening. *J Allergy Clin Immunol*. 1995;96(6 Pt 1):960-70. PubMed PMID: 8543755.
61. Ebo DG, Hagendorens MM, Bridts CH, De Clerck LS, Stevens WJ. Sensitization to cross-reactive carbohydrate determinants and the ubiquitous protein profilin: mimickers of allergy. *Clin Exp Allergy*. 2004;34(1):137-44. PubMed PMID: 14720274.
62. Santos A, van RR. Profilins: mimickers of allergy or relevant allergens? *Int Arch Allergy Immunol*. 2011;155(3):191-204.
63. Herzinger T, Schopf P, Przybilla B, Rueff F. IgE-mediated hypersensitivity reactions to cannabis in laboratory personnel. *Int Arch Allergy Immunol*. 2011;156(4):423-6.
64. Valenta R, Lidholm J, Niederberger V, Hayek B, Kraft D, Gronlund H. The recombinant allergen-based concept of component-resolved diagnostics and immunotherapy (CRD and CRIT). *Clin Exp Allergy*. 1999;29(7):896-904. PubMed PMID: 10383589.
65. Van Gasse AL, Mangodt EA, Faber M, Sabato V, Bridts CH, Ebo DG. Molecular allergy diagnosis: status anno 2015. *Clin Chim Acta*. 2015;444:54-61. doi: 10.1016/j.cca.2015.02.012. PubMed PMID: 25681645.