

This item is the archived peer-reviewed author-version of:

Personal and perceived peer use and attitudes towards use of non-prescribed prescription sedatives and sleeping pills among university students in seven European countries

Reference:

Lehne Gesa, Zeeb Hajo, Pischke Claudia R., Mikolajczyk Rafael, Bewick Bridgette M., McAlaney John, Dempsey Robert C., Van Hal Guido F., Stock Christiane, Akvardar Yildiz,- Personal and perceived peer use and attitudes towards use of non-prescribed prescription sedatives and sleeping pills among university students in seven European countries

Addictive behaviors - ISSN 0306-4603 - 87(2018), p. 17-23

Full text (Publisher's DOI): <https://doi.org/10.1016/J.ADDBEH.2018.06.012>

To cite this reference: <https://hdl.handle.net/10067/1546950151162165141>

Accepted Manuscript

Personal and perceived peer use and attitudes towards use of non-prescribed prescription sedatives and sleeping pills among university students in seven European countries

Gesa Lehne, Hajo Zeeb, Claudia R. Pischke, Rafael Mikolajczyk, Bridgette M. Bewick, John McAlaney, Robert C. Dempsey, Guido Van Hal, Christiane Stock, Yildiz Akvardar, Ondrej Kalina, Olga Orosova, Ines Aguinaga-Ontoso, Francisco Guillen-Grima, Stefanie M. Helmer



PII: S0306-4603(18)30629-4
DOI: doi:[10.1016/j.addbeh.2018.06.012](https://doi.org/10.1016/j.addbeh.2018.06.012)
Reference: AB 5597
To appear in: *Addictive Behaviors*
Received date: 21 January 2018
Revised date: 5 June 2018
Accepted date: 10 June 2018

Please cite this article as: Gesa Lehne, Hajo Zeeb, Claudia R. Pischke, Rafael Mikolajczyk, Bridgette M. Bewick, John McAlaney, Robert C. Dempsey, Guido Van Hal, Christiane Stock, Yildiz Akvardar, Ondrej Kalina, Olga Orosova, Ines Aguinaga-Ontoso, Francisco Guillen-Grima, Stefanie M. Helmer , Personal and perceived peer use and attitudes towards use of non-prescribed prescription sedatives and sleeping pills among university students in seven European countries. *Ab* (2017), doi:[10.1016/j.addbeh.2018.06.012](https://doi.org/10.1016/j.addbeh.2018.06.012)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Personal and perceived peer use and attitudes towards use of non-prescribed prescription sedatives and sleeping pills among university students in seven European countries
 Gesa Lehne^{a,b,*}, Hajo Zeeb^{b,c}, Claudia R. Pischke^{c,d}, Rafael Mikolajczyk^e, Bridgette M. Bewick^f, John McAlaney^g, Robert C. Dempsey^h, Guido Van Halⁱ, Christiane Stock^j, Yildiz Akvardar^k, Ondrej Kalina^l, Olga Orosova^l, Ines Aguinaga-Ontoso^m, Francisco Guillen-Grima^{m,n}, Stefanie M. Helmer^{c,o}

^aDepartment of Social Epidemiology, Institute of Public Health and Nursing Research, University of Bremen, 28359 Bremen, Germany

^bHealth Sciences Bremen, University of Bremen, Bremen, Germany

^cDepartment Prevention and Evaluation, Leibniz Institute for Prevention Research and Epidemiology – BIPS, 28359 Bremen, Germany

^dInstitute for Medical Sociology, Centre for Health and Society, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

^eInstitute of Medical Epidemiology, Biometry and Informatics, Martin-Luther-University Halle-Wittenberg, 06112 Halle (Saale), Germany

^fSchool of Medicine, Leeds Institute of Health Sciences, University of Leeds, Leeds, LS1 3HE, United Kingdom

^gDepartment of Psychology, Research Centre for Behaviour Change, Bournemouth University, Bournemouth, BH12 5BB, United Kingdom

^hStaffordshire Centre for Psychological Research & Centre for Health Psychology, School of Life Sciences & Education, Staffordshire University, Stoke-on-Trent, ST4 2DF, United Kingdom

ⁱEpidemiology and Social Medicine, University of Antwerp, 2610 Antwerp, Belgium

^jUnit for Health Promotion Research, University of Southern Denmark, 6700 Esbjerg, Denmark

^kDepartment of Psychiatry, Marmara University Medical School, 34722 Istanbul, Turkey

^lDepartment of Educational Psychology & Health Psychology, PJ Safarik University in Košice, 041 80 Košice, Slovak Republic

^mDepartment of Health Sciences, Public University of Navarra, 31008 Pamplona, Spain

ⁿPreventive Medicine, Clinica Universidad de Navarra, 31008 Pamplona, Spain

^oInstitute for Social Medicine, Epidemiology, and Health Economics, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, 13353 Berlin, Germany

*Corresponding author.

Gesa Lehne, MA

Department of Social Epidemiology
Institute of Public Health and Nursing Research
University of Bremen
Grazer Strasse 2a
28359 Bremen
Germany
gesa.lehne@uni-bremen.de

Hajo Zeeb, zeeb@bips.uni-bremen.de
Claudia R. Pischke, claudia.pischke@leibniz-bips.de
Rafael Mikolajczyk, rafael.mikolajczyk@uk-halle.de
Bridgette M. Bewick, B.M.Bewick@leeds.ac.uk
John McAlaney, jmcalaney@bournemouth.ac.uk
Robert C. Dempsey, Robert.Dempsey@staffs.ac.uk
Guido Van Hal, guido.vanhal@uantwerpen.be
Christiane Stock, cstock@health.sdu.dk
Yildiz Akvardar, yildiz.akvardar@marmara.edu.tr
Ondrej Kalina, ondrej.kalina@upjs.sk
Olga Orosova, olga.orosova@upjs.sk
Ines Aguinaga-Ontoso, ines.aguinaga@unavarra.es
Francisco Guillen-Grima, frguillen@unav.es
Stefanie M. Helmer, stefanie.helmer@charite.de

Abstract

Introduction: The use of non-prescribed prescription sedatives and sleeping pills (NPPSSP) among university students has been described as an important public health issue. However, the impact of perceived social norms on students' use and attitudes towards use of NPPSSP is still unclear. Our aim was to investigate whether perceptions of peer use and approval of use are associated with students' personal use and approval of NPPSSP use.

Methods: Cross-sectional data from the Social Norms Intervention for the Prevention of Polydrug Use (SNIPE) project containing 4,482 university students from seven European countries were analyzed to investigate self-other discrepancies regarding personal use and attitudes towards NPPSSP use. Associations between personal and perceived peer use and between personal and perceived approval of use were examined using multivariable logistic regression.

Results: The majority (51.0%) of students perceived their peers' NPPSSP use to be higher than their personal use. 92.6% of students perceived their peers' approval of NPPSSP use to be identical or higher than their personal approval. Students perceiving that the majority of peers had used NPPSSP at least once displayed higher odds for personal lifetime use (OR: 1.95, 95% CI: 1.49-2.55). Perceived peer approval of NPPSSP use was associated with higher odds for personal approval (OR: 5.49, 95% CI: 4.63-6.51).

Conclusions: Among European university students, perceiving NPPSSP use and approval of use to be the norm was positively associated with students' personal NPPSSP use and approval of use, respectively. Interventions addressing perceived social norms may prevent or reduce NPPSSP use among university students.

Final trial registration number: DRKS00004375 on the 'German Clinical Trials Register'.

Keywords: university students; non-medical use; sedatives; sleeping pills; perceptions; social norms

1. Introduction

The non-medical use of prescription drugs, particularly among young adults, has been recognized as an important public health issue worldwide (Martins & Ghandour, 2017). The misuse of several prescription drugs, such as stimulants, opioids, or tranquilizers, is associated with a high potential for addiction and other serious physical and psychosocial consequences (United Nations Office on Drugs and Crime, 2011). However, prescription drugs are often perceived to be safer, and more socially acceptable than most illicit drugs, because they are produced by pharmaceutical companies and usually prescribed by physicians (Bodenlos, Malorby, Noonan, Mayrsohn, & Mistler, 2014; Compton & Volkow, 2006; Hildt, Franke, & Lieb, 2011; Martins & Ghandour, 2017).

The non-medical use of prescription drugs among university students may serve as a coping strategy to manage the demands of university life and to achieve a better work-life balance (Hildt, Lieb, & Franke, 2014; Jensen, Forlini, Partridge, & Hall, 2016; Maier, Liechti, Herzig, & Schaub, 2013). The phenomenon of taking prescription drugs for the purpose of improving cognitive performance (e.g., alertness, concentration, or memory) has been termed *pharmacological cognitive enhancement* or *brain doping* (Partridge, Bell, Lucke, Yeates, & Hall, 2011). Further, evidence indicates that university students use sedatives to improve sleep or relax after stressful days, thus aiming to improve cognitive performance the next day. This is also referred to as *indirect cognitive enhancement* (Maier, et al., 2013; Maier & Schaub, 2015). Academic performance-enhancing drugs and sedatives are often used in combination: while performance-enhancing drugs are used to achieve the highest possible performance level during the day, sedatives are used to aid relaxation (Maier, et al., 2013).

Typically, peers have a significant impact on young adults' behaviors and their attitudes, and people tend to adapt their personal behavior to match that of their peers (Borsari & Carey, 2001). However, a growing body of evidence indicates that young people's perceptions of their peers' behaviors (*descriptive norms*) and attitudes towards behaviors (*injunctive norms*) are often inaccurate (Berkowitz, 2005; Perkins, 2003). University students tend to falsely believe that their peers behave or approve of behaviors differently from actual prevailing norms (*misperceptions*) (Berkowitz, 2005; Perkins, 2003), and from their personal behavior and approval of behavior (*self-other discrepancies*) (Borsari & Carey, 2001). Young people generally overestimate how riskily their peers behave. These misperceptions of other's behavior or attitudes towards behavior represent the basis for the adaptation of personal behavior and attitude towards the perceived norm (Berkowitz, 2005). Most research on misperceptions of health-related behaviors among university students originated in the U.S.A. and particularly refers to descriptive norms regarding alcohol consumption (Borsari & Carey, 2001; Perkins, 2014). In recent years, these findings were replicated in Europe (McAlaney, Bewick, & Hughes, 2011; McAlaney, et al., 2015). These studies show that exaggerated perceptions of peer alcohol consumption are associated with increased personal alcohol

consumption among university students (Borsari & Carey, 2001; McAlaney, et al., 2011; McAlaney, et al., 2015; Perkins, 2014). There is further evidence on university students' misperceptions of their peers' use of tobacco and illicit substances (e.g., marijuana, cocaine, ecstasy, and amphetamines) (Arbour-Nicitopoulos, Kwan, Lowe, Taman, & Faulkner, 2010; Bertholet, Faouzi, Studer, Daeppen, & Gmel, 2013; Dempsey, et al., 2016; Helmer, et al., 2014; Kilmer, et al., 2006; Martens, et al., 2006; Perkins, Meilman, Leichter, Cashin, & Presley, 1999; Pischke, et al., 2015), as well as regarding risky sexual behavior (Martens, et al., 2006).

Several studies have examined misperceptions or self-other discrepancies about the non-medical use of prescription drugs, as well as associations between descriptive norms and personal use, particularly regarding prescription stimulants (Helmer, et al., 2016; Kilmer, Geisner, Gasser, & Lindgren, 2015; McCabe, 2008; Sanders, Stogner, Seibert, & Miller, 2014; Silvestri & Correia, 2016), with only one study, to date, investigating prescription sedative use (Sanders, et al., 2014). Perceived approval among peers for the non-prescribed use of prescription stimulants at the same university (Helmer, et al., 2016) and perceived approval among close friends, or by the typical university student or parents (Silvestri & Correia, 2016), were positively associated with personally approving such substances among university students. The role of perceived injunctive norms regarding non-medical use of prescription sedatives, however, has not been investigated so far.

The present study aimed to investigate self-other discrepancies regarding the use and attitudes towards using non-prescribed prescription sedatives and sleeping pills (NPPSSP) in a sample of university students from seven European countries. We also aimed to investigate if perceptions of peer use (perceived descriptive norm) and peer approval of use (perceived injunctive norm) were associated with personal use and approval of NPPSSP use in our study population.

To clarify the terminology employed in this study, NPPSSP is used to describe the non-prescribed use of sedatives and sleeping pills which are only available by prescription. This does not include the use of non-prescription products, such as herbal sedatives, which can be acquired without prescription.

2. Material and Methods

2.1 Data

This analysis is based on data from the 'Social Norms Intervention for the prevention of Polydrug use' (SNIPE) project funded by the European Commission (LS/2009-2010/DPIP/AG). SNIPE was a cross-national study including students from universities in Belgium, Denmark, Germany, the Slovak Republic, Spain, Turkey, and the United Kingdom (UK). An overview of the SNIPE study is provided by Pischke and colleagues (2012). In brief, SNIPE aimed to test the feasibility of a web-based, personalized 'social norms'-feedback for the prevention of licit and illicit substance use for

European university students. Participants were recruited from one or more designated intervention and delayed-intervention control universities (21 sites in total) (McAlaney, et al., 2015). Recruitment methods aimed at increasing students' registrations on the survey website varied between countries and included, inter alia, emails, classroom announcements, social media, and printed flyers. Students who registered on the website received an email including a hyperlink to the survey webpage. Study participation was voluntary, and participants' information was pseudonymized. For the analysis reported in this manuscript, baseline data from both, students at intervention and students at delayed-intervention control universities, were considered. Statistical analysis was conducted on an anonymized dataset. For each site participating in the SNIPE project, ethical approval was obtained from the respective responsible authorities. Participants answered questions on their personal use of licit (i.e., alcohol, tobacco), and illicit substances (e.g., cocaine, ecstasy, amphetamines), as well as on their personal use of non-prescribed prescription substances to improve academic performance and NPPSSP. Further questions related to the students' personal attitudes towards use of the aforementioned substances. Moreover, perceptions of peer substance use and attitudes towards substance use were assessed. Demographic questions, such as on the participants' age, sex, migrant status, and living situation (living with or without other students), were also included.

2.2 Measurements

Students' personal use of NPPSSP was measured by asking how often they used sedatives or sleeping pills which were not prescribed, followed by a list of registered local trade names of prescription sedatives and sleeping pills as examples (e.g., diazepam, alprazolam, flunitrazepam, midazolam, stilnoct). Perceptions of peer NPPSSP use (perceived descriptive norm) were assessed by asking students how often in the last two months they think most (at least 51%) of the [female in case of a female respondent/male in case of a male respondent] students at their university have used sedatives or sleeping pills which were not prescribed, followed by a list of registered local trade names of prescription sedatives and sleeping pills as examples (e.g., diazepam, alprazolam, flunitrazepam, midazolam, stilnoct). These questions were tailored to the same sex and university of the respondents. Response options for both questions were 'Never in my/their life', 'Have used but not in the last two months', 'Once in the last two months', 'Twice in the last two months', 'Once every two weeks in the last two months', 'Weekly', 'Twice a week', 'Thrice a week', 'Four times a week', and 'Every day or nearly every day'. Furthermore, information about students' personal attitude towards NPPSSP use was collected by asking: "Which of the following best describes your attitude to using each of these substances?". Concerning students' perceptions of attitudes towards using NPPSSP among their peers (perceived injunctive norm), respondents were asked: "Which of

the following do you think best describes the attitude of most (at least 51%) of the [female/male] students at your university to the use of each of these substances?”. Response options for both questions were ‘Never ok to use’, ‘Ok to use occasionally if it doesn’t interfere with work or study’, ‘Ok to use frequently if it doesn’t interfere with work or study’, ‘Ok to use occasionally even if it does interfere with work or study’, and ‘Ok to use frequently if that is what the person wants to do’. Country, sex, age, year of study, and living situation were considered as potential determinants of NPPSSP use/attitude towards NPPSSP use.

2.3 Statistical analysis

First, frequencies of personal NPPSSP use and attitudes towards NPPSSP use were calculated and 95% bootstrap confidence intervals based on 1,000 bootstrap samples were estimated for each country, separately. Second, participants’ self-other discrepancies were classified into three groups to differentiate between students who perceived the NPPSSP use and approval of NPPSSP use of the majority of their same-sex peers as higher, identical or lower as their personal use and approval of use. Third, two binary multivariable logistic regression analyses were conducted to examine associations between perceived and personal NPPSSP use (descriptive norms model), and perceived and personal attitudes towards NPPSSP use (injunctive norms model). In the descriptive norms model, country, sex, age, year of study, living situation, perceived NPPSSP use, and personal attitude towards NPPSSP use were included as independent variables. In the injunctive norms model, all demographic variables, perceived attitude towards NPPSSP use, and personal NPPSSP use were included as independent variables. In both models, all variables were entered simultaneously (enter method). Age was included as a continuous variable, and all other variables were considered as categorical variables. Categorical variables with more than two categories (i.e., country, year of study, living situation) were each converted into a set of dichotomous variables using dummy coding. Both models were checked for the presence of multicollinearity. Tolerance (TOL) values for both models ranged from 0.90 to 1.00 indicating absence of multicollinearity between independent variables. To investigate whether sex or country moderates the associations between perception and personal NPPSSP use/attitude towards NPPSSP use, the two relevant interaction terms were added to both regression models. For significant interaction terms ($p < 0.05$), stratified analyses were conducted. All statistical analyses were performed using SPSS for windows, version 22.0.

3. Results

The SNIPE study included a total of 4,482 university students (71.4% female, mean age: 22.4 years). The Slovak Republic ($n=1,938$, 43.2%) contributed the highest number of students, followed by Turkey ($n=858$, 19.1%), Germany ($n=504$, 11.2%), Denmark ($n=464$, 10.4%), Belgium ($n=426$,

9.5%), Spain (n=185, 4.1%), and the UK (n=107, 2.4%). A detailed description of the sample characteristics is provided by Helmer et al. (2014). Information on sex and NPPSSP use was provided by 4,412 students, and 4,284 additionally answered the question regarding their attitude towards using NPPSSP.

Across all participating countries, 9.1% of the students reported having used NPPSSP at least once in life. Lifetime prevalence rates of NPPSSP use varied from 4.0% of females and 2.3% of males in Belgium to 12.5% of females and 18.2% of males in the UK. Across all countries, most students stated that 'it is never okay to use' NPPSSP with rates varying from 56.8% of females in Germany and 62.5% of males in the UK to 84.7% of females and 91.2% of males in Turkey (Table 1).

ACCEPTED MANUSCRIPT

Table 1 Personal NPPSSP use and attitude towards NPPSSP use by country and sex (% and 95% bootstrap CI)

	Belgium		Denmark		Germany		Slovak Republic	
<i>NPPSSP use (n=4,412)</i>	<i>Male (n=86)</i>	<i>Female (n=321)</i>	<i>Male (n=100)</i>	<i>Female (n=353)</i>	<i>Male (n=207)</i>	<i>Female (n=295)</i>	<i>Male (n=393)</i>	<i>Female (n=1,524)</i>
Used in the last two months	1.2 (0.0-3.8)	1.2 (0.3-2.6)	1.0 (0.0-3.3)	1.7 (0.6-3.3)	2.9 (0.9-5.4)	3.1 (1.2-5.2)	1.5 (0.5-2.9)	2.7 (1.9-3.5)
Used at least once in life	2.3 (0.0-5.8)	4.0 (2.1-6.3)	9.0 (3.6-14.7)	5.9 (3.4-8.6)	11.1 (6.7-15.6)	10.2 (6.6-13.7)	6.4 (3.9-8.9)	11.6 (10.0-13.2)
<i>Attitude towards NPPSSP use (n=4,284)</i>	<i>Male (n=85)</i>	<i>Female (n=316)</i>	<i>Male (n=95)</i>	<i>Female (n=348)</i>	<i>Male (n=203)</i>	<i>Female (n=292)</i>	<i>Male (n=384)</i>	<i>Female (n=1,489)</i>
Never ok to use	83.5 (75.0-91.5)	72.2 (67.1-77.1)	65.3 (55.9-74.0)	75.3 (70.7-79.6)	64.0 (57.1-70.4)	56.8 (51.0-62.6)	83.3 (79.4-86.9)	73.1 (70.8-75.4)
Ok to use if it doesn't interfere with work or study ^a	15.3 (7.9-23.5)	26.2 (21.6-31.6)	28.4 (20.6-37.5)	21.3 (17.3-25.9)	30.0 (23.9-37.0)	38.0 (32.4-43.7)	15.1 (11.7-18.8)	25.6 (23.4-27.8)
Ok to use ^b	1.2 (0.0-3.8)	1.3 (0.3-2.8)	6.3 (2.0-11.8)	3.4 (1.7-5.3)	5.9 (2.7-9.5)	5.1 (2.7-7.9)	1.6 (0.5-3.1)	1.3 (0.8-2.0)
	Spain		Turkey		UK			
<i>NPPSSP use (n=4,412)</i>	<i>Male (n=52)</i>	<i>Female (n=132)</i>	<i>Male (n=398)</i>	<i>Female (n=446)</i>	<i>Male (n=33)</i>	<i>Female (n=72)</i>		
Used in the last two months	1.9 (0.0-6.7)	4.5 (1.5-8.3)	2.0 (0.8-3.5)	2.5 (1.1-3.9)	12.1 (2.9-24.2)	4.2 (0.0-9.2)		
Used at least once in life	11.5 (3.8-20.5)	12.1 (6.4-18.2)	5.5 (3.6-7.9)	9.9 (7.2-12.6)	18.2 (6.5-31.4)	12.5 (5.5-21.1)		
<i>Attitude towards NPPSSP use (n=4,284)</i>	<i>Male (n=51)</i>	<i>Female (n=126)</i>	<i>Male (n=375)</i>	<i>Female (n=419)</i>	<i>Male (n=32)</i>	<i>Female (n=69)</i>		
Never ok to use	64.7 (51.1-78.3)	65.9 (57.6-73.8)	91.2 (88.4-93.9)	84.7 (81.1-88.1)	62.5 (45.7-80.0)	73.9 (62.9-83.8)		
Ok to use if it doesn't interfere with work or study ^a	33.3 (20.0-46.9)	31.7 (23.7-39.8)	6.1 (3.9-8.8)	13.6 (10.3-17.1)	34.4 (17.7-51.9)	24.6 (15.2-34.8)		
Ok to use ^b	2.0 (0.0-6.9)	2.4 (0.0-5.5)	2.7 (1.0-4.5)	1.7 (0.5-3.0)	3.1 (0.0-10.0)	1.4 (0.0-4.6)		

^a 'Ok to use occasionally if it doesn't interfere with work or study' and 'Ok to use frequently if it doesn't interfere with work or study' were collapsed into 'Ok to use if it doesn't interfere with work or study'.

^b 'Ok to use occasionally even if it does interfere with work or study' and 'Ok to use frequently if that is what the person wants to do' were combined into 'Ok to use'.

In all countries, except for Denmark (45.4%) and Turkey (43.9%), more than half (54.8%) of the students thought that at least 51% of their same sex-peers had used NPPSSP at least once in their life. Overall, 51.0% perceived their peers' NPPSSP use to be higher than their personal NPPSSP use, 46.0% to be identical, and 3.0% to be lower. With regard to attitudes towards NPPSSP use, 45.1% perceived that the majority of their peers approved of NPPSSP use. Overall, the majority of students perceived that the peer approval towards NPPSSP use was identical (62.9%) or higher (29.7%) than their personal approval (Table 2).

Table 2 Differences between personal NPPSSP use/attitude towards NPPSSP use and perceived NPPSSP use/ attitude towards NPPSSP use of the majority of peers of the same sex and university (self-other discrepancies)

	Lifetime NPPSSP use (%) (n=4,310)	Positive attitude towards NPPSSP use ^a (%) (n=4,178)
Majority of same-sex peers < personal	3.0	7.4
Majority of same-sex peers = personal	46.0	62.9
Majority of same-sex peers > personal	51.0	29.7

^a'Ok to use occasionally if it doesn't interfere with work or study', 'Ok to use frequently if it doesn't interfere with work or study', 'Ok to use occasionally even if it does interfere with work or study', and 'Ok to use frequently if that is what the person wants to do'.

After controlling for students' country, sex, age, year of study, living situation, and attitude towards NPPSSP use, the perception that the majority of same-sex peers had used NPPSSP at least once in their life was significantly associated with a higher likelihood for personal lifetime NPPSSP use (OR: 1.95, 95% CI: 1.49-2.55) (Table 3). Moreover, after controlling for all demographic variables and NPPSSP use, perceived peer approval of NPPSSP use was associated with higher odds for personal approval of NPPSSP use (OR: 5.49, 95% CI: 4.63-6.51) (Table 4).

Table 3 Associations between personal NPPSSP use and perceived lifetime NPPSSP use of peers, personal attitude towards NPPSSP use, country, age, sex, year of study, and living situation – results of a binary logistic regression (descriptive norms model)

Variables	Ever personally used NPPSSP	
	OR	(95% CI)
<i>Perceived peer NPPSSP use</i>		
Never used NPPSSP (reference)	1.00	
Ever used NPPSSP	1.95	(1.49-2.55)
<i>Personal attitude towards NPPSSP use</i>		
Never ok to use NPPSSP (reference)	1.00	
Ok to use NPPSSP ^a	7.42	(5.81-9.49)
<i>Country</i>		
Slovak Republic (reference)	1.00	
Belgium	0.24	(0.14-0.43)
Denmark	0.32	(0.20-0.52)
Germany	0.47	(0.32-0.70)
Spain	0.70	(0.41-1.22)
Turkey	0.99	(0.70-1.40)
UK	1.01	(0.52-1.94)
Age (in years)	1.04	(1.01-1.07)
<i>Sex</i>		
Female (reference)	1.00	
Male	0.82	(0.63-1.09)
<i>Year of study</i>		
1 st (reference)	1.00	
2 nd	0.78	(0.57-1.07)
3 rd	1.13	(0.83-1.54)
4 th	0.89	(0.60-1.31)
5 th	0.66	(0.39-1.10)
> 5 th	0.70	(0.35-1.41)
<i>Living situation</i>		
With other students (reference)	1.00	
Alone or with partner	2.04	(1.45-2.85)
With parents	1.06	(0.80-1.40)
Other	1.74	(0.94-3.23)

^a ‘Ok to use occasionally if it doesn’t interfere with work or study’, ‘Ok to use frequently if it doesn’t interfere with work or study’, ‘Ok to use occasionally even if it does interfere with work or study’, and ‘Ok to use frequently if that is what the person wants to do’.

Table 4 Associations between personal attitude towards NPPSSP use and perceived attitude of peers, personal NPPSSP use, country, age, sex, year of study, and living situation – results of a binary logistic regression (injunctive norms model)

Variables	Positive attitude towards NPPSSP use ^a	
	OR	(95% CI)
<i>Perceived peer attitude towards NPPSSP use</i>		
Never ok to use NPPSSP (reference)	1.00	
Ok to use NPPSSP ^a	5.49	(4.63-6.51)
<i>Personal NPPSSP use</i>		
Never used NPPSSP (reference)	1.00	
Ever used NPPSSP	7.03	(5.45-9.06)
<i>Country</i>		
Slovak Republic (reference)	1.00	
Belgium	0.99	(0.74-1.30)
Denmark	2.04	(1.49-2.80)
Germany	2.59	(2.00-3.36)
Spain	1.59	(1.09-2.34)
Turkey	0.54	(0.41-0.71)
UK	1.20	(0.72-1.99)
Age (in years)	0.97	(0.95-0.99)
<i>Sex</i>		
Female (reference)	1.00	
Male	0.84	(0.70-1.02)
<i>Year of study</i>		
1 st (reference)	1.00	
2 nd	0.84	(0.68-1.04)
3 rd	0.99	(0.78-1.24)
4 th	1.02	(0.77-1.37)
5 th	1.04	(0.73-1.47)
> 5 th	0.97	(0.60-1.54)
<i>Living situation</i>		
With other students (reference)	1.00	
Alone or with partner	0.82	(0.64-1.06)
With parents	1.06	(0.87-1.28)
Other	0.63	(0.39-1.02)

^a ‘Ok to use occasionally if it doesn't interfere with work or study’, ‘Ok to use frequently if it doesn't interfere with work or study’, ‘Ok to use occasionally even if it does interfere with work or study’, and ‘Ok to use frequently if that is what the person wants to do’.

Interaction terms in the descriptive norms model provided no evidence that the effect of perception on personal lifetime NPPSSP use was modified by country or sex. In terms of injunctive norms, significant interaction terms suggested that the effect of perception on personal attitude towards NPPSSP use was significantly modified by country, but not by sex. A stratified analysis of injunctive norms by country showed that the association between perception of peer approval and personal approval was significant for all countries, except for the UK (Table 5).

Table 5 Association between personal attitude towards NPPSSP use and perceived attitude of peers stratified by country adjusted for personal NPPSSP use, age, sex, year of study, and living situation

Country	Positive attitude towards NPPSSP use ^a	
	OR	(95% CI)
Slovak Republic	6.02	(4.64-7.81)
Belgium	2.79	(1.60-4.87)
Denmark	16.40	(9.37-28.73)
Germany	4.11	(2.69-6.29)
Spain	3.52	(1.66-7.47)
Turkey	6.41	(3.80-10.80)
UK	1.79	(0.52-6.10)

^a‘Ok to use occasionally if it doesn't interfere with work or study’, ‘Ok to use frequently if it doesn't interfere with work or study’, ‘Ok to use occasionally even if it does interfere with work or study’, and ‘Ok to use frequently if that is what the person wants to do’.

4. Discussion

In the present study with European students, we investigated self-other discrepancies regarding the use and attitudes towards the use of NPPSSP. In addition, we evaluated whether perceptions of peer use (perceived descriptive norm) and peer approval of use (perceived injunctive norm) were associated with personal use and approval of NPPSSP use. In our study, students on average perceived the NPPSSP use of their peers to be higher than their personal use and attitudes towards the use to be identical or more positive than their personal attitudes. Both, perceived descriptive and injunctive norms of peers, were associated with students' personal use and attitudes towards the use of NPPSSP, respectively.

To date, there are few studies on the use of NPPSSP among students. The only study that examined perceptions with respect to prescription sedatives by Sanders and colleagues (2014) found that 65.7% of students perceived the recreational use of prescription sedatives to be the norm among their peers despite only 2.6% of the sample reporting recreational use of these substances during the last month. More than a third of participants overestimated (26.3%) or extremely overestimated (10.2%) their peers' use, and recreational users of prescription sedatives were more likely to overestimate their peers' use of these substances (Sanders, et al., 2014). These findings are in line with our study. The results reported by Sanders and colleagues (2014), however, are based on bivariate analyses and thus did not account for further potential determinants of students' personal prescription sedative use, such as sex or age.

Our study extends the limited evidence regarding the association of perceived descriptive norms of peers with university students' personal use of NPPSSP. Indeed, our study adds to the existing evidence by revealing self-other discrepancies regarding NPPSSP use in a large sample of university students from various universities across Europe. Across all countries participating in the SNIPE study, the majority of students perceived their peers' use to be higher than their personal

use. Furthermore, we demonstrated associations between perceived peer use and students' personal use while controlling for other potential determinants of NPPSSP use ensuring further methodological rigor to our study.

The present study is the first to demonstrate discrepancies between personal and perceived peer injunctive norms regarding NPPSSP use by investigating self-other discrepancies and associations between perceived injunctive norms and students' personal approval of NPPSSP use. To date, associations between perceived injunctive norms and personal approval of using non-prescribed prescription substances have only been investigated for stimulants (Helmer, et al., 2016; Silvestri & Correia, 2016), not for sedatives or sleeping pills. Silvestri and Correia (2016), analyzing data from 959 U.S. undergraduate students, found that students' personal approval of non-medical prescription stimulant use was positively correlated with perceived approval among what students perceived to be a typical university student, close friends, as well as parents. However, the correlations between perceived parental and close friend approval with personal approval were moderate in strength with weak associations between perceived typical student approval and personal approval. This suggests that more proximal referent groups, rather than students' broader group affiliations, could be important in determining personal approval of stimulant use. Another study by Helmer and colleagues (2016), also using data from the SNIPE study, found that 38.7% of students perceived their peers to be more approving of using non-prescribed prescription stimulants to improve their academic performance than themselves. Their multivariable analysis also revealed an association between perceived peer and personal approval of using these substances. In our study, an association between perceived injunctive norms of peers and students' personal approval of using NPPSSP was found for all countries participating in the SNIPE project, except for the UK, with its comparatively small sample size.

The findings of this study align with previous observations that university students' exaggerated perceptions of peer norms also exist for prescription substances which are less commonly used and socially accepted than, for example alcohol, tobacco, or cannabis (Helmer, et al., 2016; Kilmer, et al., 2015; McCabe, 2008; Perkins, et al., 1999; Sanders, et al., 2014; Silvestri & Correia, 2016). Increased interest in the non-medical use of prescription drugs to the public and the media (Partridge, et al., 2011) may create the impression that approving and using these substances is much more common than it is in reality (McCabe, 2008; Sanders, et al., 2014). Perceiving prescription drugs to be safer, and socially acceptable because of their production by pharmaceutical companies and their prescription by physicians (Bodenlos, et al., 2014; Compton & Volkow, 2006; Hildt, et al., 2011; Martins & Ghandour, 2017) may also explain exaggerated peer norms.

The identification of perceived descriptive and injunctive norms of peers as significant predictors of students' NPPSSP use and approval of use provides empirical arguments for the important role of social norms for personal behaviors and approval of behaviors. In line with social norms theory (Berkowitz, 2005; Perkins, 2003), our findings may indicate that exaggerated perceptions of descriptive norms of peers may increase students' willingness to use NPPSSP themselves. Moreover, exaggerated perceptions of injunctive norms of peers may also lead to an increased approval of using NPPSSP in order to match personal attitudes to the perceived peer norms. Social norms interventions that challenge perceptions of descriptive and injunctive peer norms through, for example, mass media campaigns, social marketing strategies or the provision of online personalized feedback (McAlaney, et al., 2011; Perkins, 2003), may be a viable approach to prevent or reduce NPPSSP use among European university students.

There are certain limitations to the present study. The analyses are based on self-reported data collected via a confidential online survey. This is a commonly used survey technique in substance use research among university students to minimize the risk of socially desirable response behavior (Kypri, Gallagher, & Cashell-Smith, 2004). However, in general, an under- or overestimation of NPPSSP use and approval of use due to social expectation bias cannot be ruled out. Moreover, possible misunderstandings of the survey questions by survey participants, i.e., also considering the use of drugs which are available without a prescription, may have led to an overestimation of NPPSSP use and approval of use. However, since only registered local trade names of prescription sedatives and sleeping pills were provided as examples in the survey questionnaire, and given that use and approval rates of NPPSSP are in line with those for other illicit substances asked for in the SNIPE study (Helmer, et al., 2014), the risk of having misunderstood the survey questions can be considered low. On the other hand, the survey questions regarding NPPSSP may have led to an underestimation of use and approval rates since only a selection of registered local trade names of prescription sedatives and sleeping pills (e.g., diazepam, alprazolam, flunitrazepam, midazolam, stilnoct) were included. Furthermore, it is to be noted that individual email addresses were collected for the intervention provided within the study and students may have perceived that they can be identified. In addition, the number of participating students differed between countries, ranging from 107 individuals in the UK to 1,938 in the Slovak Republic. Therefore, selection bias may have differentially affected the sample composition in different countries. Finally, since the analyses are based on cross-sectional survey data, no causal relationships between perceived descriptive and injunctive norms and personal behavior and attitudes towards behavior can be deduced.

5. Conclusions

This study suggests that European university students perceive the use of NPPSSP among their peers to be higher than their personal use and peer attitudes towards the use to be identical or more positive than their personal attitudes. Furthermore, both perceived descriptive and injunctive norms of peers were shown to be associated with students' personal use and attitudes towards the use of NPPSSP, respectively. Social norms interventions may be useful to change exaggerated perceptions regarding the use and attitudes towards NPPSSP use and may prevent or reduce NPPSSP use among European university students.

Declarations of interest: none

Author Disclosures

Role of Funding Sources

The Social Norms Intervention for the Prevention of Polydrug use (SNIPE) project was funded by the European Commission (LS/2009-2010/DPIP/AG). The funder had no involvement in study design, in the collection, analysis, and interpretation of data, in the writing of the manuscript, as well as in the decision to submit the article for publication.

Contributors

GL, SMH, and HZ planned the statistical analyses. GL carried out the analyses, and wrote the manuscript. SMH participated in performing the analyses, as well as in writing and revising the manuscript. CRP, RM, BMB, JM, RCD, GVH, CS, YA, OK, OO, IA, FG contributed to the design of the study, guided the data collection, and revised the manuscript. All authors read and approved the final manuscript.

Conflict of Interest

All authors declare that they have no conflicts of interest.

Acknowledgements

We are grateful to the European Commission for funding the Social Norms Intervention for the Prevention of Polydrug use (SNIPE) project (LS/2009-2010/DPIP/AG).

References

- Arbour-Nicitopoulos, K. P., Kwan, M. Y., Lowe, D., Taman, S., & Faulkner, G. E. (2010). Social norms of alcohol, smoking, and marijuana use within a Canadian university setting. *Journal of American College Health, 59*, 191-196.
- Berkowitz, A. D. (2005). An overview of the social norms approach. In L. C. Ledermann & L. P. Steward (Eds.), *Changing the culture of college drinking: a socially situated prevention campaign* (pp. 193-214). Cresskill, NJ: Hampton Press.
- Bertholet, N., Faouzi, M., Studer, J., Daeppen, J. B., & Gmel, G. (2013). Perception of tobacco, cannabis, and alcohol use of others is associated with one's own use. *Addiction Science & Clinical Practice, 8*, 15.
- Bodenlos, J. S., Malordy, A., Noonan, M., Mayrsohn, A., & Mistler, B. (2014). Prescription drug attitudes questionnaire: development and validation. *Psychology, 05*, 1687-1693.
- Borsari, B., & Carey, K. B. (2001). Peer influences on college drinking: A review of the research. *Journal of Substance Abuse, 13*, 391-424.
- Compton, W. M., & Volkow, N. D. (2006). Abuse of prescription drugs and the risk of addiction. *Drug and Alcohol Dependence, 83 Suppl 1*, S4-7.
- Dempsey, R. C., McAlaney, J., Helmer, S. M., Pischke, C. R., Akvardar, Y., Bewick, B. M., Fawcner, H. J., Guillen-Grima, F., Stock, C., Vriesacker, B., Van Hal, G., Salonna, F., Kalina, O., Orosova, O., & Mikolajczyk, R. T. (2016). Normative perceptions of cannabis use among European university students: associations of perceived peer use and peer attitudes with personal use and attitudes. *Journal of Studies on Alcohol and Drugs, 77*, 740-748.
- Helmer, S. M., Mikolajczyk, R. T., McAlaney, J., Vriesacker, B., Van Hal, G., Akvardar, Y., Guillen-Grima, F., Salonna, F., Stock, C., Dempsey, R. C., Bewick, B. M., & Zeeb, H. (2014). Illicit substance use among university students from seven European countries: a comparison of personal and perceived peer use and attitudes towards illicit substance use. *Preventive Medicine, 67*, 204-209.
- Helmer, S. M., Pischke, C. R., Van Hal, G., Vriesacker, B., Dempsey, R. C., Akvardar, Y., Guillen-Grima, F., Salonna, F., Stock, C., & Zeeb, H. (2016). Personal and perceived peer use and attitudes towards the use of nonmedical prescription stimulants to improve academic performance among university students in seven European countries. *Drug and Alcohol Dependence, 168*, 128-134.
- Hildt, E., Franke, A. G., & Lieb, K. (2011). Pharmakologisches Neuroenhancement. Informationsquellen und Akzeptanz unter Studierenden. *Nervenheilkunde, 10*, 833-837.
- Hildt, E., Lieb, K., & Franke, A. G. (2014). Life context of pharmacological academic performance enhancement among university students - a qualitative approach. *BMC Medical Ethics, 15*, 23.

- Jensen, C., Forlini, C., Partridge, B., & Hall, W. (2016). Australian university students' coping strategies and use of pharmaceutical stimulants as cognitive enhancers. *Frontiers in Psychology, 7*, 277.
- Kilmer, J. R., Geisner, I. M., Gasser, M. L., & Lindgren, K. P. (2015). Normative perceptions of non-medical stimulant use: associations with actual use and hazardous drinking. *Addictive Behaviors, 42*, 51-56.
- Kilmer, J. R., Walker, D. D., Lee, C. M., Palmer, R. S., Mallett, K. A., Fabiano, P., & Larimer, M. E. (2006). Misperceptions of college student marijuana use: implications for prevention. *Journal of Studies on Alcohol and Drugs, 67*, 277-281.
- Kypri, K., Gallagher, S. J., & Cashell-Smith, M. L. (2004). An internet-based survey method for college student drinking research. *Drug and Alcohol Dependence, 76*, 45-53.
- Maier, L. J., Liechti, M. E., Herzig, F., & Schaub, M. P. (2013). To dope or not to dope: neuroenhancement with prescription drugs and drugs of abuse among Swiss university students. *PLoS One, 8*, e77967.
- Maier, L. J., & Schaub, M. P. (2015). The use of prescription drugs and drugs of abuse for neuroenhancement in Europe. *European Psychologist, 20*, 155-166.
- Martens, M. P., Page, J. C., Mowry, E. S., Damann, K. M., Taylor, K. K., & Cimini, M. D. (2006). Differences between actual and perceived student norms: an examination of alcohol use, drug use, and sexual behavior. *Journal of American College Health, 54*, 295-300.
- Martins, S. S., & Ghandour, L. A. (2017). Nonmedical use of prescription drugs in adolescents and young adults: not just a Western phenomenon. *World Psychiatry, 16*, 102-104.
- McAlaney, J., Bewick, B. M., & Hughes, C. (2011). The international development of the 'Social Norms' approach to drug education and prevention. *Drugs: Education Prevention and Policy, 18*, 81-89.
- McAlaney, J., Helmer, S. M., Stock, C., Vriesacker, B., Van Hal, G., Dempsey, R. C., Akvardar, Y., Salonna, F., Kalina, O., Guillen-Grima, F., Bewick, B. M., & Mikolajczyk, R. T. (2015). Personal and perceived peer use of and attitudes toward alcohol among university and college students in seven EU countries: project SNIPE. *Journal of Studies on Alcohol and Drugs, 76*, 430-438.
- McCabe, S. E. (2008). Misperceptions of non-medical prescription drug use: a web survey of college students. *Addictive Behaviors, 33*, 713-724.
- Partridge, B. J., Bell, S. K., Lucke, J. C., Yeates, S., & Hall, W. D. (2011). Smart drugs "as common as coffee": media hype about neuroenhancement. *PLoS One, 6*, e28416.
- Perkins, H. W. (2003). The emergence and evolution of the social norms approach to substance abuse prevention. In H. W. Perkins (Ed.), *The social norms approach to preventing school and*

college age substance abuse: a handbook for educators, counselors, and clinicians (pp. 3-17). San Francisco: Jossey-Bass.

Perkins, H. W. (2014). Misperception is reality: the “reign of error” about peer risk behaviour norms among youth and young adults. In M. Xenitidou & B. Edmonds (Eds.), *The complexity of social norms* (pp. 11-36). Switzerland: Springer International Publishing.

Perkins, H. W., Meilman, P. W., Leichliter, J. S., Cashin, J. R., & Presley, C. A. (1999). Misperceptions of the norms for the frequency of alcohol and other drug use on college campuses. *Journal of American College Health, 47*, 253-258.

Pischke, C. R., Helmer, S. M., McAlaney, J., Bewick, B. M., Vriesacker, B., Van Hal, G., Mikolajczyk, R. T., Akvardar, Y., Guillen-Grima, F., Salonna, F., Orosova, O., Dohrmann, S., Dempsey, R. C., & Zeeb, H. (2015). Normative misperceptions of tobacco use among university students in seven European countries: baseline findings of the 'Social Norms Intervention for the prevention of Polydrug use' study. *Addictive Behaviors, 51*, 158-164.

Pischke, C. R., Zeeb, H., Van Hal, G., Vriesacker, B., McAlaney, J., Bewick, B. M., Akvardar, Y., Guillen-Grima, F., Orosova, O., Salonna, F., Kalina, O., Stock, C., Helmer, S. M., & Mikolajczyk, R. T. (2012). A feasibility trial to examine the social norms approach for the prevention and reduction of licit and illicit drug use in European University and college students. *BMC Public Health, 12*.

Sanders, A., Stogner, J., Seibert, J., & Miller, B. L. (2014). Misperceptions of peer pill-popping: the prevalence, correlates, and effects of inaccurate assumptions about peer pharmaceutical misuse. *Substance Use & Misuse, 49*, 813-823.

Silvestri, M. M., & Correia, C. J. (2016). Normative influences on the nonmedical use of prescription stimulants among college students. *Psychology of Addictive Behaviors, 30*, 516-521.

United Nations Office on Drugs and Crime (2011). The non-medical use of prescription drugs: Policy direction issues. Vienna: United Nations. <http://www.unodc.org/documents/drug-prevention-and-treatment/nonmedical-use-prescription-drugs.pdf> Accessed 18 January 2018.

Highlights

- 9.1% of the participating European university students reported having used NPPSSP.
- Students show discrepancies between perceived peer and own NPPSSP use/attitudes.
- Perceived peer NPPSSP use was associated with higher odds for students' own use.
- Perceived approval among peers was associated with higher odds for own approval.
- Correcting misperceived peer norms may prevent or reduce NPPSSP use among students.

ACCEPTED MANUSCRIPT