

**This item is the archived peer-reviewed author-version of:**

Do spatiotemporal parameters and gait variability differ across the lifespan of healthy adults? A systematic review

**Reference:**

Herssens Nolan, Verbecque Evi, Hallems Ann, Vereeck Luc, Van Rompaey Vincent, Saeys Wim.- Do spatiotemporal parameters and gait variability differ across the lifespan of healthy adults? A systematic review  
Gait and posture - ISSN 0966-6362 - Clare, Elsevier ireland ltd, 64(2018), p. 181-190  
Full text (Publisher's DOI): <https://doi.org/10.1016/J.GAITPOST.2018.06.012>  
To cite this reference: <https://hdl.handle.net/10067/1522100151162165141>

Do spatiotemporal parameters and gait variability differ across the lifespan of healthy adults? A systematic review.

Nolan Herssens, Evi Verbecque, Ann Halleman, Luc Vereeck, Vincent Van Rompaey & Wim Saeys.

Corresponding author: Nolan Herssens, Movant, Universiteitsplein 1, Campus Drie Eiken, D.S.022, 2610 Wilrijk, Belgium. [nolan.herssens@uantwerpen.be](mailto:nolan.herssens@uantwerpen.be)

Nolan Herssens, Department of Rehabilitation Sciences and Physiotherapy/ Movant, Faculty of Medicine and Health Science, University of Antwerp, Belgium; Multidisciplinary Motor Centre Antwerp (M<sup>2</sup>OCEAN), University of Antwerp, Belgium.

Evi Verbecque, PhD, Department of Rehabilitation Sciences and Physiotherapy/ Movant, Faculty of Medicine and Health Science, University of Antwerp, Belgium.

Ann Halleman, PhD, Department of Rehabilitation Sciences and Physiotherapy/ Movant, Faculty of Medicine and Health Science, University of Antwerp, Belgium; Multidisciplinary Motor Centre Antwerp (M<sup>2</sup>OCEAN), University of Antwerp, Belgium.

Luc Vereeck, PhD, Department of Rehabilitation Sciences and Physiotherapy/ Movant, Faculty of Medicine and Health Science, University of Antwerp, Belgium; Multidisciplinary Motor Centre Antwerp (M<sup>2</sup>OCEAN), University of Antwerp, Belgium.

Vincent Van Rompaey, MD, PhD, Department of Otorhinolaryngology and Head & Neck Surgery, Antwerp University Hospital, Edegem, Belgium; Faculty of Medicine and Health Sciences, University of Antwerp, Belgium.

Wim Saeys, PhD, Department of Rehabilitation Sciences and Physiotherapy/ Movant, Faculty of Medicine and Health Science, University of Antwerp, Belgium; RevArte Rehabilitation Hospital, Edegem, Antwerp.

# Highlights

- Most mean spatiotemporal parameters seem to be affected by age
- Mean step width and variability measures remain stable over time
- Most of the reported data are based on an elderly population
- Data concerning middle-aged adults are lacking

## Abstract

**BACKGROUND:** Aging is often associated with changes in the musculoskeletal system, peripheral and central nervous system. These age-related changes often result in mobility problems influencing gait performance. Compensatory strategies are used as a way to adapt to these physiological changes.

**RESEARCH QUESTION:** The aim of this review is to investigate the differences in spatiotemporal and gait variability measures throughout the healthy adult life.

**METHODS:** This systematic review was conducted according to the PRISMA guidelines and registered in the PROSPERO database (no. CRD42017057720). Databases MEDLINE (Pubmed), Web of Science (Web of Knowledge), Cochrane Library and ScienceDirect were systematically searched until March 2018.

**RESULTS:** Eighteen of the 3195 original studies met the eligibility criteria and were included in this review. The majority of studies reported spatiotemporal and gait variability measures in adults above the age of 65, followed by the young adult population, information of middle-aged adults is lacking. Spatiotemporal parameters and gait variability measures were extracted from 2112 healthy adults

between 18 and 98 years old and, in general, tend to deteriorate with increasing age. Variability measures were only reported in an elderly population and show great variety between studies.

**SIGNIFICANCE:** The findings of this review suggest that most spatiotemporal parameters significantly differ across different age groups. Elderly populations show a reduction of preferred walking speed, cadence, step and stride length, all related to a more cautious gait, while gait variability measures remain stable over time. A preliminary framework of normative reference data is provided, enabling insights into the influence of aging on spatiotemporal parameters, however spatiotemporal parameters of middle-aged adults should be investigated more thoroughly.

## **Keywords**

Gait – Adult – Aging – Spatiotemporal Parameters – Gait Variability

## **Introduction**

Gait is often used as a way to quantify physical function, quality of life and health status [1] since walking is one of the most frequently performed physical activities in daily life [2]. Changes in gait performance have been demonstrated to correlate with risk of falling [3], cognitive impairment (e.g. dementia) [4] or even risk of early mortality [5]. The ability to walk safely and efficiently is an important predisposition in maintaining independence with older age [6]. However, aging is associated with changes in the muscular, skeletal [7] and central/peripheral nervous systems [8, 9]. Impairments such as loss of muscle strength [10] and proprioceptive feedback [11], but also deterioration of specific brain sites, for example in motor cortical regions [12] or the basal ganglia [13], related to normal aging can result in mobility problems and an increased risk of falling [14, 15].

All of these age-related changes can influence gait performance. Compensatory strategies are observed to increase stability and prevent falls [16] as a way to adapt to the physiological changes that occur with aging. Elderly populations tend to develop a more cautious gait, which is commonly characterized by a reduced walking speed, reduced stride length and an increased step width [17]. Assessment of spatiotemporal gait parameters thus provides objective data of the global performance [18, 19]. More recently, the use of gait variability measures has increased since they may be more sensitive in quantifying age-related changes in the motor control of gait [8]. Gait variability measures seem to show an increase in variability with advancing age [20].

Most frequently, spatiotemporal parameters of healthy adults are used as control values in order to determine if a certain patient population diverts from the norm, for instance when investigating the influence of neurological [21, 22], cardiovascular [23, 24] or musculoskeletal impairments on gait [25, 26]. On the other hand, spatiotemporal parameters of healthy adults have been used in order to investigate the influence of age on the way of walking. Most research focussing on the effect of aging on gait has been comparing younger with older adults [7] or different elderly populations with each other [1]. Therefore, an overview of the available literature concerning spatiotemporal parameters of adults of all ages can provide more transparency on the influence of aging on gait. Additionally, understanding the effect of aging on gait can provide important knowledge for clinicians on which gait characteristics are important for a safe, independent and efficient gait in elderly populations, and which could eventually provide guidelines for rehabilitation.

Furthermore, a recent review only discussed changes in a limited amount of spatiotemporal parameters in combination with changes in kinetics, kinematics and energy consumption with increasing age [27]. Other reviews investigated the changes in gait variability related to fear of falling [28] or analysed the differences in spatiotemporal parameters of elderly fallers and non-fallers [29], none discussing the changes in gait parameters with increasing age in healthy adults.

Therefore, the purpose of this systematic review is to investigate the differences in spatiotemporal and gait variability measures throughout healthy adult life, measured through an instrumented walkway during walking at a self-selected walking speed.

## Methods

### Protocol and registration

This systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Review and Meta-Analysis Statement (PRISMA). The protocol is available at PROSPERO (registration nr. CRD42017057720) and can be consulted online ([www.crd.york.ac.uk/prospero/](http://www.crd.york.ac.uk/prospero/)).

### Systematic Literature Search

A systematic literature search, based on the Population Intervention Comparison Outcome (PICO) method, was conducted in March 2018 in MEDLINE (Pubmed), Web of Science (Web of Knowledge), Cochrane Library for clinical trials and ScienceDirect using combinations of the following keywords: 'Adult', 'Gait' and 'Spatiotemporal Parameters'.

No filters were applied. Specifics on the search queries for the different databases are available in Appendix 1.

### Eligibility Criteria & Study Selection

To select relevant literature, the following selection criteria were applied:

1. Participants were healthy adults aged 18 and older consisting of a combination of men and women, for which evidence had to be provided : at least a statement on known neurological, cardiovascular and musculoskeletal health, BMI<30, absence of any other impairments that can influence gait (e.g. visual or vestibular impairment). The samples had to be representative for the average population of healthy people. Therefore, studies investigating e.g. athletes, high-level sports participation, soldiers, one of the sexes exclusively were

excluded from the sample. A minimum sample size of 30 participants was considered representative in accordance with the COnsensus-based Standards for the selection of health Measurement Instruments for identifying differences between relevant groups (hypothesis testing) [30].

2. Data must be collected during an instrumented overground gait analysis (e.g. GAITRite, force platforms, 3D gait analysis etc) while walking at a self-selected walking speed. Data collected using accelerometry were excluded as spatial variables and walking speed can be significantly underestimated [31]. Citations were excluded when data were collected during functional walking measures (e.g. Timed Up and Go, obstacle negotiation, climbing stairs, six-minute walk test); treadmill walking or during running, turning, stepping tasks (lateral, forward, backward, etc) or when slips/perturbations were induced.
3. Numeric values of mean spatiotemporal parameters (STP) or gait variability measures reported as primary outcome. When STP were reported for left and right separately, the results for the right side were extracted.
4. Original research, including full length articles and brief reports in which the applied methodology was reported transparently, written in English, French, Dutch or German. Reviews of any kind, meta-analyses, case studies/-series, conference proceedings, abstract only, books/book chapters, letter to the editor, study protocols, pilot studies, editorials or opinion pieces were excluded.

The selection process was conducted in two phases. In phase 1, the selection criteria were applied on the citations' title and abstract. If a study met these criteria or its eligibility could not be determined from the title and abstract, it was selected on full-text screening (phase 2) using the same eligibility criteria. After phase two, references of all included studies were screened, and included if eligible, to ensure that no relevant articles were missed. The selection process was performed by two independent reviewers (NH, EV). The selection process is presented in Figure 1.

## Risk of Bias

Risk of bias was not assessed, as for now no assessment tool for identifying risk of bias regarding normative data is available. Bias was addressed by applying rigorous selection criteria, taking representativeness of the study sample and sample size (selection bias) into account.

## Data Extraction

Data were extracted by two reviewers (NH, EV) and summarized in a methodology and evidence table. The methodology table summarizes information on study design, type of instrumented walkway, walkway length, footwear, study sample characteristics such as number of participants, age (min, max, mean and standard deviation (SD)), sex distribution (number of women), body mass index (BMI, mean and SD), body length (mean and SD) and origin.

The primary outcome measures were means and standard deviations of either spatiotemporal parameters or gait variability measures, which were plotted according to age. When results for comparable groups were available, e.g. men and women or separate groups receiving a different type of intervention but with controlled sample characteristics, a weighted mean was calculated based on the raw (baseline) data.

Outcome measures reported as step width, stride width and base of support were combined into one measure: step width, as they all reported a similar distance between the two feet in medio-lateral direction. If necessary, units were converted to from cm/s to m/s for gait speed; cm to m for spatial parameters and ms to s for temporal parameters. Applied definitions for the extracted STP are presented in Appendix 2.

## Results

### Search Results

In total, 3195 original studies were identified from the electronic databases MEDLINE (Pubmed), Web of Science (Web of Knowledge), Cochrane Library and ScienceDirect after deduplication. After



screening on title and abstract, 252 were deemed potentially relevant and were taken to screening phase two. After a more detailed screening of the full text, 18 of the 252 studies met the eligibility criteria and were included in this review. No additional studies were included after hand searching the references of the included articles. The selection process is presented in Figure 1.

INSERT FIGURE 1 HERE

### Study Characteristics

Data were extracted from 2112 healthy adults (1131 women, 53.6%) between 18 [32] and 98 [33] years old. Seven included studies reported the STP or variability measures of a young adult population (aged 19 – 40) [32, 34-39], ten studies of an elderly population (aged 65 and up) [1, 33, 40-47] and one study compared a young adult population with an elderly population [48]. Specifics on the sample characteristics are provided in Table 1.

Mean and variability measures were collected through the GAITRite system (n=8; [1, 32-34, 40-43]), OptoGait system (n=1; [35]), Gait Mat system (n=1; [44]), Vicon motion analysis system (n=5; [36-39, 46]), a Selspot II optoelectric system (n=1; [48]), a BTS 3D GA system (n=1; [47]) and a 3D – Motion Analysis Corporation system (n=1; [45]). Gait parameters were all measured during over ground walking on a walkway varying from 4.60 to 30 meters in length, at a self-selected walking speed. Only four studies reported if the subjects walked barefoot [35, 42, 44] or wearing shoes [43] (Table 1).

INSERT TABLE 1 HERE.

### Differences in Spatiotemporal Parameters as a Function of Age

Numeric values for mean STP and gait variability are presented in Table 2 and 3 respectively.

Graphical presentation of the STP as a function of age are presented in Figure 2 (mean values) and 3 (gait variability).

INSERT TABLE 2 & 3 HERE

### *Spatial parameters*

A total of 12 references included in this review reported spatial parameters. *Step length* was reported in eight of the included articles and ranged from 0.711m  $\pm$  0.011m in 60-year-olds [43] to 0.495  $\pm$  0.079m in 90-year-olds [33]. *Stride length* decreased from 1.406m in 30-year-olds [34] to 0.994m  $\pm$  0.158m in 90-year-olds [33]. *Step width* was reported to be lowest in 30-year-olds at 0.070m  $\pm$  0.047m [37] and highest in 70-year-olds at 0.151m  $\pm$  0.021m [45]. However, a great variability in reported step width is noted in the elderly population. In general step- and stride length tend to decrease with increasing age, whereas step width tends to remain stable over time (Figure 2).

Significant differences were found for step length and stride length with increasing age in elderly populations aged 65 to 85 and older ( $p < 0.001$ ) [40], when comparing 70-74 and 75-79 year-olds to elderly aged 80 and older ( $p < 0.001$ ) [1] or in elderly aged 70-98 ( $p \leq 0.006$ ) [33]. However, no significant differences in step width were found regardless of the age groups investigated [1, 40].

#### *Temporal parameters*

Temporal parameters were reported in ten of the included references. *Step time* and *stride time* both show an increase with increasing age as step and stride time were lowest for 30-year-olds at 0.50s  $\pm$  0.02s and 0.99s  $\pm$  0.03s [36] and highest in 80-year-olds at 0.57s  $\pm$  0.06s [1] for step time and 1.18s  $\pm$  0.14s [40] for stride time. *Stance time* ranged from 0.66s  $\pm$  0.07s in 20-year-olds [48] to 0.78s  $\pm$  0.11s in 80-year-olds [40], the same is noted for *single support time* as 70-year-olds are reported to have a single support time of 0.42s  $\pm$  0.04s [40], while 20-year-olds were reported to have a single support time of 0.33s  $\pm$  0.03s [39]. *Swing time* also increased from 0.36s  $\pm$  0.02s in 20-year-olds [35] to 0.42s  $\pm$  0.04s in 70-year-olds [40]. For *double support time*, 30-year-olds showed a double-limb support time of 0.19s  $\pm$  0.02s [36] which increased to 0.38s  $\pm$  0.10s in 80-year-olds [40]. In general, temporal parameters show an increase when comparing young adults with elderly, while within the elderly population temporal parameters stay fairly consistent.

However, some results are not in line with results reported by Beauchet et al. [40] as they reported a significant decrease of swing ( $p=0.049$ ) and single-limb support times ( $p=0.021$ ) between elderly groups ranging between 65 and 85 years of age. Stride time, stance time and double-limb support time increased with age ( $p<0.001$ ) and were in line with the other results found in this review (Figure 2). On the other hand, Hollman et al. [1] did not find any significant differences in temporal parameters between age groups in an elderly cohort ranging from 70 years to 85+ years of age, except for double-limb support time between 70-74 and 85-89 year-olds ( $p<0.05$ ) and 75-79 and 85-89 year-olds ( $p<0.01$ ).

INSERT FIGURE 2 HERE

#### *Combined parameters*

Combined parameters walking speed and cadence were reported in 17 of the studies included in this review. The *walking speed* of 30-year-olds was noted to be the highest at  $1.50\text{m/s} \pm 0.17\text{m/s}$  [37] and decreased with time to  $0.87\text{m/s} \pm 0.17\text{m/s}$  in 90-year-olds [33]. The same shift can be noted for *cadence* as in 90-year-olds cadence was reported to be  $105.58 \pm 7.32$  steps/min [37], while 30-year-olds displayed the highest cadence at  $122.00 \pm 4.27$  steps/min [33]. McGibbon et al. [48] compared 20-year-olds with 70-year-olds and found a significant decrease in gait speed between these two groups ( $p<0.05$ ). Hollman et al. [1] and Chui et al. [33] both investigated differences within an elderly population and both found a significant decrease in mean walking speed between elderly age-groups ( $p<0.001$ ). However, cadence was not significantly decreased ( $p=0.051$ ) in the investigated age-groups of Hollman et al. [1] but did significantly decrease ( $p<0.001$ ) in the cohort of Chui et al. [33].

#### *Gait cycle phases*

Only three of the included references reported gait cycle percentages. In an elderly population, an increase in *stance* and *double support phase* is noted:  $62.91\% \pm 1.22\%$  and  $25.81\% \pm 2.41\%$  in 70-year-olds, to  $65.31\% \pm 21.88\%$  to  $30.77\% \pm 1.38\%$  in 90-year-olds respectively [33]. The *swing phase* tends to decrease when comparing 70-year-olds to 90-year-olds ( $37.10\% \pm 1.22\%$  to  $34.71\% \pm$

1.88%), as did the *single-limb support*. However, single-limb support percentages varied greatly, as Chui et al. [33] reported the mean single-limb support of both legs combined, ranging between  $74.21\% \pm 2.47$  to  $69.42\% \pm 3.72$  in 70 to 90-year-olds ( $p < 0.05$ ), while Hollman et al. [1] reported a single-limb support for one leg, ranging between  $36.60\% \pm 2.11\%$  for 70-year-olds to  $35.52\% \pm 2.63\%$  for 80-year-olds but these differences were not significant ( $p > 0.05$ ).

#### *Gait variability measures*

Gait variability measures were expressed as coefficients of variation (CoV;  $(SD/mean) \cdot 100$ ) and were almost exclusively reported in an elderly population ranging from 70 years to 89 years of age. *Step length* and *double-limb support time CoV* show a decrease with increasing age ( $6.31\% \pm 9.40\%$  to  $6.08\% \pm 2.70\%$  and  $9.78\%$  to  $6.00\% \pm 2.10\%$  respectively [1, 40]), while *stride time CoV* ( $2.23\%$  to  $5.02\% \pm 4.82\%$ ) and *swing time CoV* ( $4.38\%$  to  $8.18\% \pm 10.21\%$ ) show an increase with increasing age [1, 34]. *Step width* and *stance time CoV* remain fairly stable in an elderly population. However, values between studies seem to differ widely (Figure 3). Hollman et al. [1] did not find any significant age-effects, however Beauchet et al. [40] did find significant age effects except for double-limb support time CoV ( $p = 0.186$ ).

INSERT FIGURE 3 HERE

## **Discussion**

The purpose of this systematic review was to describe the differences in spatiotemporal and gait variability measures throughout healthy adult life, measured through an instrumented walkway, during walking at self-selected walking speed. In total, 23 unique spatiotemporal and gait variability measures were identified in the currently included literature, gathered through an instrumented walkway or a 3D-motion analysis system. Overall findings of this literature review are: 1) most mean spatiotemporal parameters of gait seem to be affected by age, 2) mean step width and variability measures seem to remain stable over time, 3) most of the reported data are based on an elderly

population above the age of 65, followed by the young adult population, while data from the middle-aged adult population is lacking and 4) variability measures are only reported in an elderly population and show a great variety between studies.

The data presented in this study are based on healthy adults without any musculoskeletal, cardiovascular or neurological impairments. These strict selection criteria were defined as these comorbidities may impose important changes to the way of walking [21, 24, 49]. Because of these rigorous criteria, the presented data represent the effect of aging without the influence of multimorbidities that often co-occur with aging [50, 51], allowing a solid basis towards identifying gait deviations. The small amount of included papers is therefore not surprising. Although a large set of studies is available on “healthy” adults that include people that are free of any neurological, musculoskeletal or orthopedic impairments, evidence of cardiovascular health is often lacking. However, cardiovascular health also plays an important role in functioning, which is shown by altered gait patterns, e.g. in patients with intermittent claudication a slower walking speed, shorter step length and lower cadence is noted [23]. The results in this review therefore additionally provide a preliminary framework of normative reference data that enables insights into the influence of aging on STP, however data of middle-aged adults is lacking.

#### Age-related differences in mean STP

In general, the results of this systematic review indicate a reduction of gait performance with increasing age as spatial and spatio-temporal parameters tend to decrease while temporal parameters tend to increase. These features seem to suggest that as people grow older, they develop a cautious gait pattern, characterized by decreased walking speed, cadence, step length and stride length and increased step time, stride time and phases.

Walking speed, the most reported spatiotemporal parameter, declines with increasing age. This may be the result of the short step and stride lengths noticed in an elderly population, which in its turn may be the result of weakened hip extensors and ankle plantar flexors reducing the capacity to

propel the body forwards [52]. A decreased walking speed may also be the result of balance impairments [53] which results in a more cautious gait [29, 54]. This cautious gait can also be noted in the differences of the temporal parameters across age groups with increasing age, which may be the result of lower limb muscle weakness [55, 56] or might be a compensatory strategy used to increase weight-bearing stability during gait [2]. However, when looking at the differences in step width and gait cycle percentages, these remain fairly stable throughout the different age-groups as only small differences are noted with increasing age.

#### Age-related differences in gait variability

Measures related to gait variability in the included literature were almost exclusively reported in an elderly cohort above the age of 70 and, in general, remained fairly stable. Age-related differences in spatial and temporal variability measures may be elicited by a loss of lower extremity strength and range of motion, increased muscle activation variation and changes in balance [9, 57]. These differences in balance can be the result due to a decline in central motor control, worsening of the automatic stepping mechanism [8] or an insufficient postural stability in general, related to older age [3]. As stated by Dingwell et al. [58], aged healthy adults implement the same underlying stride-to-stride control strategies as young adults do. However, due to physiological changes neuromotor noise is increased, which in its turn results in a greater variability even though healthy elderly are just as successful in reducing the effect of input noise. On the other hand, as stated by Hausdorff et al. [20], gait variability measures may be able to reveal increased variability or instability even when the subsystem that contributes to gait variability only shows subtle changes.

#### Limitations of the study

Some limitations are to be considered. First, no risk of bias assessment was performed on the included references as only data from healthy control groups were extracted and implemented in this review for which an assessment tool is still lacking. Instead, selection criteria were defined in such a way that bias could be accounted for, without using a dedicated assessment tool. Additionally,

strict selection towards methods on data-collection of STP were addressed as well, as the purpose of this review was not report the effect if interventions or specific walking tasks on STP.

Secondly, a lack of information regarding cardiovascular, musculoskeletal or neurological health of the healthy adult population resulted in the exclusion of several studies reporting spatiotemporal parameters or gait variability measures.

Implications for future research and clinical relevance

Clearly, data on middle-aged people are lacking. This suggests that future research is necessary to fill the existing gap regarding normative reference values for STP in gait in the middle-aged population. Additionally, a broad range in spatiotemporal parameters is available with a high covariance among these measures which may indicate redundancy [59]. One method to narrow the range of different gait characteristics and increase transparency is to cluster the different characteristics in components of gait through a principal component analysis, as suggested by Lord et al. [59]. These domains of gait could provide a comprehensive but flexible approach to assessment as the selection of gait characteristics will depend on the gait pathology. However, in general these (mean) spatiotemporal parameters are a quick and easy way to quantify gait performance.

On the other hand, consensus regarding the methods used to analyse gait variability is lacking [60]. Articles included in this review only reported variability as the coefficient of variation, while variability also can be reported as standard deviation (SD). Lord et al. [60] suggest the use of SD to report gait variability as it provides more clarity as opposed to the coefficient of variation which is a composed parameter which can be influenced by either the mean or the standard deviation. Clearly, the exact meaning of variability measures remains to be determined as well as how variability should be expressed.

## Conclusion

The findings from this systematic review suggest a deterioration in walking in elderly populations, shown by the differences in spatiotemporal and gait variability measures. However, additional

research is necessary in middle-aged adults from the age of 30 up to the age of 59 to be able to determine when these differences start to appear and to be able to fully understand the effects of aging on gait.

### **Conflict of Interest**

None.

### **Acknowledgements**

This work was supported by the University of Antwerp Research Council [grant number ID33586], the University of Antwerp and the Antwerp University Hospital.

### **References**

1. Hollman JH, McDade EM, Petersen RC. Normative spatiotemporal gait parameters in older adults. *Gait Posture*. 2011;34(1):111-8.
2. Afiah IN, Nakashima H, Loh PY, Muraki S. An exploratory investigation of changes in gait parameters with age in elderly Japanese women. *Springerplus*. 2016;5(1):1069.
3. Maki BE. Gait changes in older adults: predictors of falls or indicators of fear. *J Am Geriatr Soc*. 1997;45(3):313-20.
4. Verghese J, Wang C, Lipton RB, Holtzer R, Xue X. Quantitative gait dysfunction and risk of cognitive decline and dementia. *J Neurol Neurosurg Psychiatry*. 2007;78(9):929-35.
5. Studenski S, Perera S, Patel K, Rosano C, Faulkner K, Inzitari M, et al. Gait speed and survival in older adults. *Jama*. 2011;305(1):50-8.



6. Callisaya ML, Blizzard L, Schmidt MD, McGinley JL, Srikanth VK. Ageing and gait variability--a population-based study of older people. *Age Ageing*. 2010;39(2):191-7.
7. McGibbon CA. Toward a better understanding of gait changes with age and disablement: neuromuscular adaptation. *Exerc Sport Sci Rev*. 2003;31(2):102-8.
8. Hausdorff JM. Gait dynamics, fractals and falls: finding meaning in the stride-to-stride fluctuations of human walking. *Hum Mov Sci*. 2007;26(4):555-89.
9. Brach JS, Studenski S, Perera S, VanSwearingen JM, Newman AB. Stance time and step width variability have unique contributing impairments in older persons. *Gait Posture*. 2008;27(3):431-9.
10. Perry MC, Carville SF, Smith IC, Rutherford OM, Newham DJ. Strength, power output and symmetry of leg muscles: effect of age and history of falling. *Eur J Appl Physiol*. 2007;100(5):553-61.
11. Skinner HB, Barrack RL, Cook SD. Age-related decline in proprioception. *Clin Orthop Relat Res*. 1984(184):208-11.
12. Seidler RD, Bernard JA, Burutolu TB, Fling BW, Gordon MT, Gwin JT, et al. Motor control and aging: links to age-related brain structural, functional, and biochemical effects. *Neurosci Biobehav Rev*. 2010;34(5):721-33.
13. Raz N, Rodrigue KM, Kennedy KM, Head D, Gunning-Dixon F, Acker JD. Differential aging of the human striatum: longitudinal evidence. *AJNR Am J Neuroradiol*. 2003;24(9):1849-56.
14. Verghese J, LeValley A, Hall CB, Katz MJ, Ambrose AF, Lipton RB. Epidemiology of gait disorders in community-residing older adults. *J Am Geriatr Soc*. 2006;54(2):255-61.
15. Rubenstein LZ, Josephson KR. The epidemiology of falls and syncope. *Clin Geriatr Med*. 2002;18(2):141-58.
16. Pavol MJ, Owings TM, Foley KT, Grabiner MD. Gait characteristics as risk factors for falling from trips induced in older adults. *J Gerontol A Biol Sci Med Sci*. 1999;54(11):M583-90.
17. Nutt JG. Classification of gait and balance disorders. *Adv Neurol*. 2001;87:135-41.

18. Gouelle A, Mégrot F, Presedo A, Husson I, Yelnik A, Penneçot G-F. The Gait Variability Index: A new way to quantify fluctuation magnitude of spatiotemporal parameters during gait. *Gait & Posture*. 2013;38(3):461-5.
19. Morris R, Hickey A, Del Din S, Godfrey A, Lord S, Rochester L. A model of free-living gait: A factor analysis in Parkinson's disease. *Gait Posture*. 2017;52:68-71.
20. Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil*. 2001;82(8):1050-6.
21. Boudarham J, Roche N, Pradon D, Bonnyaud C, Bensmail D, Zory R. Variations in kinematics during clinical gait analysis in stroke patients. *PLoS One*. 2013;8(6):e66421.
22. Roiz Rde M, Cacho EW, Pazinato MM, Reis JG, Cliquet A, Jr., Barasnevicius-Quagliato EM. Gait analysis comparing Parkinson's disease with healthy elderly subjects. *Arq Neuropsiquiatr*. 2010;68(1):81-6.
23. Gommans LNM, Smid AT, Scheltinga MRM, Cancrinus E, Brooijmans FAM, Meijer K, et al. Prolonged stance phase during walking in intermittent claudication. *J Vasc Surg*. 2017;66(2):515-22.
24. Gardner AW, Forrester L, Smith GV. Altered gait profile in subjects with peripheral arterial disease. *Vasc Med*. 2001;6(1):31-4.
25. Kiss RM, Illyes A. Comparison of gait parameters in patients following total hip arthroplasty with a direct-lateral or antero-lateral surgical approach. *Hum Mov Sci*. 2012;31(5):1302-16.
26. Magyar MO, Knoll Z, Kiss RM. The influence of medial meniscus injury and meniscectomy on the variability of gait parameters. *Knee Surg Sports Traumatol Arthrosc*. 2012;20(2):290-7.
27. Aboutorabi A, Arazpour M, Bahramizadeh M, Hutchins SW, Fadayevatan R. The effect of aging on gait parameters in able-bodied older subjects: a literature review. *Aging Clin Exp Res*. 2016;28(3):393-405.
28. Ayoubi F, Launay CP, Annweiler C, Beauchet O. Fear of falling and gait variability in older adults: a systematic review and meta-analysis. *J Am Med Dir Assoc*. 2015;16(1):14-9.


29. Mortaza N, Abu Osman NA, Mehdikhani N. Are the spatio-temporal parameters of gait capable of distinguishing a faller from a non-faller elderly? *Eur J Phys Rehabil Med*. 2014;50(6):677-91.
30. Terwee CB, Mokkink LB, Knol DL, Ostelo RWJG, Bouter LM, de Vet HCW. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Quality of Life Research*. 2012;21(4):651-7.
31. Maffioletti NA, Gorelick M, Kramers-de Quervain I, Bizzini M, Munzinger JP, Tomasetti S, et al. Concurrent validity and intrasession reliability of the IDEEA accelerometry system for the quantification of spatiotemporal gait parameters. *Gait Posture*. 2008;27(1):160-3.
32. Kodesh E, Falash F, Sprecher E, Dickstein R. Light touch and medio-lateral postural stability during short distance gait. *Neuroscience Letters*. 2015;584:378-81.
33. Chui KK, Lusardi MM. Spatial and temporal parameters of self-selected and fast walking speeds in healthy community-living adults aged 72-98 years. *J Geriatr Phys Ther*. 2010;33(4):173-83.
34. Dujmovic I, Radovanovic S, Martinovic V, Dackovic J, Maric G, Mesaros S, et al. Gait pattern in patients with different multiple sclerosis phenotypes. *Mult Scler Relat Disord*. 2017;13:13-20.
35. Gomez Bernal A, Becerro-de-Bengoa-Vallejo R, Losa-Iglesias ME. Reliability of the OptoGait portable photoelectric cell system for the quantification of spatial-temporal parameters of gait in young adults. *Gait Posture*. 2016;50:196-200.
36. Meldrum D, Shouldice C, Conroy R, Jones K, Forward M. Test-retest reliability of three dimensional gait analysis: including a novel approach to visualising agreement of gait cycle waveforms with Bland and Altman plots. *Gait Posture*. 2014;39(1):265-71.
37. Sturnieks DL, Besier TF, Hamer PW, Ackland TR, Mills PM, Stachowiak GW, et al. Knee strength and knee adduction moments following arthroscopic partial meniscectomy. *Med Sci Sports Exerc*. 2008;40(6):991-7.
38. Sturnieks DL, Besier TF, Mills PM, Ackland TR, Maguire KF, Stachowiak GW, et al. Knee joint biomechanics following arthroscopic partial meniscectomy. *J Orthop Res*. 2008;26(8):1075-80.


39. Yang F, King GA. Dynamic gait stability of treadmill versus overground walking in young adults. *J Electromyogr Kinesiol.* 2016;31:81-7.
40. Beauchet O, Allali G, Sekhon H, Verghese J, Guilain S, Steinmetz JP, et al. Guidelines for Assessment of Gait and Reference Values for Spatiotemporal Gait Parameters in Older Adults: The Biomathics and Canadian Gait Consortiums Initiative. *Frontiers in Human Neuroscience.* 2017;11.
41. Delbaere K, Sturnieks DL, Crombez G, Lord SR. Concern about falls elicits changes in gait parameters in conditions of postural threat in older people. *J Gerontol A Biol Sci Med Sci.* 2009;64(2):237-42.
42. Elboim-Gabyzon M, Rotchild S. Spatial and temporal gait characteristics of elderly individuals during backward and forward walking with shoes and barefoot. *Gait Posture.* 2017;52:363-6.
43. Jenkins ME, Almeida QJ, Spaulding SJ, van Oostveen RB, Holmes JD, Johnson AM, et al. Plantar cutaneous sensory stimulation improves single-limb support time, and EMG activation patterns among individuals with Parkinson's disease. *Parkinsonism Relat Disord.* 2009;15(9):697-702.
44. Khashan M, Mor A, Beer Y, Rath E, Morgenstern DR, Debi R, et al. Gait metric profile and gender differences in hip osteoarthritis patients. A case-controlled study. *Hip Int.* 2014;24(3):270-6.
45. Lee HJ, Lee S, Chang WH, Seo K, Shim Y, Choi BO, et al. A Wearable Hip Assist Robot Can Improve Gait Function and Cardiopulmonary Metabolic Efficiency in Elderly Adults. *IEEE Trans Neural Syst Rehabil Eng.* 2017;25(9):1549-57.
46. McClelland JA, Webster KE, Feller JA, Menz HB. Knee kinematics during walking at different speeds in people who have undergone total knee replacement. *Knee.* 2011;18(3):151-5.
47. Pistacchi M, Gioulis M, Sanson F, De Giovannini E, Filippi G, Rossetto F, et al. Gait analysis and clinical correlations in early Parkinson's disease. *Funct Neurol.* 2017;32(1):28-34.
48. McGibbon CA, Krebs DE. Discriminating age and disability effects in locomotion: neuromuscular adaptations in musculoskeletal pathology. *J Appl Physiol (1985).* 2004;96(1):149-60.

49. Al-Zahrani KS, Bakheit AM. A study of the gait characteristics of patients with chronic osteoarthritis of the knee. *Disabil Rehabil.* 2002;24(5):275-80.
50. Divo MJ, Martinez CH, Mannino DM. Ageing and the epidemiology of multimorbidity. *The European respiratory journal.* 2014;44(4):1055-68.
51. Salive ME. Multimorbidity in Older Adults. *Epidemiologic Reviews.* 2013;35(1):75-83.
52. Uematsu A, Tsuchiya K, Kadono N, Kobayashi H, Kaetsu T, Hortobágyi T, et al. A Behavioral Mechanism of How Increases in Leg Strength Improve Old Adults' Gait Speed. *PLoS ONE.* 2014;9(10):e110350.
53. Schultz AB. Muscle function and mobility biomechanics in the elderly: an overview of some recent research. *J Gerontol A Biol Sci Med Sci.* 1995;50 Spec No:60-3.
54. Giladi N, Herman T, Reider G, Il, Gurevich T, Hausdorff JM. Clinical characteristics of elderly patients with a cautious gait of unknown origin. *J Neurol.* 2005;252(3):300-6.
55. Simoneau E, Martin A, Van Hoecke J. Effects of joint angle and age on ankle dorsi- and plantar-flexor strength. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology.* 2007;17(3):307-16.
56. Brown M, Sinacore DR, Host HH. The relationship of strength to function in the older adult. *J Gerontol A Biol Sci Med Sci.* 1995;50 Spec No:55-9.
57. Hausdorff JM, Cudkowicz ME, Firtion R, Wei JY, Goldberger AL. Gait variability and basal ganglia disorders: stride-to-stride variations of gait cycle timing in Parkinson's disease and Huntington's disease. *Mov Disord.* 1998;13(3):428-37.
58. Dingwell JB, Salinas MM, Cusumano JP. Increased gait variability may not imply impaired stride-to-stride control of walking in healthy older adults: Winner: 2013 Gait and Clinical Movement Analysis Society Best Paper Award. *Gait Posture.* 2017;55:131-7.
59. Lord S, Galna B, Verghese J, Coleman S, Burn D, Rochester L. Independent domains of gait in older adults and associated motor and nonmotor attributes: validation of a factor analysis approach. *J Gerontol A Biol Sci Med Sci.* 2013;68(7):820-7.


60. Lord S, Howe T, Greenland J, Simpson L, Rochester L. Gait variability in older adults: a structured review of testing protocol and clinimetric properties. *Gait Posture*. 2011;34(4):443-50.

**Figure 1:** Flowchart of the selection process



**Figure 2:** Overview of the mean step-time parameters as a function of age

Dots are mean values; error bars represent SD of age (horizontal) or of the selected parameter (vertical)

**Figure 3:** Gait variability as a function of age

Dots are mean values; error bars represent SD of age (horizontal) or of the selected parameter (vertical)

**Table 1:** Characteristics of the individual studies

Author	Design	Type of instrumented walkway	Walkway length (m)	Footwear	Study sample characteristics								Origin		
					# Subjects	# Women	Age (years)			BMI (kg/m <sup>2</sup> )		Length (m)			
							Min	Max	Mean	SD	Mean	SD		Mean	SD
Beauchet et al. 2017	CS*	GaitRite	4.60 - 7.90		954	437	65	>85	72.8	4.8	26.2	4.1			International <sup>§</sup>
Chui et al. 2010	CS	Gaitrite	4.6		118	83	72	98	84.8 <sup>§</sup>	5.3 <sup>§</sup>	29.66 <sup>§</sup>	3.67 <sup>§</sup>			USA
Delbaere et al. 2009	CS	Gaitrite	7.2		44	27			76.8	5.2			1.68	0.09	Australia
Dujmovic et al. 2017	CC	Gaitrite	5.5		40	26			39.4	15.3					Serbia
Elboim-Gabyzon et al. 2017	CS	Gaitrite	/	barefoot	47	34			76.7	7.7	26.7	4.8	1.60	0.10	Israel
Gomez Bernal et al. 2016	CS	Opto gait	10	barefoot	126	85			27.4	1.8	22.69	2.56	1.68	0.04	Spain
Hollman et al. 2011	CS	Gaitrite	5.6		294	186	70	89	79	5	26.69	4.15	1.66	0.06	USA
Jenkins et al. 2009	CC	Gaitrite	6	shoes	40	25	49	77	64.7	7.7					USA
Khashan et al. 2014	CC	Gait Mat system	/	barefoot	40	14			64.9	10.2	26.4	3.8	1.68	0.08	Israel
Kodesh et al. 2015	CS	Gaitrite	5.17		40	25	18	40	26	4.7			1.69	0.10	Israel
Lee et al. 2017	CS	3D - Motion Analysis Corporation, Santa Rosa, CA, USA	30		30	15			74.1	7.6			1.61	0.08	South Korea
McClelland et al. 2011	CC	3D - Vicon	10		40	22			69.6	8.3			1.65	0.09	Australia
McGibbon et al. 2003	CC	3D - Selspot II optoelectric system	10		45	29			29.7	6.9	24.08	3.43	1.68	0.09	USA
					37	24			71.2	8.2	25.27	4.43	1.64	0.09	
Meldrum et al. 2014	CS	3D - Vicon	10		30	18			30	6.8	22.6		1.70		Ireland



Pistacchi et al. 2017	CC	BTS 3D GA system	10	44	22			67.0	9.4					Italy
Sturnieks et al. 2008 (b)	CC	3D - Vicon	/	42	17	24	56	37.6	7.5		1.74	0.08		Australia
Sturnieks et al. 2008 (a)	CC	3D - Vicon	10	47	18	23	56	38.2	7.9	24.9	3.6	1.74	0.09	Australia
Yang et al. 2016	CS	3D - vicon	14	54	27			23.9	4.7			1.67	0.10	USA

**Legend:** Data presented as Mean (SD); CS: Cross-sectional study design; CC: Case-control study design; m: meters; \*: combination of different cross-sectional studies; \$: Weighted mean; \$: data from Australia, Belgium, Canada, France, Japan, Luxembourg, Norway, USA, and Switzerland

**Table 2:** Overview of the mean spatio-temporal and spatial gait parameters

Author	Population Characteristics				Spatio-temporal parameters				Spatial parameters (m)						
	# Subjects	Age (years)		Walking speed (m/s)		Cadence (steps/min)		Step length		Stride length		Step width			
		min	max	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD		
Beauchet et al. 2017	711			70.6	2.40	<u>1.25</u>	<u>0.22</u>			<u>1.380</u>	<u>0.166</u>	<u>0.099</u>	<u>0.031</u>		
	207			77.6	2.60	<u>1.14</u>	<u>0.24</u>			<u>1.265</u>	<u>0.197</u>	<u>0.096</u>	<u>0.032</u>		
	36			87.7	2.80	<u>0.89</u>	<u>0.18</u>			<u>1.029</u>	<u>0.153</u>	<u>0.100</u>	<u>0.032</u>		
Chui et al. 2010	19	70	79			<u>1.39</u>	<u>0.24</u>	120.23	11.88	<u>0.689</u>	<u>0.088</u>	<u>1.380</u>	<u>0.175</u>	<u>0.095</u>	<u>0.025</u>
	77	80	89			<u>1.13</u>	<u>0.10</u>	111.89	3.91	<u>0.604</u>	<u>0.043</u>	<u>1.210</u>	<u>0.086</u>	<u>0.110</u>	<u>0.013</u>
	22	90	99			<u>0.87</u>	<u>0.17</u>	103.58	7.31	<u>0.495</u>	<u>0.079</u>	<u>0.994</u>	<u>0.158</u>	<u>0.109</u>	<u>0.033</u>
	118	70	99	84.8	5.30	<u>1.13</u>	<u>0.09</u>	111.68	3.69	<u>0.597</u>	<u>0.038</u>	<u>1.197</u>	<u>0.076</u>	<u>0.108</u>	<u>0.011</u>
Delbaere et al. 2009	44			76.9	5.10	<u>1.09</u>		106.41		<u>0.590</u>				<u>0.096</u>	
Dujmovic et al. 2017	40			39.4	15.3	<u>1.39</u>	<u>0.978 - 1.638</u>					<u>1.406</u>	<u>1.110 - 1.686</u>		
Elboim-Gabyzon et al. 2017	47			76.7	7.70	<u>0.99</u>	<u>0.03</u>	111.10	1.60			<u>1.055</u>	<u>0.022</u>		
Gomez Bernal et al. 2016	126			27.3	1.77	1.30	0.10	115.62	6.41	<u>0.681</u>	<u>0.040</u>	<u>1.358</u>	<u>0.080</u>		
Hollman et al. 2011	60	70	74			<u>1.17</u>	<u>0.18</u>	<b>108.05</b>	<b>14.60</b>	<u>0.646</u>	<u>0.086</u>	<u>1.302</u>	<u>0.157</u>	<u>0.082</u>	<u>0.033</u>
	107	75	79			<u>1.15</u>	<u>0.16</u>	<b>111.76</b>	<b>12.16</b>	<u>0.615</u>	<u>0.070</u>	<u>1.233</u>	<u>0.142</u>	<u>0.080</u>	<u>0.043</u>
	80	80	84			<u>1.06</u>	<u>0.16</u>	<b>106.76</b>	<b>8.54</b>	<u>0.596</u>	<u>0.075</u>	<u>1.203</u>	<u>0.154</u>	<u>0.094</u>	<u>0.041</u>

	47	85	89		<b>0.99</b>	<b>0.21</b>		<b>106.21</b>	<b>10.30</b>	<b>0.555</b>	<b>0.093</b>	<b>1.099</b>	<b>0.189</b>	<b>0.093</b>	<b>0.033</b>
Jenkins et al. 2009	40		64.7 3	7.66	<u>1.31</u>	<u>0.03</u>				<u>0.711</u>	<u>0.011</u>				
Kashan et al. 2014	40		64.9 0	10.2 0	<u>1.08</u>	<u>0.18</u>		109.70	9.70	<u>0.591</u>	<u>0.071</u>	<u>1.179</u>	<u>0.142</u>		
Kodesh et al. 2015	40	18	40	26.0 0	4.70	1.35	0.06								
Lee et al. 2017	30		74.0 7	4.14				105.61	5.46			<u>1.072</u>	<u>0.156</u>	<u>0.151</u>	<u>0.021</u>
McClelland et al. 2011	40		69.6 0	8.30	1.27	0.18		121.60	7.10			1.250	0.180		
McGibbon et al. 2003	45		29.7 0	6.90	1.32	0.16				0.674	0.058			0.109	0.041
	37		71.1 0	8.20	1.16	0.18				0.601	0.081			0.106	0.038
Meldrum et al. 2014	30		30.0 0	6.80	1.40	0.09		122.00	4.27	0.690	0.002	1.380	0.050	0.140	0.020
Pistacchi et al. 2017	44		67.0 0	9.42	1.33	0.06		113.84	4.30						
Sturnieks et al. 2008 (b)	42	24	56	37.6 0	7.50	1.50	0.17					1.490	0.170	0.070	0.047
Sturnieks et al. 2008 (a)	47	23	56	38.2 0	7.90	1.41	0.21	116.20	15.60			1.490	0.170	0.070	0.090
Yang et al. 2016	54		23.9 0	4.70	1.41	0.20									

**Legend:** underlined values = converted values (cm to m and ms to s); *italic values* = median values and interquartile range; **bold values** = weighted mean; m: meters; s: seconds; steps/min: steps per minute.

**Table 2 (continued):** Overview of the mean temporal gait parameters and percentage of the gait cycle

Author	Population characteristics				Temporal parameters (s)								Gait phases (%)												
	# Subjects	Age (years)			Step time		Stride time		Stance time		Single Support time		Swing time		Double Support time		Stance		Single Support		Swing		Double Support		
		<i>mi</i> <i>n</i>	<i>ma</i> <i>x</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mea</i> <i>n</i>	<i>SD</i>	<i>mean</i>	<i>SD</i>	<i>mean</i>	<i>SD</i>
Beauchet et al. 2017	711			70.6 0	2.40			<u>1.12</u>	<u>0.12</u>	<u>0.70</u>	<u>0.09</u>	<u>0.42</u>	<u>0.04</u>	<u>0.42</u>	<u>0.04</u>	<u>0.28</u>	<u>0.06</u>								
	207			77.6 0	2.60			<u>1.13</u>	<u>0.18</u>	<u>0.71</u>	<u>0.09</u>	<u>0.41</u>	<u>0.04</u>	<u>0.41</u>	<u>0.04</u>	<u>0.31</u>	<u>0.07</u>								
	36			87.7 0	2.80			<u>1.18</u>	<u>0.14</u>	<u>0.78</u>	<u>0.11</u>	<u>0.40</u>	<u>0.04</u>	<u>0.40</u>	<u>0.04</u>	<u>0.38</u>	<u>0.10</u>								
Chui et al. 2010	19	70	79														62.9 1	1.2 2	74.2 1	2.4 7	37.10 2	1.2 2	25.81 1	2.4 1	
	77	80	89														63.6 9	0.8 1	72.6 5	1.6 3	36.32 1	0.8 1	27.45 6	1.5 6	
	22	90	99														65.3 1	1.8 8	69.4 2	3.7 2	34.71 8	1.8 8	30.77 5	3.7 5	
	118	70	99	84.8 0	5.30												63.8 6	0.6 9	72.3 0	1.3 8	36.15 9	0.6 9	27.80 8	1.3 8	
Delbaere et al. 2009	44			76.9 0	5.10										0.31										
Dujmovic et al. 2017	40			39.4 0	15.3 0			<u>1.13</u>	<u>0.92</u> -					<u>0.39</u>	<u>0.3</u> <u>3-</u> <u>0.4</u> <u>5</u>	<u>0.25</u>	<u>0.16</u> <u>0.43</u>								
Elboim-Gabyzon et al. 2017	47			76.7 0	7.70																			28.70	0.6 1
Gomez Bernal et al. 2016	126			27.3 7	1.77	0.53	0.03	1.05	0.05	0.69	0.04	0.36	0.02	0.36	0.02	0.33	0.03								
Hollman et al. 2011	60	70	74			<b>0.56</b>	<b>0.06</b>	<b>1.11</b>	<b>0.11</b>	<b>0.71</b>	<b>0.09</b>	<b>0.41</b>	<b>0.05</b>	<b>0.41</b>	<b>0.04</b>	<b>0.30</b>	<b>0.06</b>	<b>63.2</b> <b>6</b>	<b>2.6</b> <b>5</b>	<b>37.0</b> <b>5</b>	<b>2.5</b> <b>7</b>	<b>36.60</b> <b>1</b>	<b>2.1</b> <b>1</b>	<b>26.76</b> <b>5</b>	<b>3.5</b> <b>5</b>
	107	75	79			<b>0.54</b>	<b>0.06</b>	<b>1.08</b>	<b>0.11</b>	<b>0.68</b>	<b>0.07</b>	<b>0.39</b>	<b>0.05</b>	<b>0.39</b>	<b>0.04</b>	<b>0.30</b>	<b>0.05</b>	<b>63.9</b> <b>3</b>	<b>2.8</b> <b>6</b>	<b>36.1</b> <b>4</b>	<b>3.9</b> <b>3</b>	<b>36.27</b> <b>8</b>	<b>2.5</b> <b>8</b>	<b>35.06</b> <b>5</b>	<b>5.2</b> <b>5</b>
	80	80	84			<b>0.57</b>	<b>0.05</b>	<b>1.13</b>	<b>0.09</b>	<b>0.72</b>	<b>0.07</b>	<b>0.40</b>	<b>0.04</b>	<b>0.40</b>	<b>0.04</b>	<b>0.32</b>	<b>0.06</b>	<b>64.1</b> <b>8</b>	<b>2.6</b> <b>5</b>	<b>36.0</b> <b>2</b>	<b>2.3</b> <b>1</b>	<b>36.01</b> <b>4</b>	<b>2.6</b> <b>4</b>	<b>28.26</b> <b>5</b>	<b>4.6</b> <b>5</b>
	47	85	89			<b>0.57</b>	<b>0.06</b>	<b>1.14</b>	<b>0.12</b>	<b>0.74</b>	<b>0.1</b>	<b>0.41</b>	<b>0.04</b>	<b>0.41</b>	<b>0.04</b>	<b>0.33</b>	<b>0.08</b>	<b>64.6</b> <b>2</b>	<b>2.5</b> <b>6</b>	<b>35.5</b> <b>5</b>	<b>2.5</b> <b>9</b>	<b>35.52</b> <b>3</b>	<b>2.6</b> <b>3</b>	<b>29.18</b> <b>1</b>	<b>4.4</b> <b>1</b>

Jenkins et al. 2009	40			64.7 3	7.66					0.41	0.0 3									
Kashan et al. 2014	40			64.9 0	10.2 0								60.5	1.7	39.4 0	1.2 0	39.50	1.7 0	21.20	2.7 0
Lee et al. 2017	30			74.0 7	4.14										35.1 5	3.5 3				
McGibbon et al. 2003	45			29.7 0	6.90					0.66	0.0 7									
	37			71.1 0	8.20					0.68	0.0 6									
Meldrum et al. 2014	30			30.0 0	6.80	0.50	0.0 2	0.99	0.03				0.40	0.0 2					0.19	0.02
Sturnieks et al. 2008 (a)	47	23	56	38.2 0	7.90					0.67	0.0 6									
Yang et al. 2016	54			23.9 0	4.70			1.06	0.10				0.33	0.0 3					0.20	0.03

Legend: underlined values = converted values (cm to m and ms to s); *italic values* = median values and interquartile range; **bold values** = weighted mean; s: seconds.

**Table 3:** overview of gait variability measures

Author	Population characteristics					Gait Variability (coefficient of variation. %)													
	# Subject	Age (years)				CoV Step length		CoV stride length		CoV BoS		CoV stride time		CoV Swing time		CoV Stance time		CoV DS time	
		Min	Max	Mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
Beauchet et al. 2017	711			70.6 0	2.40			2.20	1.10	24.60	34.7 0	2.10	1.10	4.00	1.70	3.10	1.3 0	6.80	2.90
	207			77.6 0	2.60			2.60	1.30	33.00	82.6 0	2.30	1.10	4.50	1.70	3.20	1.5 0	6.30	2.50
	36			87.7 0	2.80			3.60	2.10	28.20	23.4 0	2.80	1.30	6.00	2.70	3.50	1.7 0	6.00	2.10
Dujmovic et al. 2017	40			39.4 0	15.3 0			<u>2.20</u>	<u>1.00 - 6.82</u>			<u>2.23</u>	<u>1.20 - 11.08</u>	<u>4.38</u>	<u>2.69 - 18.63</u>			<u>9.78</u>	<u>4.48 - 34.09</u>
Hollman et al. 2011	60	70	74			<b>6.31</b>	<b>9.40</b>	<b>3.56</b>	<b>3.08</b>	<b>3.27</b>	<b>2.31</b>	<b>3.72</b>	<b>3.14</b>	<b>5.44</b>	<b>6.77</b>	<b>5.34</b>	<b>4.4 9</b>		

107	75	79	<b>5.62</b>	<b>5.94</b>	<b>4.56</b>	<b>5.32</b>	<b>3.12</b>	<b>2.33</b>	<b>4.81</b>	<b>6.20</b>	<b>7.38</b>	<b>9.02</b>	<b>5.47</b>	<b>5.63</b>
80	80	84	<b>5.53</b>	<b>2.75</b>	<b>4.07</b>	<b>2.05</b>	<b>3.46</b>	<b>2.73</b>	<b>3.89</b>	<b>1.90</b>	<b>5.69</b>	<b>2.20</b>	<b>5.02</b>	<b>2.95</b>
47	85	89	<b>6.08</b>	<b>2.70</b>	<b>5.35</b>	<b>4.60</b>	<b>2.94</b>	<b>1.13</b>	<b>5.02</b>	<b>4.82</b>	<b>8.18</b>	<b>10.21</b>	<b>5.51</b>	<b>3.74</b>

**Legend:** underlined values = converted values (cm to m and ms to s); *italic values* = median values and interquartile range; **bold values** = weighted mean