

This item is the archived peer-reviewed author-version of:

Relationship between pesticide accumulation in transplanted zebra mussel (***Dreissena polymorpha***)
and community structure of aquatic macroinvertebrates

Reference:

Bashnin Tayebah, Verhaert Vera, De Jonge Maarten, Vanhaecke Lynn, Teuchies Johannes, Bervoets Lieven.- Relationship between pesticide accumulation in transplanted zebra mussel (***Dreissena polymorpha***)) and community structure of aquatic macroinvertebrates
Environmental pollution - ISSN 0269-7491 - 252:A(2019), p. 591-598
Full text (Publisher's DOI): <https://doi.org/10.1016/J.ENVPOL.2019.05.140>
To cite this reference: <https://hdl.handle.net/10067/1599910151162165141>

1 Relationship between pesticide accumulation in transplanted zebra mussel (*Dreissena*
2 *polymorpha*) and community structure of aquatic macroinvertebrates

3 Tayebah Bashnin¹, Vera Verhaert¹, Maarten De Jonge², Lynn Vanhaecke³, Johannes

4 Teuchies¹, Lieven Bervoets¹

5 ¹ Laboratory of Systemic, Physiological and Ecotoxicological Research, Department of
6 Biology, University of Antwerp, Groenenborgerlaan, 171 2020, Antwerp, Belgium

7 ² Flemish Environment Agency, Department of water monitoring, Diestsepoort 6, bus 73,
8 3000 Leuven

9 ³ Laboratory of Chemical Analysis, Department of Veterinary Public Health and Food Safety,
10 University of Ghent, Salisburylaan 133, B-9820 Merelbeke, Belgium

11 E-mail: tayebah.bashnin@uantwerpen.be

12 **Abstract:**

13 This study examined to what degree bioaccumulated pesticides in transplanted zebra mussels
14 can give an insight to pesticide bioavailability in the environment. In addition, it was
15 investigated if pesticide body residues could be related to ecological responses (changes in
16 macroinvertebrate community composition). For this at 17 locations, 14 pesticide
17 concentrations and nine dissolved metals were measured in translocated zebra mussels and the
18 results were related to the structure of the macroinvertebrate community. Critical body burdens
19 in zebra mussel, above which the ecological status was always low, could be estimated for
20 chlorpyrifos, terbuthylazine and dimethoate being respectively 8.0, 2.08 and 2.0 ng/g dry
21 weight.

22 With multivariate analysis, changes in the community structure of the macroinvertebrates were
23 related to accumulated pesticides and dissolved metals. From this analysis, it was clear that the
24 composition of the macroinvertebrate communities was not only affected by pesticides but also
25 by metal pollution. Two different regions could be clearly separated, one dominated by metal
26 pollution, and one where pesticide pollution was more important.

27 The results of this study demonstrated that zebra mussel body burdens can be used to measure
28 pesticide bioavailability and that pesticide body burdens might give insight in the ecological
29 impacts of pesticide contamination. Given the interrelated impacts of pesticides and heavy
30 metals, it is important to further validate all threshold values before they can be used by
31 regulators.

32 **Capsule:** High zebra mussels pesticide body burden proved to be a good predictor for low
33 environmental quality

34 **Keywords:** Pesticides, Dissolved metals, Bioavailability, Ecological status, Biomonitoring

35 **1 Introduction**

36 The extensive application of pesticides for agricultural and non-agricultural uses results in
37 elevated concentrations of pesticides and their residues in surface and groundwater resources.
38 Pesticide residues have been reported as common organic contaminants worldwide (Aktar et
39 al., 2009, Ali et al., 2014, Szekacs et al., 2015). Not all pesticides are easily degradable and due
40 to their lipophilic properties, they can bioaccumulate in organisms, biomagnify in food chains,
41 and consequently influence the health of organisms including humans (Andreu and Pico, 2012,
42 Connell, 1988, Emanuela et al., 2017). Elevated pesticide levels may have adverse effects on
43 macroinvertebrate communities including the loss of certain sensitive taxa (Munze et al., 2017,
44 Schafer et al., 2012). As such, biological indices, based on the occurrence of macroinvertebrate
45 species might be a good indicator of the ecological impact of pesticides. However, other
46 stressors are present as well and there is no direct causal relationship between the measured
47 pesticide concentrations and the macroinvertebrate community. Accumulated pesticides, on the
48 other hand, can provide a better insight on bioavailability and ecological risks of pesticides.
49 Recent studies have shown that accumulated micropollutants can be used to predict ecological
50 effects on macroinvertebrate community (Bervoets et al., 2016, Luoma et al., 2010, Rainbow
51 et al., 2012) and allow to set safe body burden thresholds. The use of invertebrate body burdens
52 as an indicator for metal toxicity was evaluated in several studies. In some studies, a significant
53 relationship has been determined between metal accumulation in caddisfly (*Hydropsyche* sp.)
54 and ecological responses such as a decrease in invertebrate taxa richness (Luoma et al., 2010,
55 Rainbow et al., 2012). Also, metal concentrations in caged zebra mussels were found to be
56 predictive for the ecological quality of a river along a metal pollution gradient (De Jonge et al.,
57 2012).

58 Active biomonitoring using transplanted organisms has been used for more than two decades
59 to assess bioavailability and effects of micropollutants in aquatic ecosystems (Bervoets et al.,

60 2004, De Jonge et al., 2012, Mersch, 1996). Active biomonitoring can provide an integrative
61 measure of bioavailable pesticides in the aquatic environment and help to detect pollutants that
62 are usually present in the surface water at concentrations below detection limits of routine
63 analytical methods (Bervoets et al., 2005a, Szekacs et al., 2015). In this regard, zebra mussel
64 *Dreissena polymorpha* has been suggested as a reliable freshwater biomonitoring organism
65 (Bervoets et al., 2005b, Bourgeault et al., 2010, De Jonge et al., 2012, Kraak, 1994).

66 Understanding the relationship between pesticide accumulation in zebra mussels and their
67 effects on aquatic communities such as macroinvertebrates can provide better insight into
68 ecological risks from pesticide contamination and is crucial for accurately monitoring and
69 predicting the impacts of these contaminants in aquatic environments (Rainbow et al., 2012).

70 Although the use of invertebrate body burdens was tested in different studies as an indicator of
71 the ecological effects of metal toxicity, this approach has not been tested yet for effects of
72 organic micropollutants including pesticides.

73 The objective of this study was to investigate to what extent a selection of accumulated
74 pesticides in transplanted zebra mussels reflects bioavailable pesticide concentrations in the
75 environment, and how the tissue residues can be related to alterations in macroinvertebrate
76 community structure. Additionally, it was investigated whether we could estimate accumulated
77 pesticide concentrations (thresholds) above which ecological quality is always low. Since in
78 many cases pesticide pollution will not be the only stressor, the contribution of other stressors,
79 i.e. accumulated metals on the invertebrate communities has been investigated as well.

80 **2 Material and methods**

81 **2.1 Study area and sampling sites**

82 Adult individuals of zebra mussels (*Dreissena polymorpha*) were collected in September 2013
83 from a shallow lake, Blaarmeersen (Gent, Belgium) and transported to the lab. From former
84 analysis (2011; unpublished data) it was demonstrated that these mussels were uncontaminated

85 with regard to metals, polychlorinated biphenyls (PCBs), organic chloride pesticides (OCPs)
86 and polyaromatic hydrocarbons (PAHs). All mussels were kept in dechlorinated, aerated and
87 filtered tap water in plastic tanks (1000 liter) for at least one month at 15 °C and fed with a
88 mixture of algae (*Pseudokirchneriella subcapitata* and *Chlamydomonas reinhardtii*).

89 After this acclimatization period, zebra mussels were transplanted to cages deployed at 17 sites
90 in Flanders (Belgium) in October 2013 (Figure 1, Table 1). Nine sites were located in the
91 southeast (1-9) and 6 sites in the northeast (10-15) of Flanders. In addition, two sites in the west
92 of Flanders were selected, one in the upper part of the Scheldt (16; southwest) and one in the
93 canal Ghent Terneuzen (17; northwest). Sites 1 to 9 were selected because of a broad range in
94 pesticide pollution (Flemish Environment Agency data), whereas lower pesticide
95 concentrations were expected at sites 10 to 15 although elevated metal concentrations might be
96 present and even follow a gradient of high zinc and cadmium levels at site 10 which gradually
97 decrease until site 15 (Bervoets et al., 2005a). Sites 16 and 17 were selected because of general
98 industrial pollution.

99 Before the start of the exposure 15 mussels of comparable length (15-20 mm) were collected
100 and soft tissue was pooled for analysis of background metals and pesticides.

101 At every site, 24 individual mussels of comparable length (15-20 mm) were randomly placed
102 in 2 polyethylene cages (11 × 11 × 22 cm with a mesh size of 2 × 4 mm), which allowed free
103 circulation of water (Bervoets et al., 2004). To standardize the exposure and to prevent the
104 accumulation of sediment all cages were free floating at a maximal depth of 30 cm below the
105 water surface. After an exposure period of six weeks, all mussels were collected and processed
106 (see below).

107 **2.2 Macroinvertebrate sampling**

108 Benthic macroinvertebrates were sampled at each site using a standard invertebrate hand net
109 (500 mm-mesh, 200-300 mm frame and 500 mm bag depth). The samples were collected in

110 October 2013 covering a river length of 20 m for 5 min (Depauw and Vanhooren, 1983). In the
111 lab, samples were rinsed with tap water through a sieve of 0.5 mm to remove sediment and the
112 organic fraction. The invertebrates were transferred to a white tray with gridlines and sorted,
113 counted and transferred into small glass vials with denatured ethanol (70%) for fixation. The
114 macroinvertebrates were identified up to family or genus level according to De Pauw and
115 Vannevel (1991). The Multimetric Macroinvertebrate Index Flanders (MMIF, Gabriels et al.
116 (2010)) was used to assess the biological water quality. This index combines 5 different
117 characteristics of the macroinvertebrate community, i.e.: the total number of taxa (taxa
118 richness), the number of EPT taxa (Ephemeroptera, Plecoptera and Trichoptera), the number
119 of other sensitive taxa, not belonging to the EPT, the mean tolerance score and the Shannon-
120 Wiener diversity index. For sites 16 and 17 existing MMIF data originating from the Flemish
121 Environment Agency were used (www.vmm.be/geoview).

122 **2.3 Pesticide analysis**

123 After exposure, the zebra mussels were collected and in the lab placed in 10 L buckets at 15 °C
124 in filtered (0.45 µm) site water for 18 h to allow depuration. After removal of the byssus threads,
125 the soft tissue of the mussels was dissected and rinsed with MQ-water. All mussels were pooled
126 per site and homogenized by a rotor-stator homogenizer (Tissue Ruptor, Qiagen). The
127 homogenized tissue was transferred to polypropylene vials (50 ml) and kept at -80 °C until
128 freeze dried. For the pesticide analysis, the procedure of Wille et al. (2011) was adapted:
129 samples were freeze-dried for 48 h and the dry tissue was extracted by a combination of
130 pressurized liquid extraction (PLE) and solid phase extraction (SPE). All solvents were
131 prepared according to Wille et al. (2011). The PLE was performed on a Dionex ASE® 350
132 accelerated Extractor with Solvent Controller (Dionex Crop., Sunnyvale, CA, USA). A
133 cellulose filter (27 mm, Dionex Crop) was located on the bottom of a 22 mL stainless extraction
134 cell. Subsequently, 0.25 g of a sample (freeze-dried sample), 2 g of aluminum oxide and 4.5 g

135 of diatomaceous earth were placed in each Extraction cell. Each extraction cell was spiked with
136 the internal standards at a concentration of 100 ng/g (100 ppb). The ASE-extract obtained was
137 reduced to 0.5 mL by evaporation under nitrogen at 40 °C and subsequently diluted with 10 mL
138 of ultra-pure water. SPE was performed using Isolute EnV+ cartridges (6 mL, 200 mg, Biotage,
139 Uppsala, Sweden). The cartridges were conditioned with 10 mL of methanol and 10 mL of
140 ultra-pure water with methanol (5%). Further, the cartridges were loaded with the sample and
141 then washed with 5 mL of ultra-pure water. Elution was carried out by 5 mL of methanol and
142 acetonitrile. Next, the eluate was evaporated under nitrogen at 40 °C to 1 droplet and
143 reconstituted in 50 µL methanol and 150 µL of 2 mM aqueous ammonium carbonate.

144 Analysis for the following pesticides was carried out: dichlorvos, dimethoate, diazinon,
145 pirimicarb, linuron, metolachlor, chloridazon, chlorpyrifos, simazine, isoproturon,
146 terbuthylazine, diuron, atrazine, and kepone. Atrazine-D₅ and isoproturon-D₆ were used as an
147 internal standard. Separation of pesticides was conducted by Ultra-high performance liquid
148 chromatography (U-HPLC). The apparatus consisted of an Accela™ High-Speed LC and an
149 Accela™ Autosampler (Thermo Scientific, San Jose, CA, USA). Pesticides detection was
150 performed using a TSQ Vantage Triple-Stage Quadrupole Mass Spectrophotometer (Thermo
151 Electron) equipped with a heated electrospray ionization probe (HESI-II). Separation,
152 detection, identification and quantification of target analytes followed the same methods as
153 described in Wille et al. (2011). Based on the concentration range found in Van Praet et al.
154 (2014) we expected the accumulated concentrations to be lower than 100 ng/g. Therefore we
155 slightly modified the method of Wille et al. (2011) by spiking the samples (freeze-dried
156 mussels) with a standard mixture to obtain final concentrations in the range of 0 to 100 ng/g
157 instead of 0.1 to 250 ng/g.

158 Quality assurance was performed in the same way as explained in Wille et al. (2011). However,
159 the concentration ranges at which samples were spiked with standard solution ranged from 1 to
160 100 ppb instead of 0.1 to 100 ppb.

161 **2.4 Metal analysis**

162 Since metal pollution was known to be present at some sites, the mussel tissue was analyzed
163 for the presence of nine metals. From each pooled sample (see above) a sub-sample of about
164 0.5 g was transferred into pre-weighed acid-washed polypropylene vials and dried for at least
165 24 h at 60 °C. After re-weighing, the samples were digested using 500 µL ultra-pure nitric acid
166 (HNO₃, 69%) followed by microwave-heating (Bervoets et al., 2005a). Metal Ag, Al, Cd, Cr,
167 Cu, Fe, Ni, Pb and Zn) were measured using a quadrupole inductively coupled plasma mass
168 spectrometer (ICP-MS; Agilent 7700x ICP-MS, Santa Clara, USA). Quality control included
169 analysis of procedural blanks and certified reference material metals (mussel tissue; BCR278R)
170 from the community bureau of reference (European Union, Brussels, Belgium) (Bervoets et al.,
171 2005b). Recoveries were all within 15% of the certified values (recoveries of 90 – 112%).

172 **2.5 Statistics**

173 Relationships between the individual accumulated pesticides and the calculated indices were
174 visualized using scatterplots. Quantile regression was used to investigate the possible
175 relationship between invertebrate body burdens and calculated indices. Quantile regression
176 analysis allows to calculate the 90th quantile of ecological responses (calculated indices) as a
177 function of an environmental stressor (zebra mussel body burden) and therefore to determine
178 the maximum ecological response. Using the maximal response can reduce the influence of
179 non-modeled factors that might affect invertebrate taxa richness. The detailed information
180 regarding this model is described in De Jonge et al. (2013).

181 To find the concentration threshold, above which a good ecological status was never reached,
182 concentrations were selected where the MMIF score was higher than 0.6. The concentration
183 thresholds were set as the 95th percentile of the internal pesticides concentrations.

184 Among the detected pesticides, chlorpyrifos, dichlorvos, terbuthylazine and dimethoate were
185 used to investigate the relationship between pesticides and biological water indices since they
186 have been found in most of the sampling sites. Pesticide concentrations below the limit of
187 quantification (LOQ) were replaced by LOQ/2 (Bervoets et al., 2004, Custer et al., 2000).

188 In order to relate changes in macroinvertebrate community composition to accumulated
189 pesticides and metals, direct ordination techniques (canonical correspondence analysis; CCA)
190 were used. Before the analysis, all data were log (X+1) transformed and the gradient length
191 within the taxa data was analyzed. Detrended correspondence analysis (DCA) revealed that the
192 total gradient length of species distribution was > 2 times standard deviation. Therefore, the
193 unimodal (Gaussian distribution) response model was selected. For CCA, interspecies distance
194 and Hill's scaling were applied. Down-weighting of rare species was used to give less weight
195 to taxa which occurred only infrequently. All ordination models were calculated using
196 CANOCO version 4.5.

197 **3 Results**

198 **3.1 Background information**

199 The environmental variables (temperature (°C), pH, oxygen content (mg/L) and conductivity
200 (µS/cm) of the surface water at the sampling sites are presented in Table 1. The lowest pH was
201 6.82 observed at site 4. The oxygen concentration was the lowest with 5.3 mg/L at site 3 and
202 the highest with 11.3 mg/L at site 13. The maximum conductivity was 892 µs/cm, observed at
203 site 5. For most sites, the water oxygen level met the water quality standard (≥ 6 mg/L; ≤ 12
204 mg/L) with the exception of sites 2 and 3. Conductivity was well below the quality standard of
205 1000 µS/cm at all sites (Flemish Government, 2000).

206 **3.2 Pesticide and metal accumulation in zebra mussels;**

207 The results of the concentrations of pesticides and metals in mussels exposed at the different
208 sites are presented in table SI. 1 and SI. 3 respectively. The concentration in mussels from
209 Blaarmeersen after one month of acclimatization (called 'Start' in the tables) are reported as
210 well. Among the fourteen pesticides analyzed for in the present study, simazine and kepone
211 were below the detection limit in mussel tissue at all sites, while dimethoate, dichlorvos,
212 pirimicarb, terbuthylazine, chlorpyrifos were detectable in mussel tissue at most of the sampling
213 sites (Figure 2). The concentrations of diuron, linuron, metolachlor, isoproturon, atrazine and
214 diazinon were below the LOQ at most of the sites. The LOQ of the analyzed pesticides was 1
215 ng/g except for dimethoate and simazine which had a LOQ of 10 and 5 ng/g respectively. It
216 should be noted that high background concentrations of pirimicarb (5.48 ng/g dry weight) in
217 mussels were measured (in mussels originating from Blaarmeersen used for exposure at the
218 other sites; called "start" in SI. 1).

219 In mussels from the sites 2 and 3 nearly all measured pesticides were higher than LOQ while at
220 some sites (9,10, 11, 13, 14, 15) almost all pesticides were below the LOQ. The highest
221 concentration of pesticides found was for chlorpyrifos (2203 ng/g dry weight and 103 ng /g dry
222 weight at site 2 and 3 respectively) and for dichlorvos with concentrations up to 128 ng /g dry
223 weight. Accumulated metals are summarized in SI. 3. Elevated concentration of Cd, Pb and Zn
224 were measured in mussels from 2, 11, 12, 13, 14 and 15.

225 **3.3 Macroinvertebrate community composition**

226 In total, 34 macroinvertebrate taxa have been identified. The highest abundance was observed
227 for Gammaridae and Chironomidae representing > 95% of all invertebrates at site 9 and 12
228 respectively (SI. 2). The highest taxonomic richness for Trichoptera and Ephemeroptera was
229 observed at sites 1, 7 and 14, while no EPT families were present at sites 2-4 and 12. Plecoptera
230 was not found at pesticide nor metal polluted sites. The presence of the EPT taxa resulted in a

231 high score for 1, 7, 10 and 14. The highest total number of taxa was observed at site 1 and the
232 lowest at sites 2 and 3. According to the MMIF index 1, 9 and 10 were classified as having a
233 good biological quality ($MMIF \geq 0.7$), the sites 2, 3, 4, 12, 13 and 17 had a bad biological
234 ($MMIF \leq 0.3$) water quality and the remaining sites were considered as having a moderate
235 biological water quality (Table 1).

236 At the agricultural sites (1-9), predominant taxa were Simuliidae, Gammaridae and Tubificidae
237 while Chironomidae and Asellidae were present in higher abundance at the metal polluted sites
238 (SI. 2).

239 **3.4 Relationship between pesticide accumulation in zebra mussel and community** 240 **structure**

241 The relationship between accumulated pesticide concentrations and the biological water quality
242 index (MMIF) was investigated for 4 pesticides (dichlorvos, dimethoate, terbuthylazine and
243 chlorpyrifos), as many values for the other pesticides were below LOQ (Figure 3). No
244 significant relationships could be found (quantile regression analysis, $p > 0.05$). However, it
245 was possible to derive accumulated pesticide threshold concentrations for three pesticides
246 above which the ecological quality was always low ($MMIF < 0.6$), i.e. terbuthylazine,
247 chlorpyrifos and dimethoate. Threshold values were calculated as the 95th percentile of the
248 accumulated pesticide concentrations in zebra mussels collected at sites with good biological
249 quality ($MMIF \geq 0.7$) (Table 2).

250 The CCA diagram that relates accumulated metals and pesticides to the macroinvertebrate
251 composition (Figure 4) reveals that the macroinvertebrate taxa distribution in sampling sites
252 was influenced by accumulated pesticides and metals in zebra mussels, explaining respectively
253 26.4% and 15.2% of the variation. The sites which are dominated by pesticide pollution or
254 metal pollution are separated in the CCA diagram. The sites 1, 3, 4-7 were more influenced by

255 pesticides contamination and sites 2, 10-12 and 14-15 were more subjected to metal
256 contamination.

257 In addition, Simuliidae and Planorbidae are more associated with sites where the concentration
258 of dichlorvos, terbuthylazine and chloridazon were high, whereas sites with a high
259 concentration of Pb and Cd were mainly dominated by Chironomidae and Asellidae (Figure 4).
260 At sites 2-4 where the concentration of chlorpyrifos was high and 12 with high Zn
261 concentrations, no EPT taxa (Ephemeroptera, Plecoptera and Trichoptera) occurred.

262 **4 Discussion**

263 **4.1 Pesticide accumulation in zebra mussels**

264 Literature on bioaccumulation of pesticides in freshwater biota, mainly invertebrates is limited.
265 Wille et al. (2011) measured the same set of pesticides as this study in the marine mussel
266 (*Mytilus edulis*) for an active biomonitoring program in the Belgian coastal zone. Only
267 dichlorvos and chloridazon were found with concentrations four times lower (dichlorvos) or
268 similar (chloridazon) compared with the present study. Van Praet et al. (2014) measured the
269 same set of pesticides in the damselfly larvae *Ischnura elegans* (Zygoptera, Odonata) collected
270 from sixteen ponds in Flanders. They reported four of the measured pesticides (chloridazon,
271 dichlorvos, terbuthylazine and metolachlor) above the LOQ. In the present study, metolachlor
272 in zebra mussel tissue was below LOQ at most sites, terbuthylazine and chloridazon
273 concentrations were comparable to the concentrations in the damselfly larvae, while for
274 dichlorvos, the concentration in the present study was twice as high. Neither Wille et al. (2011)
275 nor Van Praet et al. (2014) detected simazine and kepone in invertebrates collected during their
276 studies which is in agreement with the present study.

277 In a study by Miranda et al. (2008) three pesticides (atrazine, diuron, terbuthylazine) were
278 detected in the muscle tissue of freshwater trahira fish (*Hoplias malabaricus*) from Ponta

279 Grossa Lake in south Brazil. Diuron and terbuthylazine were detected in the same range as in
280 the present study while atrazine concentration was 2 times lower.

281 It should be noted that, although pirimicarb was found at all sampling sites in the present study,
282 the measured accumulated concentration might not represent the site-specific bioavailability,
283 since the concentration was already high (highest) at the start of the exposure in the mussels
284 originating from Blaarmeersen (“start”). At most sites, however, the pirimicarb was lower in
285 the mussel after 6 weeks of exposure compared to the start, so was probably eliminated from
286 the mussel tissue. Almost all measured pesticides in this study (except dimethoate) had octanol-
287 water partition coefficient greater than 1,000 (SI. 4) and a soil half-life greater than 30 days.
288 These properties result in accumulation in sediment and aquatic biota (Andreu and Pico, 2012,
289 Van Praet et al., 2014). As most pesticides are difficult to quantify in surface water samples due
290 to their hydrophobic characteristics (Wille et al., 2011), the results of this study suggest that
291 using accumulated concentrations in zebra mussels is a good monitoring strategy.

292 **4.2 Relationship between pesticide accumulation in zebra mussel and community** 293 **structure**

294 The ecological value of the investigated water bodies was always low at high accumulated
295 concentrations of terbuthylazine, chlorpyrifos and dimethoate. Based on the results, pesticide
296 threshold values of tissue burdens ranging from 2 to 8.0 ng/g dry weight have been calculated
297 and are indicative of pesticide levels which is harmful to macroinvertebrates communities. The
298 threshold concentrations indicate safe values above which a good ecological status was never
299 reached. However, this is a conservative approach as the lower ecological quality is not directly
300 related to the presence of pesticides as also other stressors are present.

301 Mayon et al. (2006) observed a lower IBI score (Index of Biotic Integrity or Fish Index) in a
302 river polluted with pesticides, compared to a reference site. On the other hand, in a study of
303 Eaton and Lydy (2000) no relationship was found between the IBI score and the present

304 organochlorine pesticides in fish tissue on twenty sampling sites in the Arkansas river in
305 Wichita, Kansas (USA). In the present study low MMIF values were observed at sites with low
306 accumulated pesticide concentrations, indicating that other factors also negatively affect the
307 macroinvertebrate community. Bervoets et al. (2016) and Van Ael et al. (2014) pointed out that
308 the decrease in ecological quality may also be caused by others stressors such as other
309 contaminants, physical characteristics of the water body or food and habitat availability. Also,
310 the CCA analysis revealed that the macroinvertebrate community was affected not only by
311 pesticide contamination but also by metals. The pesticide-polluted and the metal-polluted sites
312 are clearly separated from each other which was expected since sites 1-9 were located in a fruit
313 cultivation area with intensive pesticide application (information Flemish Environment
314 Agency) while sites 10, 12, 13, 14 and 15 were situated in a metal contaminated river system
315 (Bervoets et al., 2005a, Michiels et al., 2017). However, even within the pesticide-polluted sites,
316 accumulated Cd, Mn, Ni and Cu concentrations at site 2 were elevated compared to
317 uncontaminated Flemish lowland rivers and comparable to values in other metal contaminated
318 sites in Flanders (Bervoets et al., 2005a).

319 Based on the quantile regression analysis, no significant negative relations were found between
320 maximal (90th quantile) ecological response and accumulated pesticides in zebra mussel. This
321 might be due to the fact that only 15 sites were considered.

322 The applicability of body tissue residues to biomonitor metal toxicity in aquatic ecosystems has
323 been described before. De Jonge et al. (2013) found significant negative relationships between
324 accumulated Cu, Pb and Zn concentrations in *Leuctra* sp., Simuliidae, *Rhithrogena* sp. and
325 Perlodidae and the maximal ecological response. In addition, De Jonge et al. (2012) observed
326 significant relationships between accumulated metal concentrations in *D. polymorpha* and
327 Chironomidae and biological community response. Luoma et al. (2010) and Rainbow et al.
328 (2012), showed that metal accumulation in larvae of caddisfly *Hydropsyche* sp. was highly

329 correlated with ecological indicators such as mayfly richness and macroinvertebrates taxa,
330 whereas Bervoets et al. (2016) could estimate safe body concentrations for a set of metals in
331 Chironomidae larvae and Tubifidae worms. Awrahman et al. (2016) stated that the
332 bioaccumulated concentrations of metals in larvae of *Hydropsyche sp.* can be used to predict
333 reductions in local mayfly (particularly ephemereleid and heptageniid) abundance.

334 According to Weber et al. (2018), the most sensitive invertebrate species to organic pesticides
335 belong to the EPT taxa. Thiery and Schulz (2004) and Colville et al. (2008) similarly found a
336 high sensitivity of Ephemeroptera to pesticide pollution mainly to chlorpyrifos. Our results are
337 in good agreement with their results.

338 As insects, Chironomidae may have been sensitive to insecticides such as chlorpyrifos.
339 However, according to (Macchi et al., 2018), they were the most abundant taxa in a stream with
340 high detected pesticides including chlorpyrifos. The result of the present study showed that
341 Chironomidae are related rather to high metal concentrations than to high pesticide levels. This
342 is in accordance with Bervoets et al. (2016) who found very high accumulated metals in
343 Chironomidae.

344 In contrast to the study by Macchi et al. (2018) and Von Der Ohe and Liess (2004) who found
345 the molluscs being the most tolerant taxa towards pesticides, we did not find molluscs at most
346 of the studied sites, which of course could be due factors other than pesticide pollution. This is
347 supported by the fact that at all sites the caged mussels survived during the 6 weeks exposure
348 period.

349 **5 Conclusion**

350 From fourteen analyzed pesticides in zebra mussel tissue, dimethoate, dichlorvos, pirimicarb,
351 terbuthylazine, chlorpyrifos were detected at most of the sampling sites. The results of pesticide
352 body burdens suggest that using the zebra mussel is a promising monitoring strategy to measure
353 bioavailable pesticides that are difficult to quantify in water.

354 The results of the present study also demonstrate that for four out of the five detectable (not for
355 dichlorvos) pesticides the ecological status was always low (MMIF <0.6) when accumulated
356 concentrations in zebra mussel were detected. Thus, we suggest that body residues of pesticides
357 in zebra mussels can be used to predict the ecological effects of pesticides on the
358 macroinvertebrate community. However, to exclude possible effects of co-occurring stress
359 factors such as dissolved metals, more research and larger database mainly are needed.
360 Additional, evidence is needed to assess the effects of the accumulated pesticides in the
361 environment.

362 **Acknowledgments:**

363 We would like to thank Melvice Epede for helping with sampling and sorting
364 macroinvertebrates. Furthermore, we would like to thank Dr. Valentine Mubiana and Steven
365 Joose for the metal analysis.

366 **References:**

- 367 AKTAR, M. W., SENGUPTA, D. & CHOWDHURY, A. 2009. Impact of pesticides use in agriculture: their
 368 benefits and hazards. *Interdisciplinary toxicology*, 2(1), 1-12.
- 369 ALI, U., SYED, J. H., MALIK, R. N., KATSOYIANNIS, A., LI, J., ZHANG, G. & JONES, K. C. 2014.
 370 Organochlorine pesticides (OCPs) in South Asian region: A review. *Science of the Total Environment*,
 371 476, 705-717.
- 372 ANDREU, V. & PICO, Y. 2012. Determination of currently used pesticides in biota. *Analytical and*
 373 *Bioanalytical Chemistry*, 404, 2659-2681.
- 374 AWRAHMAN, Z. A., RAINBOW, P. S., SMITH, B. D., KHAN, F. R. & FIALKOWSKI, W. 2016. Caddisflies
 375 Hydropsyche spp. as biomonitors of trace metal bioavailability thresholds causing disturbance in
 376 freshwater stream benthic communities. *Environmental Pollution*, 216, 793-805.
- 377 BERVOETS, L., DE JONGE, M. & BLUST, R. 2016. Identification of threshold body burdens of metals for the
 378 protection of the aquatic ecological status using two benthic invertebrates. *Environmental Pollution*,
 379 210, 76-84.
- 380 BERVOETS, L., KNAEPKENS, G., EENS, M. & BLUST, R. 2005a. Fish community responses to metal
 381 pollution. *Environmental Pollution*, 138, 338-349.
- 382 BERVOETS, L., MEREGALLI, G., DE COOMAN, W., GODDEERIS, B. & BLUST, R. 2004. Caged midge
 383 larvae (*Chironomus riparius*) for the assessment of metal bioaccumulation from sediments in situ.
 384 *Environmental Toxicology and Chemistry*, 23, 443-454.
- 385 BERVOETS, L., VOETS, J., COVACI, A., CHU, S. G., QADAH, D., SMOLDERS, R., SCHEPENS, P. &
 386 BLUST, R. 2005b. Use of transplanted zebra mussels (*Dreissena polymorpha*) to assess the
 387 bioavailability of microcontaminants in Flemish surface waters. *Environmental Science & Technology*,
 388 39, 1492-1505.
- 389 BOURGEAULT, A., GOURLAY-FRANCE, C., VINCENT-HUBERT, F., PALAIS, F., GEFFARD, A.,
 390 BIAGIANTI-RISBOURG, S., PAIN-DEVIN, S. & TUSSEAU-VUILLEMIN, M. H. 2010. Lessons
 391 from a Transplantation of Zebra Mussels into a Small Urban River: An Integrated Ecotoxicological
 392 Assessment. *Environmental Toxicology*, 25, 468-478.
- 393 COLVILLE, A., JONES, P., PABLO, F., KRASSOI, F., HOSE, G. & LIM, R. 2008. Effects of chlorpyrifos on
 394 macroinvertebrate communities in coastal stream mesocosms. *Ecotoxicology*, 17, 173-180.
- 395 CONNELL, D. W. 1988. Bioaccumulation behavior of persistent organic-chemicals with aquatic organisms.
 396 *Reviews of Environmental Contamination and Toxicology*, 102, 117-154.
- 397 CUSTER, T. W., CUSTER, C. M., HINES, R. K. & SPARKS, D. W. 2000. Trace elements, organochlorines,
 398 polycyclic aromatic hydrocarbons, dioxins, and furans in lesser scaup wintering on the Indiana Harbor
 399 Canal. *Environmental Pollution*, 110, 469-482.
- 400 DE JONGE, M., BELPAIRE, C., GEERAERTS, C., DE COOMAN, W., BLUST, R. & BERVOETS, L. 2012.
 401 Ecological impact assessment of sediment remediation in a metal-contaminated lowland river using
 402 translocated zebra mussels and resident macroinvertebrates. *Environmental Pollution*, 171, 99-108.
- 403 DE JONGE, M., TIPPING, E., LOFTS, S., BERVOETS, L. & BLUST, R. 2013. The use of invertebrate body
 404 burdens to predict ecological effects of metal mixtures in mining-impacted waters. *Aquatic Toxicology*,
 405 142, 294-302.
- 406 DE PAUW, N. & VANNEVEL, R. 1991. Macroinvertebraten en waterkwaliteit. Determineersleutels van
 407 macroin-vertebraten en beoordelingsmethoden van de waterkwaliteit. *Stichting Leefmilieu*.
- 408 DEPAUW, N. & VANHOOREN, G. 1983. Method for biological quality assessment of watercourses in
 409 Belgium. *Hydrobiologia*, 100, 153-168.
- 410 EATON, H. J. & LYDY, M. J. 2000. Assessment of water quality in Wichita, Kansas, using an index of biotic
 411 integrity and analysis of bed sediment and fish tissue for organochlorine insecticides. *Archives of*
 412 *Environmental Contamination and Toxicology*, 39, 531-540.
- 413 EMANUELA, F., GIUSEPPE, S., STEFANO, G., MATTIA, B. & MAURO, M. 2017. Comparison of Lindane
 414 and Carbaryl Pesticide Bioaccumulation in the Common Sole (*Solea solea*). *Bulletin of Environmental*
 415 *Contamination and Toxicology*, 98, 656-661.
- 416 Flemish Government 2000. Besluit van de Vlaamse Regering van 1 juni 1995 houdende vastelling van
 417 het Vlaamse reglement betreffende de milieuvergunning (Vlarem), zoals gewijzigd bij besluit
 418 van 17 juli 2000. Belgisch Staatsblad, Brussel, België.
- 419 GABRIELS, W., LOCK, K., DE PAUW, N. & GOETHALS, P. L. M. 2010. Multimetric Macroinvertebrate
 420 Index Flanders (MMIF) for biological assessment of rivers and lakes in Flanders (Belgium).
 421 *Limnologica*, 40, 199-207.

- 422 KRAAK, M. H. S., TOUSSAINT, M., LAVY, D., DAVIDS, C. 1994. SHORT-TERM EFFECTS OF METALS
423 ON THE FILTRATION-RATE OF THE ZEBRA MUSSEL DREISSENA-POLYMORPHA.
424 *Environmental Pollution*, 84, 139-143.
- 425 LUOMA, S. N., CAIN, D. J. & RAINBOW, P. S. 2010. Calibrating biomonitors to ecological disturbance: a
426 new technique for explaining metal effects in natural waters. *Integr Environ Assess Manag*, 6(2), 199-
427 209.
- 428 MACCHI, P., LOEWY, R. M., LARES, B., LATINI, L., MONZA, L., GUINAZU, N. & MONTAGNA, C. M.
429 2018. The impact of pesticides on the macroinvertebrate community in the water channels of the Rio
430 Negro and Neuquen Valley, North Patagonia (Argentina). *Environmental Science and Pollution*
431 *Research*, 25, 10668-10678.
- 432 MAYON, N., BERTRAND, A., LEROY, D., MALBROUCK, C., MANDIKI, S. N. M., SILVESTRE, F.,
433 GOFFART, A., THOME, J. P. & KESTEMONT, P. 2006. Multiscale approach of fish responses to
434 different types of environmental contaminations: A case study. *Science of the Total Environment*, 367,
435 715-731.
- 436 MERSCH, J., WAGNER, P., PIHAN, J. C. 1996. Copper in indigenous and transplanted zebra mussels in
437 relation to changing water concentrations and body weight. *Environmental Toxicology and Chemistry*,
438 15, 886-893.
- 439 MICHIELS, E. D. G., VERGAUWEN, L., HAGENAARS, A., FRANSEN, E., VAN DONGEN, S., VAN
440 CRUCHTEN, S. J., BERVOETS, L. & KNAPEN, D. 2017. Evaluating Complex Mixtures in the
441 Zebrafish Embryo by Reconstituting Field Water Samples: A Metal Pollution Case Study. *International*
442 *Journal of Molecular Sciences*, 18.
- 443 MIRANDA, A. L., ROCHE, H., RANDI, M. A. F., MENEZES, M. L. & RIBEIRO, C. A. O. 2008.
444 Bioaccumulation of chlorinated pesticides and PCBs in the tropical freshwater fish *Hoplias*
445 *malabaricus*: Histopathological, physiological, and immunological findings. *Environment International*,
446 34, 939-949.
- 447 MUNZE, R., HANNEMANN, C., ORLINSKIY, P., GUNOLD, R., PASCHKE, A., FOIT, K., BECKER, J.,
448 KASKE, O., PAULSSON, E., PETERSON, M., JERNSTEDT, H., KREUGER, J., SCHUURMANN,
449 G. & LIESS, M. 2017. Pesticides from wastewater treatment plant effluents affect invertebrate
450 communities. *Science of the Total Environment*, 599, 387-399.
- 451 RAINBOW, P. S., HILDREW, A. G., SMITH, B. D., GEATCHES, T. & LUOMA, S. N. 2012. Caddisflies as
452 biomonitors identifying thresholds of toxic metal bioavailability that affect the stream benthos.
453 *Environmental Pollution*, 166, 196-207.
- 454 SCHAFER, R. B., VON DER OHE, P. C., RASMUSSEN, J., KEFFORD, B. J., BEKETOV, M. A., SCHULZ,
455 R. & LIESS, M. 2012. Thresholds for the Effects of Pesticides on Invertebrate Communities and Leaf
456 Breakdown in Stream Ecosystems. *Environmental Science & Technology*, 46, 5134-5142.
- 457 SZEKACS, A., MORTL, M. & DARVAS, B. 2015. Monitoring Pesticide Residues in Surface and Ground
458 Water in Hungary: Surveys in 1990-2015. *Journal of Chemistry*.
- 459 THIÈRE, G. & SCHULZ, R. 2004. Runoff-related agricultural impact in relation to macroinvertebrate
460 communities of the Lourens River, South Africa. *Water Research*, 38, 3092-3102.
- 461 VAN AEL, E., BELPAIRE, C., BREINE, J., GEERAERTS, C., VAN THUYNE, G., EULAERS, I., BLUST, R.
462 & BERVOETS, L. 2014. Are persistent organic pollutants and metals in eel muscle predictive for the
463 ecological water quality? *Environmental Pollution*, 186, 165-171.
- 464 VAN PRAET, N., DE BRUYN, L., DE JONGE, M., VANHAECKE, L., STOKS, R. & BERVOETS, L. 2014.
465 Can damselfly larvae (*Ischnura elegans*) be used as bioindicators of sublethal effects of environmental
466 contamination? *Aquatic Toxicology*, 154, 270-277.
- 467 VON DER OHE, P. C. & LIESS, M. 2004. Relative sensitivity distribution of aquatic invertebrates to organic
468 and metal compounds. *Environmental Toxicology and Chemistry*, 23, 150-156.
- 469 WEBER, G., CHRISTMANN, N., THIERY, A. C., MARTENS, D. & KUBINIÖK, J. 2018. Pesticides in
470 agricultural headwater streams in southwestern Germany and effects on macroinvertebrate populations.
471 *Science of the Total Environment*, 619, 638-648.
- 472 WILLE, K., KIEBOOMS, J. A. L., CLAESSENS, M., RAPPE, K., VANDEN BUSSCHE, J., NOPPE, H., VAN
473 PRAET, N., DE WULF, E., VAN CAETER, P., JANSSEN, C. R., DE BRABANDER, H. F. &
474 VANHAECKE, L. 2011. Development of analytical strategies using U-HPLC-MS/MS and LC-ToF-MS
475 for the quantification of micropollutants in marine organisms. *Analytical and Bioanalytical Chemistry*,
476 400, 1459-1472.

477

478 **Figure Legends**


479 Figure 1. Map of (A) sampling sites (1-9), (B) sampling sites (10-15). Sites 1 to 9 are pesticide
480 polluted sites and site 10 to 15 are metal polluted sites.

481 Figure 2. Detected pesticide concentrations in zebra mussels at pesticide polluted sites (1 to 9)
482 and metal polluted sites (10 to 15).


483 Figure 3. The relationship between accumulated pesticides in zebra mussels and biological
484 water quality indices (MMIF). The dashed line indicates the threshold value for the MMIF set
485 at a score of 0.6. Due to the wide range of chlorpyrifos concentration the values were log
486 transformed.

487 Figure 4. CCA diagram of macroinvertebrate taxa composition. Direct ordination based on both
488 accumulated pesticides and metals concentration. Axes represent the first two axes of the
489 ordination analysis. Eigenvalues (%cumulative variance): axis 1:0.677 (26.4%), axis 2:0.392
490 (41.6%), axis 3:0.348 (55.2%), axis 4: 0.248 (64.8%)

491




492 Figure 1. Map of (A) sampling sites (1-9), (B) sampling sites (10-15). Sites 1 to 9 are pesticide polluted
 493 sites and site 10 to 15 are metal polluted sites.



494


495 Figure 2. Detected pesticide concentrations in zebra mussels at pesticide polluted sites (1 to 9)
 496 and metal polluted sites (10 to 15).

497



498

499 Figure 3. The relationship between accumulated pesticides in zebra mussels and biological water quality indices (MMIF). The dashed line
 500 indicates the threshold value for the MMIF set at a score of 0.6. Due to the wide range of chlorpyrifos concentration the values were log
 501 transformed.



502
 503 Figure 4. CCA diagram of macroinvertebrate taxa composition. Direct ordination based on
 504 both accumulated pesticides and metals concentration. Axes represent the first two axes of the
 505 ordination analysis. Eigenvalues (%cumulative variance): axis 1:0.677 (26.4%), axis 2:0.392
 506 (41.6%), axis 3:0.348 (55.2%), axis 4: 0.248 (64.8%)
 507

508 **Table Legends**

509 Table 1. Location, General water quality characteristic and Multimetric Macroinvertebrate
510 Index Flanders (MMIF) of the sampling sites

511 Table 2. The 95-percentile threshold concentrations (ng/g dry weight) in zebra mussel above
512 which ecological status was always low (MMIF < 0.6)

513

514 Table 1. Location, General water quality characteristics and Multimetric Macroinvertebrate
 515 Index Flanders (MMIF) of the sampling sites

Site	Name	VMMnr*	T (°C)	PH	O2 (mg/L)	O2 (%)	Conductivity (µS/cm)	MMIF
1	Demer	401000	10	7.43	9.21	86.7	506	0.7
2	Terbermenbeek	449800	8	7.06	5.09	44.8	411	0.2
3	Hoevenbeek	449750	8	7.26	5.3	46.3	545	0.15
4	Molenbeek	436920	9.7	6.82	6.94	63.6	882	0.15
5	Zevenbronnenbeek	445250	8.2	7.63	8.52	75.6	892	0.5
6	Kleine Vondelbeek	426910	9.3	7.54	9.32	80.7	733	0.5
7	Bovenheidebeek	483320	7.4	7.72	9.66	79.7	692	0.6
8	Bierbeek	483300	9	7.57	8.95	77	673	0.5
9	Paddenpoel	487200	8.8	7.55	9.74	83.1	450	0.7
10	Scheppelijke Nete	333750	NA	7.54	9.49	78	444	0.7
11	Scheppelijke Nete	333600	5.4	7.3	10.9	90	461	0.5
12	Scheppelijke Nete	333500	5.7	7.37	10.7	88	416	0.15
13	Scheppelijke Nete	NA	5.6	7.46	11.3	92	408	0.3
14	Molse Nete1	333100	5.7	7.39	10.2	84	378	0.5
15	Molse Nete2	330200	6.1	7.35	9.49	60	394	0.4
16	Scheldt	179000	NA	NA	NA	NA	NA	0.25
17	Terneuzen	NA	NA	NA	NA	NA	NA	0.25
Rf1	Blaarmeersen	NA	NA	NA	NA	NA	NA	NA

516 *VMM (Vlaamse Milieumaatschappij) is the Flemish Environment Agency which monitors (both chemically and
 517 biologically) Flemish surface waters and sediment on a regular basis

518

519 Table 2. The 95-percentile threshold concentrations (ng/g dry weight) in zebra mussel above
520 which ecological status was always low (MMIF < 0.6)

Compound	Threshold
Chlorpyrifos	8
Terbutylazine	2.08
Dimethoate	2

521

522 Supplementary data

523 SI. 1 Pesticide body burdens in zebra mussel tissue (ng/g dry weight)

Site	Dimethoate	Chloridazon	Pirimicarb	Isoproturon	Dichlorvos	Atrazine	Diuron	Linuron	Terbuthylazine	Metolachlor	Diazinon	Chlorpyrifos
1	2.20	<LOQ	1.88	<LOQ	83	<LOQ	<LOQ	<LOQ	1.02	<LOQ	<LOQ	3.05
2	1.27	<LOQ	2.27	8.24	95	<LOQ	2.23	<LOQ	7.51	2.63	1.06	2203
3	3.80	8.82	2.36	0.15	116	<LOQ	<LOQ	1.64	1	5	<LOQ	103
4	10.1	3.77	1.97	<LOQ	70	<LOQ	<LOQ	<LOQ	2.29	<LOQ	3.08	45
5	3.98	6.78	1.24	0.16	<LOQ	2.60	<LOQ	<LOQ	1.64	<LOQ	1.45	6.54
6	<LOQ	<LOQ	1.77	<LOQ	85	2.55	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	1.79
7	2.69	<LOQ	1.93	<LOQ	128	<LOQ	<LOQ	<LOQ	2.32	<LOQ	<LOQ	9.25
8	5.03	<LOQ	1.45	<LOQ	36	<LOQ	<LOQ	<LOQ	1.42	<LOQ	2.90	1.94
9	<LOQ	<LOQ	1.24	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
10	<LOQ	<LOQ	1.24	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	1.45	<LOQ	<LOQ
12	<LOQ	3.44	2.01	<LOQ	<LOQ	4.76	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	1.26
13	1.92	<LOQ	4.29	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	1.09	<LOQ	<LOQ	<LOQ
14	<LOQ	<LOQ	1.24	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
15	1.35	<LOQ	1.24	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	1.08
16	1.44	<LOQ	1.72	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
17	<LOQ	<LOQ	1.29	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
Start	1.50	<LOQ	5.48	<LOQ	<LOQ	3.31	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ

525 SI. 2 Summary of macroinvertebrate taxa and abundance. The number represent % of each
 526 species per sampling site

Order	Macroinvertebrate	Sampling sites															
	Taxa	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Oligochaeta	<i>Haplotaxidae</i>													18	14	5	
	<i>Lumbriculidae</i>	13	31			9								4		10	
	<i>Tubificidae</i>	20		33		14		1			6						
Hirudinea	<i>Erpobdella</i>	1				1										2	1
	<i>Glossiphonia.sp</i>		31										1		1		
	<i>Haementeria.sp</i>																1
	<i>Helobdella.sp</i>	1				1											
	<i>Hemiclepsis.sp</i>															7	10
Mollusca	<i>Lymnaea.sp</i>					6										2	2
	<i>Physa.sp</i>	1				1											
	<i>Planorbis.sp</i>							1									
	<i>Sphaerium.sp</i>	2			1	1											
Acari	<i>Hydracarina.sp</i>					31	67	5		1	47						
Crustacea	<i>Asellidae</i>	22	38			15			11			10	3	24	56	34	
	<i>Gammaridae</i>				85	1	1	78	77	95	21	42					
Diptera	<i>Ceratopogonidae</i>															1	
	<i>Chironomidae</i>			33									95	53	9	36	
	<i>Limoniidae</i>								1			24	1				
	<i>Ptychopteridae</i>								4	1	21						
	<i>Simuliidae</i>	25		33	8	17	28	10	9	1							
	<i>Tipulidae</i>				1		1									1	1
Megaloptera	<i>Sialis.sp</i>									1			1				
Coleoptera	<i>Dytiscidae</i>						2									2	
	<i>Gyrinidae</i>												2				
	<i>Scirtidae</i>											1					
Odonata	<i>Calopteryx.sp</i>	2				2						4		3	1		
Ephemeroptera	<i>Ephemera.sp</i>												8				
	<i>Ephemerella.sp.</i>	2				1	1										
Trichoptera	<i>Glossosomatidae</i>												5		2	1	
	<i>Hydropsychidae</i>					1										1	
	<i>Leptoceridae</i>	7				5											
	<i>Limnephilidae</i>							1	1		2						
	<i>Phryganeidae</i>											5			1		
	<i>Polycentropodidae</i>								1	1							
	<i>Rhyacophilidae</i>															1	

527

528

529 SI. 3 Concentration of metals accumulated in zebra mussel ($\mu\text{g/g}$ dry weight) at sampling site

Site	Ag	Cd	Pb	Cr	Mn	Co	Ni	Cu	Zn
1	0.017	0.570	0.350	0.001	7.96	1.05	10.4	5.04	22.2
2	0.016	7.150	2.090	0.016	59.1	7.08	72.6	76.2	140
3	0.004	0.460	0.340	0.002	12.9	1.05	9.95	5.42	185
4	0.002	0.550	0.430	0.360	12.0	0.910	10.1	4.63	136
5	0.003	0.460	0.300	0.002	11.6	0.890	9.18	4.74	170
6	0.001	0.610	0.280	0.180	12.2	1.05	9.36	5.25	175
7	0.001	0.660	0.860	0.041	20.2	1.04	10.2	6.64	225
8	0.002	0.590	0.400	0.048	12.4	0.950	10.2	5.19	182
9	0.016	0.710	0.200	0.270	10.5	0.790	8.68	5.56	228
10	0.034	0.670	2.19	0.002	28.5	1.16	8.52	4.71	308
12	0.011	4.61	5.40	0.370	21.9	0.340	21.1	13.1	2843
13	0.003	3.09	2.48	0.230	18.2	1.77	21.6	5.23	541
14	0.030	2.76	4.54	0.290	16.0	1.58	13.4	4.50	355
15	0.011	1.90	2.66	0.200	13.7	1.12	6.04	3.95	266
16	ND	0.420	0.560	0.890	ND	0.950	3.71	8.40	84.3
17	ND	8.170	0.860	1.680	ND	0.960	4.14	7.49	63.4
Start	ND	0.610	0.160	1.360	ND	1.19	1.99	8.78	85.3

530

531 SI. 4 The octanol/water partition coefficient (K_{ow}) of detected pesticides

Pesticides	Log K_{ow}	Reference
Dimethoate	0.7	Masià.A 2013
Chloridazon	3.89	Roberts.T.R 1998
Pirimicarb	3.4	Masià.A 2013
Isoproturon	2.5	Masià.A 2013
Dichlorvos	1.9	Roberts.T.R 1998
Atrazine	2.7	Masià.A 2013
Diuron	2.8	Masià.A 2013
Linuron	3	Roberts.T.R 1998
Terbuthylazine	2.3	Masià.A 2013
Metolachlor	3.4	Masià.A 2013
Diazinon	3.69	Masià.A 2013
Chlorpyrifos	4	Masià.A 2013

532