

**This item is the archived peer-reviewed author-version of:**

Priority-setting and personality : effects of dispositional optimism on preferences for allocating healthcare resources

**Reference:**

Luyten Jeroen, Kessels Roselinde, Desmet Pieter, Goos Peter, Beutels Philippe.- Priority-setting and personality : effects of dispositional optimism on preferences for allocating healthcare resources  
Social justice research - ISSN 0885-7466 - 32:2(2019), p. 186-207  
Full text (Publisher's DOI): <https://doi.org/10.1007/S11211-019-00329-5>  
To cite this reference: <https://hdl.handle.net/10067/1582020151162165141>

# **Priority-setting and personality: Effects of dispositional optimism on preferences for allocating healthcare resources**

**Jeroen Luyten<sup>1,\*</sup>, Roselinde Kessels<sup>2,3</sup>, Pieter Desmet<sup>4</sup>, Peter Goos<sup>2,5</sup>, Philippe Beutels<sup>6,7</sup>**

<sup>1</sup>Leuven Institute for Healthcare Policy, KU Leuven, Kapucijnenvoer 35, 3000 Leuven, Belgium

<sup>2</sup>Faculty of Business and Economics, University of Antwerp, Prinsstraat 13, 2000 Antwerp, Belgium

<sup>3</sup>School of Economics, University of Amsterdam, PO Box 15867, 1001 NJ Amsterdam, The Netherlands

<sup>4</sup>Rotterdam Institute for Law and Economics, Erasmus University Rotterdam, Burgemeester Oudlaan 50, Postbus 1738, 3000 DR Rotterdam, The Netherlands

<sup>5</sup>Department of Biosystems, Faculty of Bioscience Engineering, KU Leuven, Kasteelpark Arenberg 30, PO Box 2456, 3001 Heverlee, Belgium

<sup>6</sup>Centre for Health Economics Research and Modelling Infectious Diseases (CHERMID), Vaccine and Infectious Disease Institute (VAXINFECTIO), Faculty of Medicine and Health Sciences, University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Belgium

<sup>7</sup>School of Public Health and Community Medicine, The University of New South Wales, Sydney, Australia

\*Corresponding author: jeroen.luyten@kuleuven.be

Please cite this article as:

Luyten Jeroen, Kessels Roselinde, Desmet Pieter, Goos Peter, Beutels Philippe (2019). Priority-setting and personality: Effects of dispositional optimism on preferences for allocating healthcare resources, *Social Justice Research*, 1-22. <https://doi.org/10.1007/s11211-019-00329-5>

## **Abstract**

In a publicly financed health system, it is important that priority-setting reflects social values. Many studies investigate public preferences through surveys taken from samples, but to be representative, these samples must reflect value judgments of all relevant population subgroups. In this study, we explore whether, next to better-understood sources of heterogeneity such as age, education or gender, also differences in personality play a role in how people want to set limits to healthcare. We investigate the influence of dispositional optimism: whether someone anticipates a good or bad future. This is an important personality dimension that has been shown to widely reverberate into people's lives and that can also be expected to influence people's views on healthcare. To test our hypothesis, we asked a representative sample of the Belgian population ( $N=750$ ) to complete both the Revised Life Orientation Test (LOT-R) and a discrete choice experiment about allocating healthcare resources, and we investigated the relationships between both measurements. We found that more pessimistic individuals were less supportive of using patients' age as a selection criterion and more hesitant to invest in prevention. Since individual dispositions are usually not part of the criteria for selecting representative samples, our findings point at a potential non-response bias in studies that elicit social values.

**Keywords** Resource allocation, Preferences, Equity, Prevention, Fair innings, Responsibility, Optimism, Pessimism

## 1. Introduction

Health policy decision-makers with limited budgets unavoidably have to set priorities over different healthcare interventions (Daniels and Sabin 2008). These decisions imply several complex trade-offs. For instance, deeply grained intuitions to ‘rescue’ individual, identifiable patients need to be reconciled with more calculated, utilitarian approaches that focus on cost-effectiveness, maximizing health at the population level. A judgment needs to be made about which illness is more severe, e.g. comparing physical with mental illness. A decision is needed about whether there is a role for personal characteristics of patients such as how old a patient is or whether she has taken good care of her health in the past (Olsen, Richardson et al. 2003). A balance must be struck between investing resources in prevention versus in treatment (Faust and Menzel 2012). The common element in these questions is that they require value judgments. There is no obvious right or wrong way to make these decisions, and reasonable people can disagree in their answers.

Therefore, many think that there is an important role for the preferences and values of the population in setting limits to public healthcare. An extensive body of research has emerged describing how people make trade-offs and set priorities [for reviews, see (Schwappach 2002, Dolan, Shaw et al. 2005, Shah 2009, Gu, Lancsar et al. 2015)]. In the UK, the National Institute for Health and Care Excellence (NICE) regularly organizes “Citizens Councils” to formulate policy guidance with respect to social values in healthcare (NICE 2008). In Belgium, the Federal Healthcare Knowledge Centre recruited a large group of volunteers to develop ‘citizen labs’ to answer the dilemmas of healthcare rationing (Cleemput 2014, Standaard 2014). Similar examples can be found for many other countries, including Canada, the Netherlands or Australia (Mooney and Blackwell 2004, OMHLTC 2013). The values that emerge from these studies are increasingly used to assist decision-makers in priority-setting, e.g. in reimbursement decisions. They can be used in a deliberative approach, considering evidence on social values next to other pieces of information (such as medical or economic evidence) or in more algorithmic approaches where social value judgments are quantified, e.g. in the form of equity weights or inequality aversion estimates, which can explicitly be incorporated into decision models (e.g. in a distributive cost-effectiveness analysis) [see e.g. (Lancsar, Wildman et al. 2011, Asaria, Griffin et al. 2016, Cookson, Ali et al. 2018)].

The validity of these studies claiming to elicit the social values of a population depends on the representativeness of the samples used in the study. Usually, sample representativeness is assessed in terms of age, gender, educational attainment, geographical spread and perhaps a few other socio-demographic variables. However, also differences in less visible characteristics can be important. One of these is someone’s personality. The role of personality traits in shaping how people think, feel and

behave has long been recognized in the field of psychology (Corr and Matthews 2009). Personality traits have been linked to a substantial series of important life outcomes, including marital status, occupational attainment and even mortality [e.g. (Roberts, Kuncel et al. 2007)]. If personality influences how someone makes the value judgments inherent to priority-setting, then this has implications for the representativeness of studies that infer social values from (often online) samples of volunteers. These studies might attract particular personality types, leading to social values that are not necessarily representative of a population. Indeed, research has shown that particular personality types are more likely to participate in survey research than others, leading to significant non-response bias (Marcus and Schutz 2005, Smith, Edens et al. 2012). For instance, one study found that non-responders to survey research were less agreeable, less extravert, had a lower openness to experience and lower narcissism than volunteers in both self- and observer ratings (Marcus and Schutz 2005).

In this study, we explore the effect on preferences for priority-setting of one well-known personality difference between individuals: whether someone generally expects to have a good or a bad future. This difference is coined ‘dispositional optimism’ in the psychology literature and there are strong reasons to believe that it matters to people’s views on healthcare (see Section 2). We added an established instrument to measure dispositional optimism, the Revised Life Orientation Test (LOT-R), to a discrete choice experiment (DCE) about priority-setting that was executed in a sample ( $N=750$ ) of the general population in Belgium. Section 2 summarizes the literature on dispositional optimism and suggests ways in which this personality dimension can influence preferences for healthcare priority-setting. Section 3 describes the methods used. Section 4 presents the results and Section 5 assesses the relevance of our findings and provides further points for discussion.

## **2. Dispositional optimism**

The categorization of individuals in optimists and pessimists is deeply rooted in popular culture and folk psychology (e.g. the glass half-full or half-empty metaphor) (Carver and Scheier 2014). In previous decades, it has become the subject of extensive scientific research, mainly through the development of the Life Orientation Test (LOT), a self-report measure of optimism and pessimism, which was revised in 1994 (LOT-R) (Scheier and Carver 1985, Scheier, Carver et al. 1994). This body of research resolves that ‘dispositional optimism’ is a personality trait, which remains relatively stable over one’s lifespan, provided it is not substantially manipulated or exposed to disruptive life changes (Carver, Scheier et al. 2010). There is converging evidence that optimism has a neurobiological basis, with pessimistic and optimistic views being primarily determined by higher activity in the right and the left cerebral hemisphere (Hecht 2013). Moreover, studies that investigate the genetic origins of dispositional

optimism find heritability estimates of 20-30% (Mosing, Gordon et al. 2009, Rius-Ottenheim, Kromhout et al. 2012a-b). However, nurture matters too with for instance higher socio-economic status in childhood being predictive of a more optimistic nature in adulthood (Heinonen, Raikkonen et al. 2006).

There is ample evidence that dispositional optimism strongly affects our mental, physical, economic and social state (Carver, Scheier et al. 2010, Hecht 2013). Optimism is associated with increased protection against stroke (Kim, Park et al. 2011), carotid artery blockage (Matthews, Raikkonen et al. 2004) and rehospitalization after coronary artery bypass grafting (Tindle, Chang et al. 2009) and yields better immune responses (Segerstrom 2007). One study followed a cohort of 95,000 healthy women and found that more optimistic individuals are less likely to develop coronary heart disease (CHD) or die from CHD-related causes or any cause over an 8-year period (Tindle, Chang et al. 2009). Optimists are also less likely to smoke, more likely to exercise and have more healthy diets (Carver and Scheier 2014). Furthermore, outside of the health domain, optimism is also associated with better socio-economic outcomes. More optimistic students (measured before starting higher education) have lower dropout rates in their college years (Nes, Evans et al. 2009) and later earn more than their less optimistic counterparts (Segerstrom 2007). Also, optimists indicate greater satisfaction in their romantic relationships, and so do their partners (Srivastava, McGonigal et al. 2006). They have a broader social network (Andersson 2012) and are more resilient towards developing loneliness late in life (Rius-Ottenheim, Kromhout et al. 2012a-b). They even invest in different stock portfolios (Puri and Robinson 2007) and make different financial and accounting decisions as managers (Heaton 2002). One study suggests that optimists are also more vulnerable to problematic gambling behaviour (Gibson and Sanbonmatsu 2004). Issues of reverse causation (that success determines optimism and not vice versa) and unmeasured confounding (that both optimism and success are caused by an omitted variable) cannot be fully ruled out in these studies due to their observational nature and often relatively short follow-up periods. However, many studies use extensive strategies to mitigate these issues and are confident that the causal relationship runs (at least partly) from optimism to the variable under study.

An explanation for the more positive outcomes associated with optimism is that optimists and pessimists are fundamentally different in their problem-solving attitude (Carver, Scheier et al. 2010, Hecht 2013). Optimism is shown to be associated with more efficient scanning for risks, more effective coping with adversity and taking a more proactive approach in confronting possible problems. The mechanism behind this difference in attitude, it is argued, operates mainly through differences between optimists and pessimists in their motivation for reaching objectives (Carver and Scheier 2014). Optimists are more confident that goals (small or big ones) are achievable, see fewer impediments and therefore do more effort, whereas pessimists easier disengage. This attitudinal difference reverberates into different degrees of dedication to goals, more experience in reaching these goals, which will in turn reinforce motivation and become a source of support in achieving new goals or coping with adversity. Moreover,

research shows that in the simultaneous pursuit of multiple goals, where trade-offs in investing effort are needed, optimists also do better in allocating more effort to high-priority goals and disengaging from either low-priority goals or goals with unfavourable odds (Segerstrom 2007, Geers, Wellman et al. 2009, Geers, Wellman et al. 2010, Pavlova and Silbereisen 2013). There is also a relationship between optimism and time preference. Although both are related concepts, optimism deals with someone's valuing of positive or negative future events, whereas time preference readjusts this value for distance in time with events further ahead receiving a lower value than more proximate events. Research has investigated the link between these two concepts and has found that higher levels of optimism are associated with higher discounting for time preference (Berndsen and van der Pligt 2001). The underlying mechanism, it is argued, is that optimists are keener on immediate gains because they are optimistic that these gains will be followed by additional gains, whereas they are keener to delay losses because they are more confident that these will be avoidable in the future.

Based on the literature described above, we hypothesized that dispositional optimism can also be an important driver of how people approach the problem of setting limits to healthcare. More specifically, we expect dispositional optimism to influence 1) the extent to which individuals value prevention over treatment, 2) the extent to which they take into account the known effectiveness of an intervention, 3) their valuation of health gains occurring in distant versus proximal futures (time preference) and 4) the extent to which they favour age-based rationing. First, the more proactive problem-solving attitude of optimists (Carver, Scheier et al. 2010, Hecht 2013) suggests that optimists will prefer higher investments in prevention and in avoiding bad outcomes in the future rather than paying for a wait-and-treat scenario later on. Observations from health behaviour research also show that pessimists are less prevention-oriented when it comes to their own health behaviours. Studies have documented this in the context of, e.g., preventing heart attacks (Radcliffe and Klein 2002), taking vitamins, eating low-fat foods and enrolment in a cardiac rehabilitation program after a bypass grafting (Scheier, Matthews et al. 1999) and sexual risk-taking behaviour (Taylor, Kemeny et al. 1992). Second, optimists' easier disengagement from goals with unfavourable odds (Geers, Wellman et al. 2009) suggests that optimists will set lower priorities for less effective healthcare, or for healthcare of which the health benefits are more uncertain. Third, the higher levels of time discounting of optimists (Berndsen and van der Pligt 2001) suggest that optimists will set higher priorities for healthcare with more immediate benefits. And, finally, as optimists, by definition, expect a better future, they may therefore also anticipate a lower need of healthcare at older age. Research indeed shows that at older age pessimists have worse health outcomes than optimists (Carver, Scheier et al. 2010, Carver and Scheier 2014) and fewer economic and social resources: lower incomes, a smaller social network, fewer close relationships and they perceive less social support from their partners, relatives and friends (Brissette, Scheier et al. 2002, Carver, Lehman et al. 2003, Macleod and Conway 2005), all of which are important sources of health and well-being at

older age [e.g. (Giles, Glonek et al. 2005, Sirven and Debrand 2012)]. This might affect their views on age-based priority-setting.

To test these hypotheses, we conducted a large-scale discrete choice experiment among a sample of the Belgian population. In the next sections, we summarize our study and the answers it gave regarding those hypotheses.

### 3. Methods

#### 3.1 Sample

A sample from an online panel of 10,753 Belgians was recruited via a market research company of which a total of 3,160 individuals (30%) agreed to participate in our study. From this group, 750 respondents were retained, by random filling of predetermined quota for age, gender, province, rural versus urban spread and level of educational attainment. Only participants aged 18-75 years were included. **Table 1** summarizes the sample's representativeness relative to the Belgian population. All these respondents provided answers to the LOT-R and DCE questions.

**Table 1** Characteristics of the sample relative to those of the Belgian population. Source Belgian Data: Federale Overheidsdienst Economie (FOD-economie 2012)

	<b>Sample (%)</b>	<b>Belgian population (%)</b>
<b>Language</b>		
Dutch	56	56
French	44	44
<b>Gender</b>		
Male (M)	50	50
Female (F)	50	50
<b>Gender per age group<sup>a</sup></b>		
18 – 25 M	6	6
18 – 25 F	6	6
25 – 34 M	9	9
25 – 34 F	10	9
35 – 44 M	10	11
35 – 44 F	10	10
45 – 54 M	10	10
45 – 54 F	11	10
55 – 64 M	9	8
55 – 64 F	10	8
65 – 74 M	6	6
65 – 74 F	4	6
<b>Level of education<sup>b</sup></b>		
None or primary	8	19
Lower secondary	10	20
Higher secondary	31	33
Higher non-university	35	18



(Post-)university	15	10
<b>Province</b>		
Antwerp	15	16
West Flanders	10	11
East Flanders	13	13
Limburg	8	8
Hainault	13	12
Liege	10	10
Luxemburg	3	2
Namur	5	4
Brussels	10	10
Flemish Brabant	11	10
Walloon Brabant	3	3
<b>Smoking status<sup>c</sup></b>		
Never smoked	45	54
Ex-smoker	30	22
Smoker	25	25

<sup>a</sup>Age: the percentages reported are proportions in the selected population (18-75), representing 71% of the total Belgian population

<sup>b</sup>Education: the percentages reported for the Belgian population are for the age group 15 years or older. The percentages for our sample are only for the age group between 18 and 75 years. The overrepresentation of higher educated respondents in our sample as compared to the total population can be explained by our exclusion of the age group between 15 and 18 years that is too young for higher education, and the age group 75 years or older for which higher education was less democratically accessible

<sup>c</sup>Smoking percentages from the population are based upon the study (SIPH, 2008) and are representative of the population aged 15 years or older

### 3.2 The Revised Life Orientation Test (LOT-R)

The standard instrument to measure dispositional optimism is the Revised Life Orientation Test (LOT-R) (Scheier, Carver et al. 1994). This is a revised version of the earlier Life Orientation Test (Scheier and Carver 1985) and focuses more on the conceptual core of the trait (i.e. expectations about one's future). The scale consists of ten items, three of which measure optimism, three measure pessimism and four of which are filler items to disguise the underlying purpose of the test (see **Figure 1**). Respondents have to answer on a Likert scale (in our case, a five-point scale), ranging from 'strongly disagree' to 'strongly agree'. Numerous studies have documented the reliability and the validity of the scale, reporting adequate measures of internal consistency, test-retest reliability and construct and predictive validity [for a review, see (Carver, Scheier et al. 2010)]. In empirical studies, test-retest correlations range from 0.58 to 0.79 over periods ranging from several weeks, years to more than a decade (Carver, Scheier et al. 2010).

1. In uncertain times, I usually expect the best.
[2. It's easy for me to relax.]
3. If something can go wrong for me, it will.
4. I'm always optimistic about my future.
[5. I enjoy my friends a lot.]
[6. It's important for me to keep busy.]
7. I hardly ever expect things to go my way.
[8. I don't get upset too easily.]
9. I rarely count on good things happening to me.
10. Overall, I expect more good things to happen to me than bad.

Note: Items 2, 5, 6 and 8 are fillers.

**Figure 1** The Revised Life Orientation Test (LOT-R)

There is discussion in the literature about whether optimism and pessimism are two polar opposites on a one-dimensional continuum (RobinsonWhelen, Kim et al. 1997, Rauch, Schweizer et al. 2007, Segerstrom, Evans et al. 2011, Chiesi, Galli et al. 2013) or whether both are two separable dimensions, one pertaining to affirmation versus disavowal of optimism and the other to affirmation or disavowal of pessimism (Glaesmer, Rief et al. 2012). Although virtually all bipolar trait scales that contain both positively and negatively framed items typically form two dimensions in factor analyses, for some researchers these statistical grounds are enough to also treat optimism and pessimism as two conceptually different dimensions (Carver, Scheier et al. 2010, Carver and Scheier 2014). Others think the split is a product of method variance in responding and question the conceptual possibility of people to be pessimistic and optimistic at the same time. Studies aimed at settling the issue reached opposite conclusions: some claim a unidimensional view is best (RobinsonWhelen, Kim et al. 1997, Rauch, Schweizer et al. 2007, Segerstrom, Evans et al. 2011, Chiesi, Galli et al. 2013), others claim the two dimensions should be treated separately (Glaesmer, Rief et al. 2012). To accommodate for these different views and in line with recommendations in the literature (Carver, Scheier et al. 2010), we used three different LOT-R outcomes in our analyses: a total LOT-R score (the one-dimensional construct, termed LO henceforth) based on the three optimistic and the three pessimistic items, and two separate constructs: an optimism score (OPT) based only on the three optimistic items and a pessimism score (PES) based on the three pessimistic items. To obtain the LO and OPT scores, we summed the scores of the items under consideration. To obtain the PES score, we summed the scores of the pessimistic items and reversed the scaling so that a high PES score stands for a more pessimistic attitude.

### 3.3 Discrete choice experiment on priority-setting

DCEs are a widely used survey method to quantify individuals' preferences (Ryan, Gerard et al. 2008). Participants are presented with a series of choices, usually between two goods described by the same attributes but differing in their attribute levels. By observing respondents' preferred choices, researchers can infer how the value of the competing options is determined by the attributes of the product. We carried out a DCE in which all 750 respondents had to complete 14 choice sets consisting of two

competing healthcare interventions of which they were told that only one could be subsidized (completely) by the government. We described the health programs in terms of seven attributes: effectiveness of the program, severity of the illness, when health benefits are expected to occur (timing), possibility of adverse effects from the intervention, age group of patients, link between disease and patients' lifestyle and the curative or preventive nature of the program (see the first column in **Table 3**). In all other respects (e.g. costs), both programs were equal. These seven attributes were chosen because they represent salient, generic dimensions of healthcare, comprising a wide range of possible healthcare programs. The attributes that are most relevant to our hypotheses were effectiveness, timing, curative or preventive type and patients' age group. Our expectation was that respondents differing in their LOT-R score will attach a different importance to these attributes in their choices which healthcare program to subsidize.

**Appendix A** shows one of the 42 choice sets that we constructed (14 per respondent, 3 survey versions, in total 10,500 choice observations (14\*750)). The full descriptive results of the DCE and its conclusions for all attributes regardless of LOT-R are summarized elsewhere (Luyten, Kessels et al. 2015).

### **3.4 Other variables in the survey**

Background information collected from respondents included their age, sex, height, weight (to calculate the Body Mass Index), province, language, professional group, educational attainment, household size, age of youngest family member (indicating whether respondents have children as well as their age), experience as healthcare worker, smoking status and experience with severe illness (personal or within the family). Respondents also provided self-assessments of their health through a standardized Health-Related Quality of Life instrument, the EQ-5D-5L (Herdman, Gudex et al. 2011).

### **3.5 Statistical analysis**


We assessed the internal consistency of the LOT-R using Cronbach's alphas and described the correlation between the LO, OPT and PES scores using Pearson's correlation coefficients. We constructed multiple linear regression equations to identify significant associations between LO, OPT and PES and the other respondent characteristics surveyed (see Section 4.1). We analysed the relationship between the DCE data and the three LOT-R scores using a multinomial logit model (MNL), also called McFadden's conditional logit model (McFadden 1974) (see Section 4.2). This model allows assessing the relative weight of each of the seven attributes in predicting a choice, but also, by adding an interaction term – for instance with LO – it allows assessing whether respondents who differ in LO also differ in their valuation of the attributes. To control for confounding via other respondent variables

associated with LO, OPT and PES, we also added interactions with these other variables in the model. In **Appendix B**, we provide further information about the MNL model.

## 4. Results

### 4.1 Psychometric properties of and variables associated with LOT-R

**Figure 2** summarizes the LO, PES and OPT variables using boxplots. LO values ranged from 8 to 30 and PES and OPT values from 3 to 15. Our sample had an average (and median) life orientation score of 20. Respondents scored higher on the OPT than on the PES scale. The average (and median) optimism score was 11; the average (and median) pessimism score was 9. The internal consistency of the three variables was acceptable [Cronbach's  $\alpha = 0.75$  (LO), 0.78 (PES) and 0.72 (OPT)]. A factor analysis clearly indicated OPT and PES as two unique factors (describing about 67% of the variation). The OPT score correlated weakly with the three pessimistic items (between 0.24 and 0.30), and the PES score correlated weakly with the optimistic items (between -0.36 and -0.17).



**Figure 2** Boxplots of LO, PES and OPT scores ( $N=750$ )

Results from multiple linear regression analysis (see **Table 2**) revealed that five variables were positively associated with a higher LO score: a higher EQ-5D-5L score, being older, non-smoker, a university degree and having a relatively younger-aged youngest household member. Higher OPT scores (optimistic items only) are associated with higher EQ-5D-5L scores, older age and having a younger-aged youngest household member. Higher PES scores are associated with lower EQ-5D-5L scores, younger age, smoking and having no or only a lower secondary education degree.

**Table 2** Variables associated with an individual's LOT-R score

	LO		OPT		PES	
	Estimate	P-value	Estimate	P-value	Estimate	P-value
EQ-5D-5L score	5.7778	<0.0001	3.3713	<0.0001	-2.5052	<0.0001
Respondent age	0.0485	<0.0001	0.0286	<0.0001	-0.0138	0.0096
Smoking status						
Yes	-0.4975	0.0005	NS	NS	0.4593	<0.0001
No	0.4975	0.0005	NS	NS	-0.4593	<0.0001
Level of education						
None or lower secondary education	-0.6833	0.0062	NS	NS	0.5572	0.0005
Higher secondary education	-0.1865	0.3529	NS	NS	0.1791	0.1623
Higher non-university degree	0.0752	0.6997	NS	NS	-0.1374	0.2685
(Post-)university degree	0.7946	0.0022	NS	NS	-0.5989	0.0003
Age youngest household member	-0.0157	0.0276	-0.0095	0.0272	NS	NS


NS means that the variable is non-significant at the 5% level

Variables that we also investigated but were found to be non-significant at the 5% level are the following: gender, province, occupational status, language, experience as healthcare worker, personal experience with severe illness, experience with severe illness in the family, respondent's body length and respondent's weight

#### 4.2 Associations between dispositional optimism and preferences for priority-setting

We estimated three models, shown in **Table 3**. Model I is the basic model of the DCE analysis that best describes the average preference of the sample, quantifying the extent to which the utility of an intervention (or, in other words, the respondent's choice) depends on the seven attributes of the intervention [see (Luyten, Kessels et al. 2015) for further discussion]. As can be inferred from the largest estimates (in magnitude) of the levels of each attribute, the average respondent's choice can mainly be predicted on the basis of the patient's age and lifestyle, followed by concerns about effectiveness and severity of illness. Models II and III focus on the link between these priority-setting preferences and respondents' LOT-R scores. Introducing the LO variable in model II in interaction with each of the seven attributes, we found that it interacts significantly with two attributes: 'patient's age' ( $p=0.07$ ) and 'type of intervention' ( $p=0.01$ ). However, using the OPT and PES subscales instead of the complete LO scale revealed that the OPT variable does not interact with any of the attributes, whereas the PES variable interacts with the same attributes 'patient's age' ( $p<0.01$ ) and 'type of intervention' ( $p<0.01$ ) with which LO interacts (see Model III). Because OPT and PES were identified as only weakly correlated clusters of which only PES interacted with the DCE attributes, we focus the rest of our analysis on PES instead of LO. However, this does not necessarily mean that we only investigate pessimism as the separation between optimism and pessimism may be a mere measurement artefact. The interactions with PES

remained significant when we controlled for the effect of the respondent variables that are associated with PES (EQ-5D-5L, age, education and smoking status), indicating an independent relationship between pessimism and preferences for age-based priority-setting and prevention versus cure. As indicated by the likelihood ratio (LR) test statistics in **Table 3**, demonstrating the predictive power of the attributes and attributes' interactions, we see that the interactions with PES are substantial. The predictive power of the PES interactions was 12% for PES\*age (LR Chi-Square=21) and 5% for PES\*type (LR Chi-Square=10) as compared to the predictive power of the lifestyle attribute, i.e. the attribute with highest predictive power in Model III (LR Chi-Square=184). As shown in **Figure 3**, the most influential attribute (based on the LR test) is the patient's health-related lifestyle, which is about twice as important as the intervention's effectiveness and about four times as important as the patient's age and severity of illness. Among the least important attributes (or attribute interactions) are adverse effects, the intervention's type and the PES interactions with patient's age and type, where the former has more impact than the latter. Time span is the attribute that is least important. Model III in **Table 3** shows that the main effects and the PES interaction effects with 'Age of Patient' are of opposite sign. The same holds for the main effects and the PES interaction effects with 'Type of Intervention'. To assess whether the overall effects disappear when the main and PES interaction effects are combined for 'Age of Patient' or 'Type of Intervention', we tested for their significance. We found that the main and interaction effects are jointly significant at a 5% level for both attributes, illustrating that they do not cancel each other out.



**Figure 3** Importance of the seven attributes and their significant interactions with PES (MODEL III in **Table 3**) to the social value of a healthcare program relative to the most important attribute 'Lifestyle of Patient', the importance of which is set to 100

**Table 3** Estimates of coefficients in the MNL models and overall significances of the attributes using p-values obtained from likelihood ratio tests

Term	MODEL I			MODEL II Z = LO			MODEL III Z = PES		
	Estimate	LR Chi-square	P-value	Estimate	LR Chi-square	P-value	Estimate	LR Chi-square	P-value
Age of Patient (80-90 years)	-0.6200	205.919	<0.0001	-0.3547	7.967	0.0928	-1.0583	59.451	<0.0001
Age of Patient (60-70 years)	-0.0185			0.4075			-0.3567		
Age of Patient (40-50 years)	0.1210			0.0493			0.1727		
Age of Patient (20-30 years)	0.2363			0.1307			0.3516		
Age of Patient (0-10 years)	0.2812			-0.2327			0.8906		
Lifestyle of Patient (fully)	-0.3730	184.143	<0.0001	-0.3737	184.558	<0.0001	-0.3744	184.763	<0.0001
Lifestyle of Patient (partly)	0.0574			0.0582			0.0591		
Lifestyle of Patient (not at all)	0.3156			0.3155			0.3153		
Effectiveness (33%)	-0.2440	88.775	<0.0001	-0.2436	88.386	<0.0001	-0.2434	88.120	<0.0001
Effectiveness (66%)	0.0117			0.0119			0.0118		
Effectiveness (100%)	0.2323			0.2317			0.2316		
Severity of Illness (not severe)	-0.2351	46.545	<0.0001	-0.2343	46.295	<0.0001	0.2340	46.122	<0.0001
Severity of Illness (severe)	0.0758			0.0758			0.0759		
Severity of Illness (lethal)	0.1594			0.1585			0.1581		
Adverse Effects (often)	-0.1383	27.869	<0.0001	-0.1374	27.485	<0.0001	-0.1367	27.112	<0.0001
Adverse Effects (rarely)	0.0776			0.0774			0.0769		
Adverse Effects (never)	0.0607			0.0600			0.0598		
Time span (after 20 years)	-0.0617	4.771	0.0920	-0.0609	4.807	0.0904	-0.0606	4.892	0.0867
Time span (after 5 years)	0.0039			0.0026			0.0015		
Time span (within a year)	0.0579			0.0583			0.0591		
Type (preventive)	0.0127	0.348	0.5552	-0.1736	4.799	0.0285	0.1848	9.952	0.0016
Type (curative)	-0.0127			0.1736			-0.1848		
Z*Age of Patient (80-90 years)	NA	NA	NA	-0.0135	8.661	0.0702	0.0497	21.259	0.0003
Z*Age of Patient (60-70 years)	NA			-0.0216			0.0386		
Z*Age of Patient (40-50 years)	NA			0.0036			-0.0058		
Z*Age of Patient (20-30 years)	NA			0.0053			-0.0133		
Z*Age of Patient (0-10 years)	NA			0.0261			-0.0692		
Z*Type (preventive)	NA	NA	NA	0.0095	5.984	0.0144	-0.0194	9.909	0.0016
Z*Type (curative)	NA			-0.0095			0.0194		
-2*LogLikelihood	14006			13987			13968		
BIC	14145			14172			14153		

Model I: basic model with attributes of the DCE only, Model II: Model I + significant interactions with total LOT-R score (LO), Model III: Model I + significant interactions with partial LOT-R score (PES)

Coefficient estimates corresponding to the last level of an attribute, either as a main effect or involved in an interaction, are italicized and calculated as minus the sum of the estimates for the other levels of that attribute

NA means not applicable

The models are the final models after a stepwise removal of interactions between Z and the attributes lifestyle of patient, effectiveness, severity of illness, adverse effects and time span, which are non-significant at the 5% level


The modeling results are robust to the inclusion of control variables such as 'Gender' (M/F) and 'Language' (Dutch/French)



In terms of model interpretation, the following example can be instructive to assess the effect size of the PES interactions. When asked to choose between (1) a curative healthcare program for 0-10 years old with 100% effectiveness, no side effects, for a serious disease with no link with lifestyle occurring within a year, and (2) the exact same program targeted at 80-90 instead of 0-10 years old, someone with a maximal pessimism score has a marginal probability of choosing the younger patient group over the older one of 55%. Someone with minimal pessimism will have a chance of choosing the youngest group of 88%. Regarding the interaction between PES and type, when asked to choose between a curative healthcare program for 30-50 years old with 100% effectiveness and no side effects, for a severe disease with no link with lifestyle occurring within a year, and (2) the exact same program that is preventive instead of curative, someone with maximal pessimism will have 45% chance of preferring the preventive program. Someone with minimal pessimism will have a 60% chance of preferring the preventive program.

We found no difference in optimists' and pessimists' views on priority-setting when it comes to the criteria of the program's effectiveness, health-related lifestyle, severity of illness or risk of adverse effects.

**Figure 4** shows the relation between respondents having different PES scores and the different utilities they attribute to health programs according to the recipient's age (Panel A) or the preventive or curative nature of the program (Panel B). The solid lines indicate the main effect (for a PES score of zero), showing younger age groups taking priority over older ones, and prevention over cure. The dashed lines indicate how this main effect changes when we add the interaction with the respondent's PES score. The higher the respondent's PES score, the more the bonus for interventions in young people diminishes and the more equal the prioritization of different age groups becomes. The more pessimistic the respondent, the greater cure will be valued, and the lower the level of pessimism, the higher the utility of prevention. A PES score of ten represents a point of indifference, with scores below preferring prevention and scores above preferring cure.



**Figure 4** Marginal utility values for the levels of the attributes ‘Age of Patient’ (A) and ‘Type of Intervention’ (B) in interaction with respondents’ pessimism scores (PES, dashed lines) and as main effects only, independent of PES (solid lines)

## 5. Discussion

It is widely acknowledged that decisions to prioritize or deprioritize healthcare should, in one way or another, take into account social values. A wide body of research has emerged that investigates how people think about setting limits to healthcare. Typically, studies use samples that are representative of a population in terms of basic, socio-demographic characteristics such as age or gender but a sample could be non-representative in many more ways. In this study, we explored the influence of personality as a source of heterogeneity in how people want priorities to be set and potentially bias in how samples are recruited. We investigated whether differences in the personality trait of dispositional optimism translated into different views on how to set healthcare priorities. Our general hypothesis that dispositional optimism matters was confirmed, but not fully in the way we expected.

On the one hand, respondents who had a more negative outlook on the future were indeed less likely to favour prevention over cure (with cure even being more desirable for very pessimistic individuals) and had a lower willingness to prioritize younger generations (e.g. through age-weighting of health benefits). These findings are in line with those from other studies. As hypothesized in Section 2, studies showing that pessimists have a less proactive problem-solving attitude indeed suggest a lower appreciation of prevention. This is also confirmed in research showing that pessimists are less prevention-oriented when it comes to their own health behaviours [see e.g. (Radcliffe and Klein 2002, Scheier, Matthews et al. 1999, Taylor, Kemeny et al. 1992)]. In our survey, we also observed that pessimists were more likely to smoke. The finding regarding age-based priority-setting is compatible with the view that pessimists expect a worse future for themselves. Many studies indeed document that pessimists face a less healthy

old age than optimists and need more healthcare [see e.g. (Carver, Scheier et al. 2010, Carver and Scheier 2014, Brissette, Scheier et al. 2002, Carver, Lehman et al. 2003, Macleod and Conway 2005)].

On the other hand, we did not find any relationship between levels of dispositional optimism and time preference, as indicated by a non-significant interaction between respondents' dispositional optimism and the timing attribute in the DCE. Also, although our initial expectation was that optimists would be more in favour of a differentiation according to (cost-)effectiveness because of their easier disengagement from goals with unfavourable odds, no effect was found in our DCE regarding the effectiveness of health programs. Moreover, another noteworthy finding was that the effect of optimism was driven by the pessimism items: the pessimistic items and not the optimistic ones mattered. As mentioned in the Methods section, there is an ongoing debate about the dimensionality of the LOT-R. Whereas factor analyses have often revealed two separate dimensions, hereby providing statistical arguments for why our variables PES and OPT may diverge, scholars have also argued that the divergence between both constructs may have methodological reasons. Item wording and valence may explain different effects of pessimism and optimism scales [see e.g. (McPherson and Mohr 2005, Kam and Meyer 2012)]. Apart from methodological reasons and statistical observations, however, the literature does not provide many conceptual arguments for why the concept of pessimism and optimism are different and why different effects should be expected for PES and OPT. The fact that we observe effects for PES yet not for OPT may therefore give an indication of conceptual differences between OPT and PES, but it may also be explained by the question wording of both scales.

To our knowledge, our study is the first to establish a relation between personality and how people set priorities in healthcare. Yet our study also has several limitations. Although dispositional optimism is a well-established individual difference in the literature, it is likely entangled with other dispositional variables. Our study only allowed controlling for a limited number of more common factors such as age, education or health. Future studies could explore (interactions with) other personality variables. Also, we surveyed our respondents on a complex topic, in a single recording. Although our sample was broadly representative of the Belgian population according to usual demographics, we recruited respondents from an online panel, excluding those who are younger than 18 and older than 75 years. Membership to the panel, as we have discussed, may be associated with unobservable characteristics. Moreover, there are general criticisms against using DCEs to elicit social preferences (Bryan and Dolan 2004). Future studies could use more longitudinal and experimental study designs in which the consistency of the results can be assessed and where optimism can be manipulated in a treatment and control group. This can provide meaningful insights into how optimism or pessimism can be (transiently) stimulated and whether this leads to higher or lower support for particular forms of healthcare. Use of qualitative methods to understand the motivations behind respondents' choices can also be insightful. However, notwithstanding these limitations, we think that our results and suggested

explanations open up possibilities for further research. Although we acknowledge that selecting representative samples based on personality can be impractical or even unrealistic, our results do call for more discussion on how to understand the representativeness of social values studies.

**Acknowledgements** We are grateful to Dr. Michael Shiner and two anonymous reviewers for their comments on an earlier version of this manuscript.

**Authors' contributions** JL, PD and PB framed the research question and set up the experiment. RK and PG designed the experiment. RK and JL analysed the data. All authors were involved in the writing of the text.

**Funding** The authors acknowledge funding from the Research Foundation – Flanders (FWO, project numbers G098911N and G043815N, and Roselinde Kessels' postdoctoral fellowship) and Pfizer's European HTAcademy prize competition (2011) and the Antwerp Study Centre for Infectious Diseases (ASCID) at the University of Antwerp. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

## **Compliance with ethical standards**

**Conflict of interest** The authors declare that they have no conflict of interest.

**Ethical approval** All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**Ethical standards** The Committee for Medical Ethics of the University of Antwerp reviewed the study protocol, the questionnaire and the information letter for participants and approved them on 16 March 2015. The market research company Ipsos conducted the survey and provided the responses for analysis in anonymous form only.

## Appendix A: Example of a choice set

Medical interventions A and B are exactly equally expensive, and they apply to a similar number of patients. If you were forced to make a choice, which of both interventions should be reimbursed by the government? To make it easier for you, we have highlighted in yellow the characteristics that differ between both interventions. There are no right or wrong answers; we are interested in your opinion.

	<b>A</b>	<b>B</b>
<b>What type of intervention is it?</b>	Curative (meant to cure patients who are ill)	Preventive (meant to prevent healthy persons from becoming ill)
<b>How big is the probability of success of the intervention?</b>	2 in 3 is successful	Always successful
<b>How often do adverse effects occur?</b>	Often	Often
<b>How severe is the illness for which the intervention is developed?</b>	Not lethal, but everyone who gets the disease will experience a severe and lasting reduction in quality of life	Lethal, everyone who gets the disease will die from it
<b>Does the patient cause the disease through his or her own lifestyle?</b>	Not at all	Not at all
<b>How long does it take before the patient becomes ill/ shows signs/symptoms of illness?</b>	Within a year	Within a year
<b>At what age does the patient become ill?</b>	0 to 10 years	40 to 50 years
<b>YOUR PREFERENCE:</b>	<input type="radio"/>	<input type="radio"/>

## Appendix B: MNL model

Formally, the MNL model employs random utility theory which describes the utility that a respondent attaches to intervention  $j$  ( $j = 1, 2$ ) in choice set  $s$  ( $s = 1, \dots, 14$ ) as the sum of a systematic and a stochastic component:

$$U_{js} = \mathbf{x}'_{js}\boldsymbol{\beta} + \varepsilon_{js}.$$

In the systematic component,  $\mathbf{x}'_{js}\boldsymbol{\beta}$ ,  $\mathbf{x}_{js}$  is a vector containing the attribute levels of intervention  $j$  in choice set  $s$ . Additionally, in our analysis, this vector includes the interactions between the attribute levels and the LOT-R score or any other respondent variable under investigation. The vector  $\boldsymbol{\beta}$  is the vector of parameter values indicating the importance respondents attach to the different attribute levels and interactions. The stochastic component  $\varepsilon_{js}$  is the error term capturing the unobserved sources of utility. Under the assumption that the error terms are independently and identically Gumbel distributed, the MNL probability that a respondent chooses intervention  $j$  in choice set  $s$  is

$$p_{js} = \frac{\exp(\mathbf{x}'_{js}\boldsymbol{\beta})}{\exp(\mathbf{x}'_{1s}\boldsymbol{\beta}) + \exp(\mathbf{x}'_{2s}\boldsymbol{\beta})}.$$

To estimate the parameter vector  $\boldsymbol{\beta}$ , we used a maximum likelihood estimation approach, which maximizes the probability of obtaining the responses from the selected data sample. A positive estimate has a positive effect on the total utility, whereas a negative estimate has a negative effect. We computed the overall significance of the attributes and interactions by means of likelihood ratio (LR) tests. Such tests evaluate the difference in goodness of fit between nested models. More specifically, they compare the goodness of fit of an unrestricted or full model to a restricted model in which one or more parameters have been set to zero.

## References

- Andersson, M. A. (2012). "Dispositional Optimism and the Emergence of Social Network Diversity." Sociological Quarterly **53**(1): 92-115.
- Asaria, M., S. Griffin and R. Cookson (2016). "Distributional Cost-Effectiveness Analysis: A Tutorial." Med Decis Making **36**(1): 8-19.
- Berndsen, M. and J. van der Pligt (2001). "Time is on my side: Optimism in intertemporal choice." Acta Psychologica **108**(2): 173-186.
- Brissette, I., M. F. Scheier and C. S. Carver (2002). "The role of optimism in social network development, coping, and psychological adjustment during a life transition." J Pers Soc Psychol **82**(1): 102-111.
- Bryan, S. and P. Dolan (2004). "Discrete choice experiments in health economics. For better or for worse?" Eur J Health Econ **5**(3): 199-202.
- Carver, C. S., J. M. Lehman and M. H. Antoni (2003). "Dispositional pessimism predicts illness-related disruption of social and recreational activities among breast cancer patients." J Pers Soc Psychol **84**(4): 813-821.
- Carver, C. S. and M. F. Scheier (2014). "Dispositional optimism." Trends in Cognitive Sciences **18**(6): 293-299.
- Carver, C. S., M. F. Scheier and S. C. Segerstrom (2010). "Optimism." Clin Psychol Rev **30**(7): 879-889.
- Chiesi, F., S. Galli, C. Primi, P. I. Borgi and A. Bonacchi (2013). "The Accuracy of the Life Orientation Test-Revised (LOT-R) in Measuring Dispositional Optimism: Evidence From Item Response Theory Analyses." Journal of Personality Assessment **95**(5): 523-529.
- Cleemput, I. (2014). "incorporating societal preferences in reimbursement decisions: relative importance of decision criteria according to Belgian citizens." KCE Reports **234**.
- Cookson, R., S. Ali, A. Tsuchiya and M. Asaria (2018). "E-learning and health inequality aversion: A questionnaire experiment." Health Econ **27**(11): 1754-1771.
- Corr, P. and G. Matthews (2009). The cambridge handbook of personality psychology. New York, Cambridge University Press.
- Daniels, N. and J. Sabin (2008). Setting Limits Fairly: Learning to share resources for health. New York, Oxford University Press.
- Dolan, P., R. Shaw, A. Tsuchiya and A. Williams (2005). "QALY maximisation and people's preferences: a methodological review of the literature." Health Economics **14**(2): 197-208.
- Faust, H. and P. Menzel (2012). Prevention vs Treatment: what's the right balance?. New York, Oxford University Press.
- Geers, A. L., J. A. Wellman and G. D. Lassiter (2009). "Dispositional optimism and engagement: the moderating influence of goal prioritization." J Pers Soc Psychol **96**(4): 913-932.
- Geers, A. L., J. A. Wellman, L. D. Seligman, L. A. Wuyek and L. A. Neff (2010). "Dispositional optimism, goals, and engagement in health treatment programs." J Behav Med **33**(2): 123-134.
- Gibson, B. and D. M. Sanbonmatsu (2004). "Optimism, pessimism, and gambling: the downside of optimism." Pers Soc Psychol Bull **30**(2): 149-160.
- Giles, L. C., G. F. Glonek, M. A. Luszcz and G. R. Andrews (2005). "Effect of social networks on 10 year survival in very old Australians: the Australian longitudinal study of aging." J Epidemiol Community Health **59**(7): 574-579.
- Glaesmer, H., W. Rief, A. Martin, R. Mewes, E. Brahler, M. Zenger and A. Hinz (2012). "Psychometric properties and population-based norms of the Life Orientation Test Revised (LOT-R)." British Journal of Health Psychology **17**: 432-445.
- Gu, Y., E. Lancsar, P. Ghijben, J. R. Butler and C. Donaldson (2015). "Attributes and weights in health care priority setting: A systematic review of what counts and to what extent." Soc Sci Med **146**: 41-52.
- Heaton, J. B. (2002). "Managerial optimism and corporate finance." Financial Management **31**(2): 33-45.
- Hecht, D. (2013). "The neural basis of optimism and pessimism." Exp Neurobiol **22**(3): 173-199.

- Heinonen, K., K. Raikkonen, K. A. Matthews, M. F. Scheier, O. T. Raitakari, L. Pulkki and L. Keltikangas-Jarvinen (2006). "Socioeconomic status in childhood and adulthood: associations with dispositional optimism and pessimism over a 21-year follow-up." *J Pers* **74**(4): 1111-1126.
- Herdman, M., C. Gudex, A. Lloyd, M. Janssen, P. Kind, D. Parkin, G. Bonsel and X. Badia (2011). "Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L)." *Qual Life Res* **20**(10): 1727-1736.
- Kam, C. and J. P. Meyer (2012). "Do optimism and pessimism have different relationships with personality dimensions? A re-examination." *Personality and Individual Differences* **52**(2): 123-127.
- Kim, E. S., N. Park and C. Peterson (2011). "Dispositional optimism protects older adults from stroke: the Health and Retirement Study." *Stroke* **42**(10): 2855-2859.
- Lancsar, E., J. Wildman, C. Donaldson, M. Ryan and R. Baker (2011). "Deriving distributional weights for QALYs through discrete choice experiments." *J Health Econ* **30**(2): 466-478.
- Luyten, J., R. Kessels, P. Goos and P. Beutels (2015). "Public preferences for prioritizing preventive and curative health care interventions: a discrete choice experiment." *Value in Health* **18**(2).
- Macleod, A. K. and C. Conway (2005). "Well-being and the anticipation of future positive experiences: The role of income, social networks, and planning ability." *Cogn Emot* **19**(3): 357-374.
- Marcus, B. and A. Schutz (2005). "Who are the people reluctant to participate in research? Personality correlates of four different types of nonresponse as inferred from self- and observer ratings." *J Pers* **73**(4): 959-984.
- Matthews, K. A., K. Raikkonen, K. Sutton-Tyrrell and L. H. Kuller (2004). "Optimistic attitudes protect against progression of carotid atherosclerosis in healthy middle-aged women." *Psychosom Med* **66**(5): 640-644.
- McFadden, D. (1974). Conditional logit analysis of qualitative choice behavior. *Frontiers in Econometrics* P. Zarembka. New York, Academic Press: 105-142.
- McPherson, J. and P. Mohr (2005). "The role of item extremity in the emergence of keying-related factors: an exploration with the life orientation test." *Psychol Methods* **10**(1): 120-131.
- Mooney, G. H. and S. H. Blackwell (2004). "Whose health service is it anyway? Community values in healthcare." *Med J Aust* **180**(2): 76-78.
- Mosing, M. A., S. D. Gordon, S. E. Medland, D. J. Statham, E. C. Nelson, A. C. Heath, N. G. Martin and N. R. Wray (2009). "Genetic and environmental influences on the co-morbidity between depression, panic disorder, agoraphobia, and social phobia: a twin study." *Depress Anxiety* **26**(11): 1004-1011.
- Nes, L. S., D. R. Evans and S. C. Segerstrom (2009). "Optimism and College Retention: Mediation by Motivation, Performance, and Adjustment(1)." *Journal of Applied Social Psychology* **39**(8): 1887-1912.
- NICE (2008). Social Value Judgements: Principles for the development of NICE guidance <https://www.nice.org.uk/proxy/?sourceUrl=http%3a%2f%2fwww.nice.org.uk%2fmedia%2fC18%2f30%2fSVJ2PUBLICATION2008.pdf>.
- Olsen, J. A., J. Richardson, P. Dolan and P. Menzel (2003). "The moral relevance of personal characteristics in setting health care priorities." *Soc Sci Med* **57**(7): 1163-1172.
- OMHLTC (2013). A report on the ontario citizen's council QALYs and drug funding decisions in Ontario. O. m. o. h. a. l.-t. care.
- Pavlova, M. K. and R. K. Silbereisen (2013). "Dispositional optimism fosters opportunity-congruent coping with occupational uncertainty." *J Pers* **81**(1): 76-86.
- Puri, M. and D. T. Robinson (2007). "Optimism and economic choice." *Journal of Financial Economics* **86**(1): 71-99.
- Radcliffe, N. M. and W. M. P. Klein (2002). "Dispositional, unrealistic, and comparative optimism: Differential relations with the knowledge and processing of risk information and beliefs about personal risk." *Personality and Social Psychology Bulletin* **28**(6): 836-846.
- Rauch, W. A., K. Schweizer and H. Moosbrugger (2007). "Method effects due to social desirability as a parsimonious explanation of the deviation from unidimensionality in LOT-R scores." *Personality and Individual Differences* **42**(8): 1597-1607.
- Rius-Ottenheim, N., D. Kromhout, A. J. de Craen, J. M. Geleijnse, R. C. van der Mast, F. G. Zitman, R. G. Westendorp, P. E. Slagboom and E. J. Giltay (2012a). "Parental longevity correlates with



- offspring's optimism in two cohorts of community-dwelling older subjects." Age (Dordr) **34**(2): 461-468.
- Rius-Ottenheim, N., D. Kromhout, R. C. van der Mast, F. G. Zitman, J. M. Geleijnse and E. J. Giltay (2012b). "Dispositional optimism and loneliness in older men." Int J Geriatr Psychiatry **27**(2): 151-159.
- Roberts, B. W., N. R. Kuncel, R. Shiner, A. Caspi and L. R. Goldberg (2007). "The Power of Personality: The Comparative Validity of Personality Traits, Socioeconomic Status, and Cognitive Ability for Predicting Important Life Outcomes." Perspectives on Psychological Science **2**(4): 313-345.
- RobinsonWhelen, S., C. T. Kim, R. C. MacCallum and J. K. KiecoltGlaser (1997). "Distinguishing optimism from pessimism in older adults: Is it more important to be optimistic or not to be pessimistic?" Journal of Personality and Social Psychology **73**(6): 1345-1353.
- Ryan, M., K. Gerard and M. Amaya-Amaya (2008). "Using Discrete Choice Experiments to Value Health and Health Care Preface." Using Discrete Choice Experiments to Value Health and Health Care **11**: Xiii-Xiv.
- Scheier, M. F. and C. S. Carver (1985). "Optimism, Coping, and Health - Assessment and Implications of Generalized Outcome Expectancies." Health Psychology **4**(3): 219-247.
- Scheier, M. F., C. S. Carver and M. W. Bridges (1994). "Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test." J Pers Soc Psychol **67**(6): 1063-1078.
- Scheier, M. F., K. A. Matthews, J. F. Owens, R. Schulz, M. W. Bridges, G. J. Magovern and C. S. Carver (1999). "Optimism and rehospitalization after coronary artery bypass graft surgery." Arch Intern Med **159**(8): 829-835.
- Schwappach, D. L. (2002). "Resource allocation, social values and the QALY: a review of the debate and empirical evidence." Health Expect **5**(3): 210-222.
- Seegerstrom, S. C. (2007). "Optimism and Resources: Effects on Each Other and on Health over 10 Years." J Res Pers **41**(4).
- Seegerstrom, S. C., D. R. Evans and T. A. Eisenlohr-Moul (2011). "Optimism and pessimism dimensions in the Life Orientation Test-Revised: Method and meaning." Journal of Research in Personality **45**(1): 126-129.
- Shah, K. K. (2009). "Severity of illness and priority setting in healthcare: a review of the literature." Health Policy **93**(2-3): 77-84.
- Sirven, N. and T. Debrand (2012). "Social capital and health of older Europeans: causal pathways and health inequalities." Soc Sci Med **75**(7): 1288-1295.
- Smith, S. T., J. F. Edens, M. Epstein, P. G. Stiles and N. G. Poythress, Jr. (2012). "Do the psychopathic personality traits of fearless dominance and self-centered impulsivity predict attitudes about and influences on research participation?" Behav Sci Law **30**(1): 1-15.
- Srivastava, S., K. M. McGonigal, J. M. Richards, E. A. Butler and J. J. Gross (2006). "Optimism in close relationships: How seeing things in a positive light makes them so." Journal of Personality and Social Psychology **91**(1): 143-153.
- Standaard, D. (2014). "G20 000 voor medicijnen." [http://www.standaard.be/cnt/dmf20140302\\_01006486](http://www.standaard.be/cnt/dmf20140302_01006486).
- Taylor, S. E., M. E. Kemeny, L. G. Aspinwall, S. G. Schneider, R. Rodriguez and M. Herbert (1992). "Optimism, coping, psychological distress, and high-risk sexual behavior among men at risk for acquired immunodeficiency syndrome (AIDS)." J Pers Soc Psychol **63**(3): 460-473.
- Tindle, H. A., Y. F. Chang, L. H. Kuller, J. E. Manson, J. G. Robinson, M. C. Rosal, G. J. Siegle and K. A. Matthews (2009). "Optimism, cynical hostility, and incident coronary heart disease and mortality in the Women's Health Initiative." Circulation **120**(8): 656-662.