This item is the archived peer-reviewed author-version of:

Mutual HIV status disclosure is associated with consistent condom use in public sector ART clients in Free State province, South Africa : a short report

Reference:

le Roux Booysen Frederik, Wouters Edwin, de Walque Damien, Over Mead.- Mutual HIV status disclosure is associated with consistent condom use in public sector ART clients in Free State province, South Africa : a short report
Aids care - ISSN 0954-0121 - 29:11(2017), p. 1386-1390
Full text (Publisher's DOI): <https://doi.org/doi:10.1080/09540121.2017.1290210>
To cite this reference: <http://hdl.handle.net/10067/1407360151162165141>

Title:

Mutual HIV status disclosure is associated with consistent condom use in public sector ART clients in Free State province, South Africa: a short report

Acknowledgement: We thank the study participants and the fieldwork staff as well as the Free State Department of Health (FSDOH) and National Health Laboratory Service (NHLS). This work was supported by: Research Committee of the World Bank; The Bank Netherlands Program Partnership; WB-DfiD 'Evaluation of the Community Response to HIV and AIDS'; Programme to Support Pro-Poor Policy Development (PSPPD), a partnership between the Presidency, Republic of South Africa and the European Union; Health Economics and Aids Research Division (HEARD) at the University of Kwazulu-Natal (UKZN); University of the Free State (UFS); and South Africa's National Research Foundation (NRF). The findings, interpretations and conclusions in the paper are those of the authors and do not reflect the views of funding agencies.

Abstract:

Risky sexual behaviour in PLWHA on antiretroviral therapy threatens both prevention and treatment efforts, but disclosure promises to support safer sexual practices. This paper investigates the association between HIV self-disclosure and consistent condom use in a cohort of public sector patients on antiretroviral (ARV) treatment. Using data from the FEATS cohort study, logistic regression analysis shows that knowledge of your partner's HIV status is positively associated with consistent condom use (OR 2.73, 95% CI 1.37-5.43, $p=0.004$) and so too mutual HIV disclosure (OR 3.38, 95% CI 1.60-7.18, $p=0.001$). Prevention and treatment programmes, through couple HIV counselling and testing (CHCT) and other assistance programmes, should focus on supporting the mutual disclosure of HIV status among PLWHA on ARV treatment.

Keywords:

antiretroviral therapy, ART, condom use, HIV disclosure, HIV/AIDS, South Africa

Word count: 1,519

Introduction

PLWHA on antiretroviral (ARV) treatment experience improvements in their life expectancy, health and quality of life, and concomitantly in sexual activity (Berhan & Berhan, 2012; Hanif, Bastos, Malta, Bertoni, Winch & Kerrigan, 2014; Kaye, Kakaire, Osinde, Lule & Kakande, 2013). Risky sexual behaviour in HIV-positive individuals on ART is of critical importance, given that it stands to facilitate the on-going transmission of the virus in the non-infected and the re-infection of the treated with drug-resistant strains of the HIV virus (Hanif et al., 2014; Kaye et al., 2013; Shewamene, Legesse, Tsega, Bhagavathula & Endale, 2015).

One potential avenue for impacting sexual behaviour is disclosure. In the Disclosure Processes Model (DPM), sexual practices are described as a longer-term, health-related dyadic outcome of self-disclosure (Chaudoir & Fisher, 2010). The disclosure process, therefore, stands to open avenues for negotiating condom use (Bachanas et al., 2013; Conserve et al., 2015; Deribe, Woldemichael, Wondafrash, Haile & Amberbir, 2008; Li et al., 2013), thus contributing to safer sexual practices.

Although some studies have investigated HIV disclosure by PLWHA on ART (Ebuenyi et al., 2014; Hirsch et al., 2014; Seid, Wasie & Admassu, 2012; Yaya et al., 2015) and numerous the sexual behaviour of PLWHA on ART (Akinyemi, Awolude, Adewole & Kanki, 2010; Anene, Ighodaro & Olufemi, 2014; Ayiga, 2012; Crepaz & Marks, 2004; Diabaté et al., 2013; Eisele et al., 2008; Kaida et al., 2008; Kennedy, Medley & Sweat, 2007; Mawji et al., 2012; Peltzer & Ramlagan, 2010; Sarna et al., 2008; Wilson & Minkoff, 2001), only a handful of cross-sectional studies from sub-Saharan Africa have investigated the HIV disclosure – sexual behaviour nexus (Adebayo et al., 2014; Bachanas et al., 2013; Fekadu, Addisie & Mellie, 2014). The current analysis also investigates the association between HIV self-disclosure and consistent condom use, but uses longitudinal data for a cohort of public sector patients on ARV treatment.

Methods

Participants and procedures:

The Effective Aids Treatment and Support in the Free State (FEATS) study is a longitudinal, cohort study. As part of the evaluation component of the study, which comprised a randomized control trial of a peer adherence and nutritional support intervention (Booyesen, De Walque, Over, Hashimoto & De Reuck, 2016), a total of 655 adult ARV patients was recruited into the study at twelve primary health care clinics in the Free State province of South Africa. Patients were questioned on their sexual behaviour and on HIV disclosure. At baseline, ninety-six percent (630) of patients completed the survey. A total of 453 (71.9%) and 413 (65.5%) of patients were re-interviewed during two follow-ups (N=1,496), conducted 14.9 [IQR 13.5-15.9] and 9.8 [IQR 8.8-10.6] months apart, respectively. The aggregate attrition was 44.1%, with the 352 patients in the analytical sample interviewed in all three survey rounds (N=1,056).

Ethical clearance:

Ethical approval for the study was obtained from the Medical School of the University of the Free State (ETOVS 145/07). Written informed consent was obtained from all study participants.

Measures:

Consistent condom use, the dependent variable, means reporting always using a condom during sex over the past six months. In the case of HIV disclosure, the main independent variable, a distinction is made between “*self-disclosure*”; disclosing your HIV status to your partner, “*partner disclosure*”; knowing your partner’s HIV status, and “*mutual disclosure*”; both knowing your partner’s HIV status and having disclosed your own HIV status to your partner. In addition, a categorical variable was constructed, where the variable takes on a value of “1” (neither partner disclosed), “2” (one partner only disclosed) or “3” (both partners disclosed). Classification is based on the response to two questions, i.e. “Does your partner know your HIV status” and “Do you know the HIV status of your partner”.

Control variables include the following: sex, age (years), marital status, education, health-related quality of life (EQ-VAS), substance use (alcohol, tobacco or other drugs), ART optimism (standardised index), ARV treatment duration (months), survey round, and district.

Analysis:

The socio-demographic characteristics of the sample are first described. Following a bivariate analysis, logistic regression models are employed to investigate the link between HIV disclosure (independent variable) and consistent condom use (dependent variable). The criterion for statistical significance is $p \leq 0.05$. Analysis was conducted using *Stata14*.

Results

Participants:

Study participants are predominantly female (76.0%) with a mean age of 37 years on average. Few patients have tertiary education (1.4%), while almost one in five persons have completed grade 12 (18.1%). Only 3.5% have no education, 30.9% primary, and 45.9% some secondary. Two thirds of study participants are single (66.6%) while 23.8% live with their spouse or partner.

Sexual behaviour and disclosure (Table 1/2):

Half of patients were sexually active (50.1%). Consistent condom use was high (79.2%) (Table 1). A majority of patients have disclosed their own HIV status (87.5%) or knew their partner's HIV status (75.1%). Mutual disclosure is common (74.2%). According to Table 1, the percentage of respondents reporting that they know their partner's HIV status increased significantly post-baseline ($p=0.009$), while the proportion of cases where the HIV status of one partner only was known concomitantly declined ($p=0.017$). Condom use was consistently higher where HIV status had been disclosed to or by a partner ($p \leq 0.004$) or to both partners (<0.001) (Table 2).

Regression analysis (Table 3):

Where a patient knows their partner's HIV status, the likelihood of consistent condom use increases 2.73 fold (CI 1.37-5.43, $p=0.004$). The association is even stronger for mutual disclosure (OR 3.38, CI 1.60-7.18, $p=0.001$). Age also predicts consistent condom use, but only in one regression model, model 1. An increase in age of one year, increases the likelihood of consistent condom use by a marginal 3.6% (CI 1.00-1.07, $p=0.032$). In the accompanying bivariate analysis, disclosure of one's own HIV status was also positive and statistically significant (OR 2.33, CI 1.20-4.51, $p=0.012$).

Discussion

Levels of disclosure fall within the range reported in the literature: 60.9% in Togo (Yaya et al., 2015) and 93.1% in Ethiopia (Seid et al., 2012). Consistent condom use too is similar to rates reported in other studies, ranging from 60.8% in Assela, Ethiopia (Fekadu et al., 2014) to 78.9% in north-western Ethiopia (Shewamene et al., 2015). Studies generally ignore partner HIV status (Venkatesh, Flanigan & Mayer, 2011). As hypothesised in the Disclosure Processes Model (DPM) (Chaudoir & Fisher, 2010; Chaudoir et al., 2011), this study finds that knowledge of your partner's HIV status and mutual HIV disclosure are associated with safer sexual practices, as did Bachanas et al. (2013) in a study in Kenya, Namibia and Tanzania.

This finding highlights the importance of disclosure assistance. Walcott, Hatcher, Kwena and Turan (2013) emphasise that PLWHA be given different options for HIV disclosure assistance, options that promote the relationship building skills and knowledge of communication strategies necessary for mutual disclosure (Hightow-Weidman et al., 2013), including couple HIV counselling and testing (CHCT) (Walcott et al., 2013). A recent systematic review found evidence in support of media- and computer-based as well as facilitator- and group-based disclosure programmes (Conserve et al., 2015). Kennedy, Fonner, Armstrong, O'Reilly and Sweat (2015), in another systematic review, find mixed evidence and modest effects, with the strongest evidence for partner notification interventions.

This study has various limitations. Firstly, the results are not representative of the larger South African ART programme. Secondly, reports on sexual behaviour and HIV disclosure are based on self-report and subject to social desirability bias. Thirdly, partners' HIV status was not recorded. As a result, it is not possible to investigate condom use in sero-concordant and non-concordant couples. Fourth, the relationship between disclosure and condom use is bi-directional and causality cannot be inferred from this analysis. Disclosure is a complex process (Chaudoir, Fisher & Simoni, 2011) and has various positive and negative consequences (Conserve, Groves & Maman, 2015; Genet, Sebsibie & Gultie, 2015). This study, in the fifth instance, cannot reflect on the extent to which the study participants' disclosure experience was positive or negative nor can it shed light on the full complexity of the disclosure process as experienced by study participants. Sixth, disclosure, through serosorting, may also encourage risky sexual behaviour in HIV-positive sero-concordant couples. Finally, no data was collected on desire to have children, which is a key factor in explaining condom use.

Conclusion

Prevention and treatment programmes, through couple HIV counselling and testing (CHCT) and other disclosure assistance programmes, should focus on supporting the mutual disclosure of HIV status among PLWHA on antiretroviral (ARV) treatment.

Disclosure statement

The authors have no conflict of interest.

References

Adebayo, A.M., Ilesanmi, O.S., Omotoso, B.A., Ayodeji, O.O., Kareem, A.O. & Alele, F.O. (2014). Disclosure to sexual partner and condom use among HIV positive clients attending ART clinic at a tertiary health facility in South West Nigeria. *Pan African Medical Journal*, 19, 245. doi: 10.11604/pamj.2014.18.245.4371

Akinyemi, J.O., Awolude, O.A., Adewole, I.F. & Kanki, P.J. (2010). Condom use among antiretroviral therapy patients in Ibadan, Nigeria. *Journal of Infectious Diseases in Developing Countries*, 4(8), 495-502.

Anene, A.K., Ighodaro, I. & Olufemi, I.S. (2014). The Sexual Behaviours of HIV Positive Patients Receiving Antiretroviral Therapy in HIV Treatment Centre in Nigeria. *Journal of AIDS & Clinical Research*, 5(12), 1000392. doi: 10.4172/2155-6113.1000392

Ayiga, N. (2012). Rates and Predictors of Consistent Condom-Use by People Living with HIV/AIDS on Antiretroviral Treatment in Uganda. *Journal of Health, Population and Nutrition*, 30(3), 270-280.

Bachanas, P., Medley, A., Pals, S., Kidder, D., Antelman, G., Benech, I. ... Bukuku, M. (2013). Disclosure, Knowledge of Partner Status, and Condom Use Among HIV-Positive Patients Attending Clinical Care in Tanzania, Kenya and Namibia. *Aids Patient Care and STDs*, 27(7), 425-433. doi: 10.1089/apc.2012.0388

Berhan, A. & Berhan, Y. (2012). Is the Sexual Behaviour of HIV Patients on Antiretroviral therapy safe or risky in Sub-Saharan Africa? Meta-Analysis and Meta-Regression. *AIDS Research and Therapy*, 9, 14. Retrieved from <http://www.aidsrestherapy.com/content/9/1/14>

Booyesen, F., De Walque, D., Over, M., Hashimoto, S. & De Reuck, C., (2016). *A Randomized Control Trial of a Peer Adherence and Nutritional Support Program*

for Public Sector Antiretroviral Patients (World Bank Policy Research Working Paper 7760). Washington, DC: The World Bank Development Research Group.

Chaudoir, S.R. & Fisher, J.D. (2010). The Disclosure Processes Model: Understanding Disclosure Decision Making and Postdisclosure Outcomes Among People Living With a Concealable Stigmatized Identity. *Psychological Bulletin*, 136(2), 236-256. doi: 10.1037/a0018193

Chaudoir, S.R., Fisher, J.D. & Simoni, J.M. (2011). Understanding HIV disclosure: A review and application of the Disclosure Processes Model. *Social Science & Medicine*, 72, 1618-1629. doi: 10.1016/j.socscimed.2011.03.028

Conserve, D.F., Groves, A.K. & Maman, S. (2015). Effectiveness of Interventions Promoting HIV Serostatus Disclosure to Sexual Partners: A Systematic Review. *AIDS & Behavior*, 19, 1763-1772. doi: 10.1007/s10461-015-1006-1

Crepaz, N., Hart, T.A. & Marks, G. (2004). Highly active antiretroviral therapy and sexual risk behaviour: a meta-analytic review. *JAMA*, 292, 224-236. doi: 10.1001/jama.292.2.224

Deribe, K., Woldemichael, K., Wondafrash, M., Haile, A. & Amberbir, A. (2008). Disclosure experience and associated factors among HIV positive men and women clinical service users in southwest Ethiopia. *BMC Public Health*, 8, 81. doi: 10.1186/1471-2458-8-81

Diabaté, S., Chamberland, A., Zannou, D.M., Geraldo, N., Azon-Kouanou, A., Massinga-Loembe, M., ... Alary, M. (2013). Sexual behaviour after antiretroviral therapy initiation in female sex workers and HIV-positive patients from the general population, Cotonou, Benin. *AIDS Care*, 25(11), 1426-1432. doi: 10.1080/09540121.2013.772279

Ebuenyi, I.D., Ogoina, D., Ikuabe, P.O., Harry, T.C., Inatimi, O. & Chukwueke, O.U. (2014). Prevalence Pattern and Determinants of Disclosure of HIV Status in an

Antiretroviral Therapy Clinic in the Niger Delta Region of Nigeria. *African Journal of Infectious Disease*, 8(2), 27-30. doi:

Eisele, T.P., Matthews, C., Chopra, M., Brown, L., Silvestre, E., Daries, V. & Kendall, C. (2008). High Levels of Risk Behavior Among People Living with HIV Initiating and Waiting to Start Antiretroviral Therapy in Cape Town South Africa. *AIDS & Behavior*, 12, 570-577. doi: 10.1007/s10461-007-9279-7

Fekadu, H., Adissie, M. & Mellie, H. (2014). Sero Status Disclosure and Condom use among Plwhas on Art in Assela Town Health Facilities, Oromiya Region. *Journal of AIDS & Clinical Research*, 6(1), 1000414. doi: 10.4172/2155-6113.1000414

Genet, M., Sebsibie, G. & Gultie, T. (2015). Disclosure of HIV seropositive status to sexual partners and its associated factors among patients attending antiretroviral treatment clinic follow up at Mekelle Hospital, Ethiopia: a cross-sectional study. *BMC Research Notes*, 8, 109. doi: 10.1186/s13104-015-1056-5

Hanif, H., Bastos, F.I., Malta, M., Bertoni, L., Winch, P.J. & Kerrigan, D. (2014). Where does Treatment Optimism Fit in? Examining Factors Associated with Consistent Condom Use Among People Receiving Antiretroviral Treatment in Rio de Janeiro, Brazil. *AIDS & Behavior*, 18, 1945-1954. doi: 10.1007/s10461-014-0711-5

Hightow-Weidman, L.B., Phillips, G., Outlaw, A.Y., Wohl, A.R., Fields, S., Hildalgo, J. & LeGrand, S. (2013). Patterns of HIV Disclosure and Condom Use Among HIV-Infected Young Racial/Ethnic Minority Men Who Have Sex With Men. *AIDS & Behavior*, 17, 360-368. doi: 10.1007/s10461-012-0331-x

Hirsch, A., Forrest, J.I., Kanter, S., O'Brien, N., Salters, K.A., McCandless, L. ... Hogg, R.S. (2014). Factors Associated with Disclosure of HIV Status Among a Cohort of Individuals on Antiretroviral Therapy in British Columbia, Canada. *AIDS & Behavior*, 18, 1014-1026. doi: 10.1007/s10461-013-0623-9

Kaida, A., Gray, G., Bastos, F.I., Andia, I., Maier, M., McIntyre, J., ... Hogg, R. (2008). The relationship between HAART use and sexual activity among HIV-positive women of reproductive age in Brazil, South Africa, and Uganda. *AIDS Care*, 20(1), 21-25. doi :10.1080/09540120701426540

Kaye, D.K., Kakaire, O., Osinde, M.O., Lule, J.C. & Kakande, N. (2013). The impact of highly active antiretroviral therapy on high-risk behaviour of HIV-infected patients in sub-Saharan Africa. *Journal of Infectious Diseases in Developing Countries*, 7(6), 436-447. doi: 10.3855/jidc.2644

Kennedy, C.E., Fonner, V.A., Armstrong, K.A., O'Reilly, K.R. & Sweat, M.D. (2015). Increasing HIV serostatus disclosure in low and middle-income countries: a systematic review of intervention evaluations. *AIDS*, 29, S7-S23. doi: 10.1097/QAD.0000000000000671

Kennedy, C., O'Reilly, K., Medley, A. & Sweat, M. (2007). The impact of HIV treatment on risk behaviour in developing countries: a systematic review. *AIDS Care*, 19, 707-720. doi: 10.1080/09540120701203261

Li, G., Lu, H., Li, X., Sun, Y., He, X., Fan, S. ... Shao, Y. (2013). Mutual HIV Disclosure among HIV-Negative Men Who have Sex with Men in Beijing, China, 2010. *Archives of Sexual Behaviour*, 42, 1267-1273. doi: 10.1007/s10508-012-9944-4

Mawji, E., McKinnon, L., Wachichi, C., Chege, D., Thottingale, P., Kariri, A., ... Kaul, R. (2012). Does antiretroviral therapy initiation increase sexual risk taking in Kenyan female sex workers? A retrospective, case-control study. *BMJ Open*, 2, e000565. doi: 10.1136/bmjopen-2011-000565

Peltzer, K. & Ramlagan, S. (2010). Safer sexual behaviours after 1 year of antiretroviral treatment in Kwazulu-Natal, South Africa: a prospective cohort study. *Sexual Health*, 7(2), 135-141. doi: 10.1071/SH09109

Sarna, A., Luchters, S.M.F., Geibel, S., Kaai, S., Munyao, P., Shikely, S., ... Temmerman, M. (2008). Sexual risk behaviour and HAART: a comparative study of HIV-infected persons on HAART and on preventive therapy in Kenya. *International Journal of STD & AIDS*, 19, 85-89. doi: 10.1258/ijsa.2007.007097

Seid, M., Wasie, B. & Admassu, M. (2012). Disclosure of HIV Positive Result to a Sexual Partner among Adult Clinical Services Users in Kemissie District, Northeast Ethiopia. *African Journal of Reproductive Health*, 16(1), 97-104. doi:

Shewamene, Z., Legesse, B., Tsega, B., Bhagavathula, A.S. & Endale, A. (2015). Consistent condom use in HIV/AIDS patients receiving antiretroviral therapy in northwestern Ethiopia: implication to reduce transmission and multiple infections. *HIV/AIDS – Research and Palliative Care*, 7, 119-124. doi: 10.2147/HIV.S79122

Venkatesh, K.K., Flanigan, T.P. & Mayer, K.H. (2011). Is expanded HIV treatment preventing new infections? Impact of antiretroviral therapy on risky sexual behaviors in the developing world. *AIDS*, 25, 1939-1949. doi: 10.1097/QAD.0b013e32834b4ced

Walcott, M.M., Hatcher, A.M., Kwen, Z. & Turan, J.M. (2013). Facilitating HIV status disclosure for pregnant women and partners in rural Kenya: a qualitative study. *BMC Public Health*, 13, 1115. doi: 10.1186/1471-2458-13-1115

Wilson, T.E. & Minkoff, H. (2001). Condom Use Consistency Associated With Beliefs Regarding HIV Disease Transmission Among Women Receiving HIV Antiretroviral Therapy. *Journal of Acquired Immune Deficiency Syndromes*, 27, 289-291.

Yaya, I., Saka, B., Landoh, D.E., Patchali, P.M., Patassi, A.A., Aboubakari, A. ... Pitché, P. (2015). HIV Status Disclosure to Sexual Partners, among People Living with HIV and AIDS on Antiretroviral Therapy at Sokodé Regional Hospital, Togo. *PLoS ONE*, 10(2), e0118157. doi: 10.1371/journal.pone.0118157

Table 1: Condom use and disclosure (%), by survey round

	Survey round			Total	p-value
	1	2	3		
Condom use (%)	77.1	83.0	78.6	79.6	0.381
Disclosed own status (%)	88.0	88.3	89.1	88.5	0.937
Partner's status known (%)	68.4	79.5	78.1	75.7	0.044
Disclosure (%)					
Neither partner	10.7	10.5	9.4	10.2	0.054
One partner only	22.1	10.5	13.6	15.1	
Both partners	67.1	78.8	76.9	74.7	
Total	100.0	100.0	100.0	100.0	

Note: Condom use refers to consistent condom use, i.e. reportedly using a condom "every time" in the past six months.

Numbers may not add up due to rounding.

Table 2: Consistent condom use (%), by HIV disclosure status

	Percentage (%)	p-value
Disclosed own status:		
No	66.1	0.004
Yes	81.9	
Partner's status known:		
No	65.0	<0.001
Yes	84.7	
Disclosure:		
Neither partner	67.3	<0.001
One partner only	63.6	
Both partners	85.2	

Note: Consistent condom use refers to reportedly using a condom "every time" in the past six months.

Table 3: Logistic regression models – consistent condom use

	Bivariate		Model 1		Model 2	
	OR	p-value	OR	p-value	OR	p-value
Female	0.654	0.148	0.724	0.316	0.729	0.413
Age (years)	1.024	0.145	1.036	0.032	1.035	0.056
Marital status (comparison = single):						
Not cohabiting	2.466	0.139	1.956	0.205	1.962	0.334
Cohabiting	0.829	0.460	0.552	0.052	0.543	0.083
Education (comparison = none)						
Primary	1.206	0.926	1.279	0.709	1.264	0.913
Secondary	1.084	0.969	1.234	0.751	1.236	0.923
Grade 12 or higher	1.321	0.891	1.186	0.813	1.236	0.923
EQ-VAS	1.002	0.728	1.000	0.946	0.999	0.977
Substance use (yes)	0.731	0.252	0.712	0.319	0.712	0.357
ART optimism	1.086	0.571	1.063	0.712	1.075	0.660
ARV treatment duration (months)	1.004	0.733	1.038	0.529	1.044	0.414
Individual disclosure:						
Disclosed own status (yes)	2.332	0.012	1.419	0.426		
Knows partner's status (yes)	2.988	<0.001	2.736	0.004		
Mutual disclosure:						
Disclosure (comparison = neither partner)						
One partner only	0.850	0.642			0.981	0.966
Both partners	2.810	0.001			3.389	0.001
Sample (N)			488		488	
Wald chi2 (p-value)			35.77 (0.011)		39.66 (0.003)	
Pseudo R ²			0.081		0.087	

Note: Results are for pooled regression models and are adjusted for district and survey round. The likelihood ratio (LR) test (not reported here) indicates that the pooled regression model in each case outperforms the random effects (RE) panel model ($p > 0.05$). P-values are based on bootstrapped standard errors.