

**This item is the archived peer-reviewed author-version of:**

Contrasting response of European forest and grassland energy exchange to heatwaves

**Reference:**

Teuling Adriaan J., Seneviratne Sonia I., Stöckli Reto, Reichstein Markus, Moors Eddy, Ciais Philippe, Luysaert Sebastiaan, van den Hurk Bart, Ammann Christof, Bernhofer Christian, ....- Contrasting response of European forest and grassland energy exchange to heatwaves  
Nature geoscience - ISSN 1752-0894 - 3:10(2010), p. 722-727  
Full text (Publisher's DOI): <https://doi.org/10.1038/NGEO950>  
To cite this reference: <https://hdl.handle.net/10067/853690151162165141>

# **Contrasting response of European forest and grassland energy exchange to heatwaves**

## **Supplementary information**

Teuling et al. (2010)

**Supplementary Table 1** | List of flux tower sites, site characteristics, and site references for additional site information (see full references on following pages). For the sites without reference we refer to [www.fluxdata.org](http://www.fluxdata.org) for additional information.  $T_{max}$  is the mean summer (JJA) daily maximum temperature excluding heatwave months June-August 2003 and July 2006.


Abbr.	Site name	Lat.	Long.	Elev. (m)	Clim.	IGBP class	Years	$T_{max}$ (°C)	Site reference
AT-Neu	Neustift/Stubai Valley	47.117	11.318	970.	Cfb	Grassland	2002–2006	21.41	Hammerle et al. (2008)
BE-Bra	Brasschaat (De Inslag Forest)	51.309	4.521	16.	Cfb	Mixed forest	1997–2006	20.60	Gielen et al. (2010)
BE-Lon	Lonzee	50.552	4.745	165.	Cfb	Cropland	2004–2006	21.29	Moureaux et al. (2006)
BE-Vie	Vielsalm	50.306	5.997	450.	Cfb	Mixed forest	1996–2006	18.89	Aubinet et al. (2001)
CH-Oe1	Oensingen 1	47.286	7.732	450.	Cfb	Grassland	2002–2006	22.49	Ammann et al. (2007)
DE-Bay	Bayreuth-Waldstein	50.142	11.867	775.	Cfb	Evergreen needleleaf forest	1996–1999	17.76	Staudt et al. (2007)
DE-Geb	Gebesee	51.100	10.914	161.	Cfb	Cropland	2004–2006	22.05	Anthoni et al. (2004)
DE-Gri	Grillenburg	50.950	13.513	385.	Cfb	Grassland	2005–2006	20.65	Gilmanov et al. (2006)
DE-Hai	Hainich	51.079	10.452	430.	Cfb	Deciduous broadleaf forest	2000–2006	19.14	Kutsch et al. (2010)
DE-Har	Hartheim	47.934	7.601	201.	Cfb	Evergreen needleleaf forest	2005–2006	23.58	Schindler et al. (2006)
DE-Kli	Klingenberg	50.893	13.523	480.	Cfb	Cropland	2004–2006	19.70	Prescher et al. (2010)
DE-Meh	Mehrstedt 1	51.275	10.656	286.	Cfb	Grassland	2003–2006	21.52	Don et al. (2009)
DE-Tha	Tharandt (Anchor Station)	50.964	13.567	380.	Cfb	Evergreen needleleaf forest	1996–2006	20.32	Grünwald et al. (2007)
DE-Wet	Wetzstein	50.453	11.458	785.	Cfb	Evergreen needleleaf forest	2002–2006	17.55	Rebmann et al. (2010)
DK-Sor	Soroe (LilleBogeskov)	55.487	11.646	40.	Cfb	Deciduous broadleaf forest	1996–2006	18.75	Pilegaard et al. (2003)
FR-Fon	Fontainebleau	48.476	2.780	90.	Cfb	Deciduous broadleaf forest	2005–2006	22.29	see <a href="http://www.fluxdata.org">www.fluxdata.org</a>
FR-Gri	Grignon	48.844	1.952	125.	Cfb	Cropland	2005–2006	22.77	Hibbard et al. (2005)
FR-Hes	Hesse Forest (Sarrebouurg)	48.674	7.065	300.	Cfb	Deciduous broadleaf forest	1997–2006	21.78	Granier et al. (2008)
FR-LBr	Le Bray	44.717	−0.769	61.	Cfb	Evergreen needleleaf forest	1996–2006	24.97	Berbigier et al. (2001)
FR-Lq1	Laqueuille	45.644	2.737	1040.	Cfb	Grassland	2003–2006	18.67	Allard et al. (2007)
FR-Lq2	Laqueuille extensive	45.639	2.737	1040.	Cfb	Grassland	2003–2006	18.67	Allard et al. (2007)
IT-Lav	Lavarone	45.955	11.281	1353.	Cfb	Evergreen needleleaf forest	2000–2006	18.37	Marcolla et al. (2003)
IT-MBo	Monte Bondone	46.016	11.047	1550.	Cfb	Grassland	2003–2006	16.29	Gianelle et al. (2009)
IT-Mal	Malga Arpaco	46.117	11.703	1730.	Cfb	Grassland	2003–2006	15.01	see <a href="http://www.fluxdata.org">www.fluxdata.org</a>
IT-Ren	Renon/Ritten	46.588	11.435	1730.	Dfb	Evergreen needleleaf forest	1999–2006	17.20	Montagnani et al. (2009)
NL-Ca1	Cabauw1	51.971	4.927	0.7	Cfb	Grassland	2003–2006	20.60	Jacobs et al. (2007)
NL-Hor	Horstermeer	52.029	5.068	−2.2	Cfb	Grassland	2004–2006	22.49	Jacobs et al. (2007)
NL-Loo	Loobos	52.168	5.744	25.	Cfb	Evergreen needleleaf forest	1996–2006	20.58	Dolman et al. (2002)
UK-PL3	Pang/Lambourne	51.45	−1.267	115.	Cfb	Deciduous broadleaf forest	2005–2006	23.02	see <a href="http://www.fluxdata.org">www.fluxdata.org</a>
UK-Ham	Hampshire	51.121	−0.861	80.	Cfb	Deciduous broadleaf forest	2004–2005	20.88	see <a href="http://www.fluxdata.org">www.fluxdata.org</a>

**Supplementary Table 1** (cont.) | Full site references.


- Allard, V., et al., The role of grazing management for the net biome productivity and greenhouse gas budget (CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub>) of semi-natural grassland, *Agr. Ecosyst. Environ.* **121**, 47–58 (2007).
- Ammann, C., Flechard, C.R., Leifeld, J., Neftel, A., & Fuhrer, J., The carbon budget of newly established temperate grassland depends on management intensity, *Agr. Ecosyst. Environ.* **121**, 5–20 (2007).
- Anthoni, P.M., et al., Forest and agricultural land-use-dependent CO<sub>2</sub> exchange in Thuringia, Germany, *Glob. Change Biol.* **10**, 2005–2019 (2004).
- Aubinet, M., Chermanne, B., Vandenhaute, M., Longdoz, B., Yernaux, M., & Laitat, E., Long term carbon dioxide exchange above a mixed forest in the Belgian Ardennes, *Agr. Forest Meteorol.* **108**, 293–315 (2001).
- Berbigier, P., Bonnefond, J.M., & Mellmann, P., CO<sub>2</sub> and water vapour fluxes for 2 years above Euroflux forest site, *Agr. Forest Meteorol.* **108**, 183–197 (2001).
- Dolman, A.J., Moors, E.J., & Elbers, J.A., The carbon uptake of a mid latitude pine forest growing on sandy soil. *Agr. Forest Meteorol.* **111**, 157–170 (2002).
- Don, A., Rebmann, C., Kolle, O., Scherer-Lorenzen, M. & Schulze E.-D., Impact of afforestation-associated management changes on the carbon balance of grassland, *Glob. Change Biol.* **15**, 1990–2002 (2009).
- Gianelle, D., Vescovo, L., Marcolla, B., Manca, G., & Cescatti, A., Ecosystem carbon fluxes and canopy spectral reflectance of a mountain meadow, *Int. J. Remote Sens.* **30**(2), 435–449 (2009).
- Gielen, B., Verbeeck, H., Neiryneck, J., Sampson, D.A., Vermeiren, F., & Janssens, I.A., Decadal water balance of a temperate Scots pine forest (*Pinus sylvestris* L.) based on measurements and modelling, *Biogeosciences* **7**, 1247–1261 (2010).
- Gilmanov, T.G., et al., Partitioning European grassland net ecosystem CO<sub>2</sub> exchange into gross primary productivity and ecosystem respiration using light response function analysis, *Agr. Ecosyst. Environ.* **121**, 93–120 (2006).
- Granier, A., Breda, N., Longdoz, B., Gross, P., & Ngao, J., Ten years of fluxes and stand growth in a young beech forest at Hesse, North-eastern France, *Ann. For. Sci.* **65**(7), 704, doi:10.1051/forest:2008052 (2008).
- Grünwald, T., & Bernhofer, Ch., A decade of carbon, water and energy flux measurements of an old spruce forest at the Anchor Station Tharandt, *Tellus* **59B**, 387–396 (2007).
- Hammerle, A., Haslwanter, A., Tappeiner, U., Cernusca, A., & Wohlfahrt G., Leaf area controls on energy partitioning of a temperate mountain grassland, *Biogeosciences* **5**, 421–431 (2008).
- Hibbard, K.A., Law, B.E., Reichstein, M., & Sulzman, J., An analysis of soil respiration across northern hemisphere temperate ecosystems, *Biogeochemistry* **73**, 29–70 (2005).
- Jacobs, C.M.J., et al., Variability of annual CO<sub>2</sub> exchange from Dutch grasslands, *Biogeosciences* **4**, 803–816 (2007).
- Kutsch, W.L., Persson, T., Schrumpf, M., Moyano, F.E., Mund, M., Andersson, S., & Schulze E.-D., Heterotrophic soil respiration and soil carbon dynamics in the deciduous Hainich forest obtained by three approaches, *Biogeochemistry*, doi:10.1007/s10533-010-9414-9 (2010).
- Marcolla, A., Pitacco, A., & Cescatti, A., Canopy architecture and turbulence structure in a coniferous forest, *Bound.-Lay. Meteorol.* **108**, 39–59 (2003).
- Montagnani, L., Manca, G., Canepa, E., Georgieva, E., Acosta, M., Feigenwinter, C., Janous, D., Kerschbaumer, G., Lindroth, A., Minach, L., Minerbi, S., Mölder, M., Pavelka, M., Seufert, G., Zeri, M., & Ziegler, W., A new mass conservation approach to the study of CO<sub>2</sub> advection in an alpine forest, *J. Geophys. Res.* **114**, D07306, doi:10.1029/2008JD010650 (2009).
- Moureaux, C., Debacq, A., Bodson, B., Heinesch, B., & Aubinet, M., Annual net ecosystem carbon exchange by a sugar beet crop, *Agr. Forest Meteorol.* **139**, 25–39 (2006).
- Pilegaard, K., Mikkelsen, T.N., Beier, C., Jensen, N.O., Ambus, P., & Ro-Poulsen, H., Field measurements of atmosphere-biosphere interactions in a Danish beech forest, *Boreal Environ. Res.* **8**, 315–333 (2003).
- Prescher, A.-K., Grünwald, T., & Bernhofer, Ch., Land use regulates carbon budgets in eastern Germany: from NEE to NBP, *Agr. Forest Meteorol.* **150**, 1016–1025, doi:10.1016/j.agrformet.2010.03.008 (2010).
- Rebmann, C., Zeri, M., Lasslop, G., Mund, M., Kolle, O., Schulze, E.-D., & Feigenwinter, Ch., Treatment and assessment of the CO<sub>2</sub>-exchange at a complex forest site in Thuringia, Germany, *Agr. Forest Meteorol.* **150**, 684–691, doi:10.1016/j.agrformet.2009.11.001 (2010).

Schindler, D., Turk, M., & Mayer, H., CO<sub>2</sub> fluxes of a Scots pine forest growing in the warm and dry southern upper Rhine plain, SW Germany, *Eur. J. Forest Res.* **125**, 201–212 (2006).


Staudt, K., & Foken, T., Documentation of reference data for the experimental areas of the Bayreuth Center for Ecology and Environmental Research (BayCEER) at the Waldstein site, in: Arb. Ergebnisse No. 35, Dep. Micrometeorol., University of Bayreuth (2007).



**Supplementary Figure 1** | Distribution of the monthly average 500 hPa height anomaly (Z500) with respect to the climatology over the years 1948–2002. **a**, June 2003. **b**, July 2003. **c**, August 2003. **d**, July 2006. Data come from the NCEP/NCAR Reanalysis and have been obtained from the Climate Explorer ([climexp.knmi.nl](http://climexp.knmi.nl)). The atmospheric circulation regime in June 2003 is dominated by a deep anomalous trough covering the North Atlantic, while heatwave conditions in July and August 2003 and July 2006 were caused by an Omega Blocking situation with an extensive high located over Northern Europe.




**Supplementary Figure 2** | Functional relationships between midday surface fluxes of sensible heat ( $H$ ), latent heat ( $\lambda ET$ ), and ground heat ( $G$ ) and environmental conditions. **a**, Incoming shortwave radiation ( $SW_{\downarrow}$ ). **b**, Daily maximum air temperature ( $T_{\max}$ , see Supplementary Table 1 for non-heatwave site averages). Curves have been derived using locally weighted polynomial regression (LOESS) on all midday data (9:00–13:00 UTC, heatwave days included) in the months June–August for all sites. Uncertainty bounds reflect 5% and 95% percentiles of the LOESS regression as determined by bootstrapping. The upper plot displays the relations per land use type separately; in the lower plot the relations are shown for  $H$  and  $\lambda ET$  separately. Relationships with radiation are stronger than with temperature as indicated by the smaller uncertainty bounds. Note that grassland and forest have different maxima in their  $T, \lambda ET$ -relationships, with forest becoming temperature limited at lower temperatures. The sensible heat flux over grassland shows no clear relation with temperature over the observed range of temperatures, whereas it strongly increases with temperature over forest.




**Supplementary Figure 3** | Relation between heatwave day (HWD) anomalies of sensible (H) and latent ( $\lambda$ ET) heat flux for forest (triangles) and grass-/cropland (circles) sites. Lines indicate levels of available energy. Site abbreviations indicate the HWD medians. The single-component Gaussian density contours reveal a distinct clustering between forest and grass-/cropland sites.





**Supplementary Figure 4** | Reduction in uncertainty of the estimated summer (June–August) climatology as a function of the number of days (D) and the number of years (Y) of the averaging period as determined by the random combination method. Data are taken from flux tower sites with the longest continuous records (Hainich, Hesse Forest, Loobos) and the Rietholz bach experimental site in Switzerland ([www.iac.ethz.ch/url/rietholz bach](http://www.iac.ethz.ch/url/rietholz bach)). Note that the exponents will converge to  $-0.50$  for uncorrelated data. The strong temporal autocorrelation makes it more effective to increasing the number of years to reduce the uncertainty than increasing the number of days around the day of interest (roughly by a factor 3). For temperature, estimating the climatology from 5 years of data and using a 15-day window reduces the uncertainty by 73% of the interannual variability of daily temperature. For fluxes, which have a lower temporal autocorrelation, this reduction is 77%. Here we use at least 2 years of data and a 15-day window, corresponding to minimum reductions of 58% and 65%, respectively.



**Supplementary Figure 5** | Observed summer soil moisture evolution for the site pairs shown in Figure 2. **a**, Cabauw and Loobos (distance 60 km). **b**, Mehrstedt and Hainich (distance 26 km). **c**, Grillenburg and Tharandt (distance 4 km). Thick lines indicate observations for the year 2006, grey lines show observations for 2004 and 2005 as reference. Arrows indicate the dates. Soil moisture observations are made at approximately 30 cm depth using Time Domain Reflectometry. Note that soil moisture levels at the peak of the 2006 heatwave were not necessary below 2004 and 2005 levels for the forest sites that show a large increase in sensible heat flux. No soil moisture observations are available for Grillenburg. The plots also illustrate the difficulty when comparing soil moisture across sites: volumetric soil moisture values can be in completely different ranges due to differences in soil texture, and the measurements represent only a small part of the total volume of soil over which plants take up their water. Moreover, the uptake of plants is a dynamic process and might not be represented well by measurements at a single depth.