

Role of Nanoplanktons in Marine food-webs

Neelava Das^{1*} Ashish Pandey²

¹Former Laboratory for Molecular Plant Physiology and Biotechnology, Department of Biology, University of Antwerp, Groenenborgerlaan, B2020, Belgium

²Alkem Health Science Ltd., Samardung, South Sikkim 737116, India

*E-mail address: neelavamicrobiologist@gmail.com

Keywords: Nanoplanktons, remineralization, top–down control, bacterial consumers

ABSTRACT

Nanoplanktons are ubiquitous protozoan zooplankton in a size range of 2 to 20 μm , play key ecological roles in aquatic ecosystems. Heterotrophic nanoflagellates are distributed through the continental shelf and margin area of the oceans as well as deep-sea. These organisms contribute significantly to the total living biomass within these systems, serve as the major top–down control on bacterial assemblages, and are an important source of mortality for microalgae and other heterotrophic nanoflagellates. From many recent studies, it is generally accepted that HNF is one of the most important bacterial consumers. They also function as important remineralizers of organic matter and nutrients in aquatic systems. In accordance with these important ecological roles, heterotrophic nanoflagellates have been the subject of considerable study both in the field and laboratory.

1. MARINE FOOD WEBS AND THE ‘MICROBIAL LOOP’

Aquatic ecosystems cover about 70% of the surface of our globe and marine environment makes up ~ 97% of the aquatic ecosystem. Marine eco-niches are characterized by diverse and contrasting physical, chemical and biological characteristics - from the shallow coastal to extreme deep-sea habitats. Diverse organisms are inhabitants of the marine ecosystem. From the smallest viruses (<0.2 microns) and bacteria (<2 microns) to single-cell marine plants called phytoplanktons (2 – 200 microns) to the biggest of mammals – the blue whales (25meter), all play a role in the sustainability as these marine organisms remain interdependent on each other through “marine food-webs”. Thus marine food-webs represents a ‘network of food chains’ or feeding relationships by which energy and nutrients are passed on from one species of living organisms to another for growth and reproduction (Fig.1) .




Fig. 1 Marine food-webs

2. STRUCTURE AND FUNCTIONING OF MARINE FOOD-CHAINS

The marine food chains are described in more quantitative terms of the ‘ecological pyramids’ or the ‘trophic prisms’ which describes the various stages within the ecological food chain based on numbers, biomass or energy. At the individual trophic levels of the prism, marine organisms can be broadly classified as the producers, consumers and decomposers, wherein matter is cycled among organisms and passed on from producers to consumers at the second trophic level and subsequently remineralized by the decomposers to be again brought back to the system for the producers (Fig. 2) and shapes the biogeochemical cycling of carbon and nutrients on our planet.




Fig. 2 Cycling of matter and energy flow through marine food chains

Thus, at the base of ecological pyramids (Fig. 3) we have the Primary Producers of “marine food-webs” which can be single-cell marine plants called “**Phytoplanktons**” in the sunlit upper layers of the ocean called the euphotic zone, or even “**chemo- autotrophic bacteria**” in deep-sea environments of hydrothermal vents.




Fig. 3 Trophic prism comparing a sunlit phytoplankton based community with a deep sea microbe based Hot Vent community

The vast area of the pelagic ocean and euphotic zone, however makes “**Phytoplanktons**” the most important primary producers of our planet. They form the basis of an intricately linked “food web”. Through process of photosynthesis and respiration “Phytoplanktons” are responsible for gaseous exchanges of carbon-dioxide and oxygen with the atmosphere.

Until a few decades ago, marine pelagic food webs were depicted as containing only three major groups of organisms: algal cells were the primary producers at the bottom of the food chain, they were grazed by mesozooplankton (0.2 mm – 20 mm in size), which in turn were grazed by fish. This is called the classic food chain or the grazing food chain and results in high productivity that supports the fisheries of the world.




Fig. 4 Classical marine food-chains

This view of a marine pelagic food chain was challenged during the 1970s and 1980s by scientists such as Pomeroy and Azam who showed that there was an alternative pathway of carbon flow that led from bacteria to protozoans (nanoflagellates and ciliates) to metazoa, with dissolved organic matter (DOM) being utilized as substrate by the bacteria. DOM can enter the pelagic environment from a variety of sources: excretion of dissolved organic carbon (DOC) by algal cells, algal cell lysis, 'sloppy grazing' by mesozooplankton or diffusion from fecal pellets. This food web paradigm was called the microbial loop (Fig. 5). This food web is thus driven by recycled carbon.




Fig. 5 Role of Microbes in the marine environment

3. MICROBIAL FOOD WEB MODEL

The microbial food web refers to the combined trophic interactions among microbes in aquatic environments. An important pathway sustaining the marine microbial food-web is the “microbial loop”. It describes the process by which bacteria can uptake dissolved organic material (DOM) that cannot be directly ingested by larger organisms. DOM includes phytoplankton photosynthates, zooplankton wastes and cytoplasm that leaks out of phytoplankton cells. Nanoflagellates and ciliates eat these marine bacteria, helping to recycle organic matter back into the marine food web. Bacteria also help to facilitate phytoplankton growth by releasing nutrients.




Fig. 6 The Microbial food-web model

A large degree of complexity exists in the interactions that can take place between the micro-organisms in pelagic food webs which includes:

- a) **Preferential Grazing** - For example, young copepod life stages and ciliates may compete for the same food sources, but adult copepods have been observed to graze preferentially on ciliates and so may give their offspring a competitive advantage.
- b) **Omnivory** - It has been discovered that omnivory is very common. For example, both copepods and ciliates can be primary and secondary consumers, with ciliates preying on autotrophic and heterotrophic flagellates as well as cyanobacteria and bacteria, and copepods grazing on phytoplankton and ciliates.

The Microbial Food web components

The microbial components of webs can be divided into two main size classes-

1. **Picoplankton** (<2 μm) including autotrophic prokaryotic and eukaryotic phytoplankton cells and heterotrophic microbes (bacteria and small flagellates).
2. **Nanoplankton** (2-20 μm) with nanophytoplankton, large flagellates and small ciliates. Flagellates and ciliates are the most important picoplanktonivorous protozoa in most aquatic Environments (McManus & Fuhrman, 1988, Sander et al; 1992).




Fig. 7 Size classes of the microbial food web components

4. Position of Nanoflagellates in Trophic Prism

The position of nanoflagellates in trophic prism is based on its nutritional status as either autotrophs or heterotrophs. Thus nanoflagellates are residing under either in the producer or primary consumer. So their numbers also vary for their different type of grazing strategy and energy utilization. Like this way autotrophic nanoflagellates are coming under the producer rather heterotrophic nanoflagellates are grouped as the primary consumer.

5. Role of Nanoflagellates and ciliates in the Microbial Food web

Heterotrophic Bacteria remains at the base of the Microbial food-web. The extent of role of bacteria in an ecosystem to act as only "remineralizers" of nutrients or as direct nutritional source for higher trophic levels depends on several factors controlling their production and abundance. Nanoflagellates are reported bacteriovores and hence control their abundance and production. Nanoflagellates can also graze on the phototrophic picoplanktons and Ciliates can graze on both bacteria and small flagellates. Thus, the role of Nanoplanktons (Nanoflagellates and ciliates) is important to understand the pattern of the energy flow through microbial loop to higher zooplanktons and fisheries. Due to the scarcity of diagnostic morphological features, the destruction of delicate forms by fixation and the selectivity of culturing efforts we have little knowledge about the diversity, autecology and biogeography of heterotrophic nanoflagellates (Lim *et al.* 1999). Chryomonads or chrysophyceans (class Chrysophyceae) are phototrophic and/or heterotrophic nanoflagellates that comprise a major component of the aquatic food web in both marine and freshwater systems e.g. colorless chryomonads together with bicosoecids constitute between 20 - 50% of the annual average biomass of pelagic HNF (Arndt *et al.* 2000).

6. Current Status and Significance:

The population genetics study revealed that the importance of heterotrophic nanoflagellates related to climate change, therefore in Polar Regions, heterotrophic nanoflagellates are major consumers of bacteria and contribute significantly to the carbon flux from DOC via bacteria to larger organisms such as ciliates and metazoans. New research explored that the growth rates of *P. imperforate*, a heterotrophic nanoflagellate increased significantly with increasing temperature in Polar Regions (Lee and Fenchel 1972). It has been examined that growth rates of cultured heterotrophic nanoflagellates from permanently cold environments (Lee and Fenchel, 1972; Choi and Peters, 1992; Mayes *et al.*, 1997). Now a days culture of the heterotrophic nanoflagellate species have been examined to determine the effects of a wide range of physical and chemical parameters on growth rate and other physiological and biogeochemical processes.

5. CONCLUSION

Several studies have shown that Heterotrophic bacteria consume dissolved organic material (DOM) originating from phytoplankton photosynthetic activity, converting it to particulate organic material (POC) (Azam *et al.*, 1983). Studies have shown that heterotrophic bacteria can consume

10–40% of primary production in the eastern subarctic Pacific (Kirchman *et al.*, 1993) and North Atlantic (Ducklow *et al.*, 1993; Li *et al.*, 1993). Heterotrophic bacteria are known to be fed primarily by heterotrophic nano- flagellates (HNF), which in turn are cropped mainly by microzooplankton (Sherr and Sherr, 1988; Weisse and Scheffel-Möser, 1991). Thus, in addition to micro-zooplankton herbivory, HNF bacterivory can be regarded as a fundamental process in controlling carbon flow through microbial food webs in the ocean.

References

- [1] Adrian, R. 1991. Filtering and feeding rates of cyclopoid copepods feeding on phytoplankton. *Hydrobiologia* 210: 217-223.
- [2] Azam F, Fenchel T, Field JG, Gray JS, Meyer-Reil LA & Thingstad F (1983) The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series* 10: 257–263
- [3] Ballintine, J. 1953. Comparison of the different methods of estimating nanoplankton. *J. Mar. Biol. Assoc. U. K.* 32: 129-147
- [4] Barsdate, R. J., T. Fenchel and R. T. Prentki. 1974. Phosphorous cycle of a model ecosystem: significance for decomposer food chains and effect of bacterial grazers. *Oikos*. 25: 239-251
- [5] Beaver, J.R. and T.L. Crisman. 1989. The role of ciliated protozoa in pelagic freshwater ecosystems. *Microbial Ecology* 17: 111-136.
- [6] Christoffersen K, B. Riemann, A. Klysner, and M. Sondergaard. 1993. Potential role of fish predation and natural populations of zooplankton in structuring a plankton community in eutrophic lake water. *Limnology and Oceanography* 38: 561-573.
- [7] Christoffersen, K., B. Riemann, L.R. Hansen, A. Klysner, and H.B. Sorensen 1990. Qualitative importance of the microbial loop and plankton community structure in a eutrophic lake during a bloom of cyanobacteria. *Microbial Ecology* 20: 253-272.
- [8] Cole JJ, Findlay S & Pace MJ (1988) Bacterial production in fresh and saltwater ecosystems: a cross-system overview. *Marine Ecology Progress Series* 43: 1–10
- [9] Cole, J.J. and N.F. Caraco. 1993. The pelagic microbial food web of oligotrophic lakes., p. 101-111. *In* T.E. Ford [ed.], *Aquatic microbiology. An ecological approach*. Blackwell Scientific Publications
- [10] Daggett, P., and T. A. Nerad. 1982. Axenic cultivation of *Bodo edax* and *Bodo ancinatus* and observations on feeding rate in monoaxenic culture. *Abst.30. J. Protozool.* 29: 290-291.
- [11] Davis, P. G. and J. McN. Sieburth. 1982. Differentiation of phototrophic and heterotrophic nanoplankton populations in marine waters by epifluorescent microscopy. *Annl. Inst. Oceanogr., Paris* 58(S): 249-260.
- [12] Dussart, G. 1965. Les differentes categories de plancton. *Hydrobiologia* 26: 72- 74. Estep, K., P. G. Davis, M. D. Keller and J. McN Sieburth. 1986. How important are algal nanoflagellates in bacterivory. *Limnol. Oceanogr.* 31: 646-650.
- [13] Fenchel, T. 1982a. Ecology of heterotrophic microflagellates I. Some important forms. *Mar. Ecol. Prog. Ser.* 8: 211-233.
- [14] Fenchel, T. 1982b. Ecology of heterotrophic microflagellates II. Bioenergetics and growth. *Mar. Ecol. Prog. Ser.* 8: 225-231.
- [15] Fenchel, T. 1982c. Ecology of heterotrophic microflagellates II. Adaptations to heterogeneous environment. *Mar. Ecol. Prog. Ser.* 9: 25-33

- [16] Fenchel, T. 1982d. Ecology of heterotrophic microflagellates IV. Quantitative, importance and occurrence as bacterial consumers. *Mar. Ecol. Prog. Ser.* 9: 35-42.
- [17] Finlay K, Roff JC (2004) Radiotracer determination of the diet of calanoid copepod nauplii and copepodites in a temperate estuary. *Ices Journal of Marine Science* 61:552-562.
- [19] Gasol JM (1994) A framework for the assessment of top-down vs bottom-up control of heterotrophic nanoflagellate abundance. *Mar Ecol Prog Ser* 113: 291- 300.
- [20] Gasol JM, Simons AM, Kalff J (1995) Patterns in the top-down versus bottom- up regulation of heterotrophic nanoflagellates in temperate lakes. *J Plankton Res* 17: 1879-1903.
- [21] Goldman, J. C. and D. A. Caron. 1985. Experimental studies on an omnivorous microflagellate: implications for grazing and nutrient regeneration in the marine microbial food chain. *Deep Sea Res.* 32: 899-915
- [22] Goldman, J. C., D. A. Caron, O. Ketil Andersen and M. R. Dennett. 1985. Nutrient cycling in a microflagellate food chain I. Nitrogen dynamics. *Mar. Ecol. Prog. Ser.* 24: 231-242.
- [23] Gude H. 1979. Grazing by protozoa as a selection factor for activated sludge. *Microb. Ecol.* 5: 225-237.
- [24] Haas L. W. and K. L. Webb. 1979. Nutritional mode of several non-pigmented microflagellates from the York river estuary, Virginia. *J. Exp. Mar. Biol.Ecol.* 39: 125-134.
- [25] Hilliard, D. K. 1971. Notes on the occurrence and taxonomy of some planktonic Chrysophytes in an Alaskan lake, with comments on the genus *Bicoeca*. *Arch. Protistenk.* D.113: 98-122.
- [26] Hobbie, J. E., R. J. Daley and S. Jasper. 1977. Use of Nuclepore filters for counting bacteria by fluorescent microscopy. *Appl. Environ. Microbiol.* 33: 1225- 1228.
- [27] Jansson M, Bergström A-K, Blomqvist P, Isaksson A & Jonsson A (1999) Impact of allochthonous organic carbon on microbial food web carbon dynamics and structure in Lake Öträsket. *Archiv für Hydrobiologie* 144: 409-428
- [28] Jurgens, K., S.A. Wickham, K.O. Rothhaupt, and B. Santer. 1996. Feeding rates of macro- and microzooplankton on heterotrophic nanoflagellates. *Limnology and Oceanography* 41: 1833-1839.
- [29] Kopylov, A. I. and E. F. Moiseev. 1980. Effect of colourless flagellates on the determination of bacterial production in seawater. *Hydrobiology* 252:503-505
- [30] Kopylov, A. I., A. F. Pasternak and Y. V. Moiseev. 1981. Consumption of flagellates by planktonic organisms. *Oceanology* 2 1: 269-271
- [32] Landry M. R. and R. P. Hassett. 1982. Estimating the grazing impact of marine microzooplankton. *Mar. Biol.* 67: 283-288
- [33] Leadbeater, B. S. C. and C. Morton. 1974. A microscopical study of a marine species of *Codnosia* James -Clark (Choanoflagellate) with special reference to the ingestion of bacteria. *Bot. J. Linn. Soc.* 6: 337-347.
- [34] Levine, N. D et al. 1980. A Newly Revised Classification of Protozoa. *J. Protozool.* 27: 37-58.
- [35] Lighthart, B. 1969. Planktonic and benthic bacterivorous protozoa at 11 stations in the Puget Sound and adjacent Pacific Ocean. *J. Fish. Res. Bd. Can.* 26:299-306.
- [36] Perry, J.J., J.T. Staley, and S. Lory. 2002. *Microbial life*. Sinauer Associates, Inc. Reuman DC, Cohen JE (2004) Trophic links' length and slope in the Tuesday Lake food web with species' body mass and numerical abundance *J Anim Ecol* 73: 852-866
- [37] Rassoulzadegan F, Lavalpeuto M, Sheldon RW (1988) Partitioning of the food ration of marine ciliates between picoplankton and nanoplankton. *Hydrobiologia* 159: 75-88

-
- [38] Rassoulzadegan F, Sheldon RW (1986) Predator-prey interactions of nanozooplankton and bacteria in an oligotrophic marine-environment *Limnol Oceanogr* 31: 1010-1021
- [39] Sherr EB, Sherr BF (2002) Significance of predation by protists in aquatic microbial food webs. *Antonie Van Leeuwenhoek* 81: 293-308
- [40] Sieburth J McN, Davis PG (1982) The role of heterotrophic nanoplankton in the grazing and nurturing of planktonic bacteria in the Sargasso and Caribbean seas. *Ann Inst Oceanogr Paris* 58: 285-296.
- [41] Sieburth, J. McN. and P. G. Davis. 1982. The role .of heterotrophic nanoplankton in the grazing and nurturing of planktonic bacteria in the Sargasso and Caribbean Seas.
- [42] *Annls. Inst. Oceanogr. Paris.* 58(S): 285-296.
- [43] Stockner JG & Porter KG (1988) Microbial food webs in freshwater planktonic ecosystems. *In* *Complex Interactions in Lake Communities* (Ed Carpenter SR), Springer Verlag, New York, pp69–84.

(Received 18 July 2015; accepted 22 July 2015)

Volume 43

10.18052/www.scipress.com/ILNS.43

Role of Nanoplanktons in Marine Food-Webs

10.18052/www.scipress.com/ILNS.43.38