Short Communication

Trend analysis suggested a change in subspecies among *Mycobacterium avium* isolated from pigs in Belgium, 1967–2013

K. Soetaert, C. Vluggen, L. Duytschaever, J. Denoël, V. Roupie, F. Smeets, N. Bruffaerts, K. Huygen, D. Fretin, M. Diels, L. Rigouts, C. Saegerman, V. Mathys

Mycobacterium avium is one of the most frequently isolated non-tuberculous mycobacteria, responsible of human and porcine infections. The species *M. avium* consists of four subspecies; *M. avium* subspecies *avium* (MAA), *M. avium* subspecies *silvaticum* (MAS), *M. avium* subspecies *paratuberculosis* (MAP) and *M. avium* subspecies *hominissuis* (MAH) (Thorel and others 1990, Mijls and others 2002, Turenne and others 2007). Given its pathogenicity for both human beings and pigs, MAH represents an increasing public health concern (Pate and others 2011, Iwamoto and others 2012, Muwonge and others 2014). In pigs, these mycobacteria provoke granulomatous lesions mainly in lymph nodes

of the digestive tract, potentially reducing the value of carcasses (Alvarez and others 2011).

Recently, the authors reported on the predominance of MAH among Belgian patients (98.9 per cent) and slaughter pigs (100 per cent) infected by *M. avium* (Vluggen and others 2016). Genotyping of these isolates revealed a large genetic diversity and the absence of a link between genotypes and the place of residence (human) or the farm of origin (pigs) suggesting an environmental source of infection (Vluggen and others 2016).

As porcine *M. avium* isolates collected in Belgium during an earlier period of time (1967–1968 and 1992–1996) were available in the Belgian Co-ordinated collections of micro-organisms (BCCM)/Institute of Tropical Medicine (ITM) collection, the authors performed subspecies determination and genotyping on this panel in order to compare the results with those from the more recent isolates sampled in 2012–2013 (Vluggen and others 2016), to investigate the evolution of *M. avium* subspecies distribution over time in infected domestic pigs.

All 37 porcine *M. avium* isolates available from the collection maintained at BCCM/ITM (Belgium) were used for this study. This panel comprised isolates sampled, isolated and provided by the Belgian veterinary centre Het Centrum voor Onderzoek in Diergeneeskunde en Agrochemie – Centre d'Étude et de Recherches Vétérinaires et Agrochimiques (CODA-CERVA) during the periods 1967–1968 (n=12) and 1992–1996 (n=25).

The *M. avium* species identification of the selected strains was confirmed by PCR detection of the insertion sequence (IS) 1245 (Guerrero and others 1995, Vluggen and others 2016) and the *M. avium* subspecies characterisation was determined by analysing the presence/absence of IS901 (Godfroid and others 2005, Vluggen and others 2016). Genotyping of the *M. avium* isolates was performed by multispacer sequence typing (MST) as described previously (Cayrou and others 2010, Vluggen and others 2016). The MST types were determined by consulting the MST database (Méditerranée infection 2015). Results were compared with those recently reported on a panel of current Belgian porcine *M. avium* isolates (n=52) (Vluggen and others 2016).

Based on the presence of IS1245 (specific for *M. avium* strains) and the presence/absence of IS901 (present in MAA/MAS but absent in MAH/MAP), 11 isolates among the 37 analysed were identified as MAH and 26 as MAA/MAS (Table 1).

Overall, 10 different MST types were identified; 5 among MAH and 5 among MAA (Table 1), with MST type 12 being the most prevalent among MAH (6/11) and MST types 2 and 3 the most prevalent in MAA isolates (14/26 and 6/26, respectively). The authors identified four new MST types which were recorded in the MST database developed by the 'Université de la Méditerranée' as MST type 117, 118, new 1 and new 2.

Comparison of MST types from porcine isolates obtained in the 1960s–1990s with those sampled in 2012–2013, revealed the presence of both MAA (n=26, 70.3 per cent) and MAH (n=11, 29.7 per cent) subspecies in the old panels versus only MAH isolates in the recent panel, and a change is observed in MST type over time (Table 2).

The present study was undertaken to determine the evolution of *M. avium* subspecies infecting Belgian pigs over time. Unlike the omnipresence of the subspecies *hominissuis* reported among Belgian pigs infected by *M. avium* in 2012–2013, the authors observed in their older panels of isolates the presence of both species, MAA and MAH. Moreover, the MST type changed over time.

In Europe, various mycobacteria have been recently reported from pigs at slaughterhouses, with 11 per cent of the isolates belonging to the *M. avium* complex (MAC) (EFSA 2011). Mycobacteria from MAC are ubiquitous bacteria that can be found in water, food, soil and other environmental samples and

Veterinary Record (2017)

doi: 10.1136/vr.103951

K. Soetaert, Lic,
C. Vluggen, Lic,
V. Mathys, PhD,
Bacterial Diseases Service, Operational
Direction Communicable and Infectious
Diseases, Scientific Institute of Public
Health (WIV-ISP), Brussels, Belgium
L. Duytschaever, PhD,
J. Denoël, DMV,
F. Smeets, Tech,
C. Saegerman, DMV, MSc Epid, PhD,
Prof., DipIECVPH,
Research Unit for Epidemiology and
Risk Analysis applied to Veterinary
Sciences (UREAR-ULg), Fundamental
and Applied Research, Animal and
Health (FARAH) Center, Liège, Belgium
V. Roupie, PhD,
D. Fretin, PhD,
Bacterial Zoonoses of Livestock Unit,
Operational Direction Bacterial
Diseases, Veterinary and Agrochemical
Research Centre (CODA-CERVA),
Brussels, Belgium

L. Duytschaever, is also at Bacterial
Zoonoses of Livestock Unit, Operational

Direction Bacterial Diseases, Veterinary
and Agrochemical Research Centre
(CODA-CERVA), Brussels, Belgium

N. Bruffaerts, PhD,
K. Huygen, PhD,
Immunology Service, Operational
Direction Communicable and
infectious Diseases, Scientific Institute
of Public Health (WIV-ISP), Brussels,
Belgium

M. Diels, Tech,
L. Rigouts, PhD, Prof.,
BCCM/ITM Mycobacteria Collection,
Mycobacteriology Unit, Institute of
Tropical Medicine, Antwerp, Belgium

L. Rigouts, is also at Department
Biomedical Sciences, University of
Antwerp, Antwerp, Belgium

KS and CV contributed equally.

Email for correspondence:
vmathys@wiv-isp.be

Provenance: Not commissioned;
externally peer reviewed

Accepted February 16, 2017

Short Communication

TABLE 1: Distribution of *Mycobacterium avium* subspecies and their MST of 37 Belgian porcine isolates sampled in the years 1967–1968 and 1992–1996

Subspecies identification based on the IS profiles and MST types	IS1245/IS901 profile	Isolates sampled in 1967–1968	Isolates sampled in 1992–1996	Total
<i>M avium</i> subspecies <i>hominissuis</i>	+/-	2 2 MST type 12	9 4 MST type 12 2 MST type 13 1 MST type 15 1 MST type 118 (new) 1 MST New 2	11
<i>M avium</i> subspecies <i>avium/silvaticum</i>	+/+	10 3 MST type 2 4 MST type 3 3 MST type 117 (new)	16 11 MST type 2 2 MST type 3 2 MST type 4 1 MST New 1	26
Total		12	25	37

—, absence of the fragment; +, presence of the fragment; IS, insertion sequence; MST, multispacer sequence typing

TABLE 2: MST types identified among Belgian porcine *Mycobacterium avium* isolates in this study (1967–1968 and 1992–1996 sample) in comparison with the study from [Vluggen and others \(2016\)](#) (1) (2012–2013 sample)

<i>M avium</i> subspecies	MST type	1967–1968 panel	1992–96 panel	2012–2013 panel	Total
MAA	2	3	11		14
	3	4	2		6
	4		2		2
	117 (new)	3			3
MAH	New 1		1		1
	12	2	4	28	34
	13		2		2
	15		1		1
	118 (new)		1		1
	New 2		1		1
	20			2	2
	22			7	7
	30			1	1
	51			2	2
	85			3	3
	91			1	1
	102			2	2
	103			1	1
	104			3	3
	109			1	1
	110			1	1
Total		12	25	52	89

MAA, *Mycobacterium avium* subspecies *avium*; MAH, *Mycobacterium avium* subspecies *hominissuis*; MST, multispacer sequence typing

are considered opportunistic pathogens for numerous animal species, mainly birds and pigs, as well as for human beings ([Radomski and others 2010](#)). In particular, MAA is the aetiological agent of tuberculosis in birds, including wood pigeons ([Mijs and others 2002](#)). One hypothesis to explain the decrease of MAA in domestic pigs over time is the increase of biosafety measures designed to reduce the risk of introduction, establishment and spread of animal diseases, and their gradual implementation in the pig sector after the epidemic of classical swine fever in the beginning of the 1990s ([Arrêté royal 1953](#), [Vanthemsche and Sagerman 1994](#)). These measures probably induced the decrease of entrance of birds in the piggeries.

In The Netherlands, a decrease in prevalence of MAA infections was observed after the arrest of compost use in piggeries ([Komijn and others 2007](#)). However, in Belgium, the compost is still being used ([Arrêté royal 2003](#)).

Strains with particular IS profiles (absence of IS1245 usually present in *M avium* strains or false PCR detection of IS901 provoked by the presence of ISMav6) have been reported in different

studies realised on contemporary strains ([Ichikawa and others 2009](#), [Vluggen and others 2016](#)). No such strains were found among the 1960's sample. Although limited by the small sample size of the present study, this observation suggests that these variants may have appeared more recently.

During the authors' 2012–2013 study, MST 12 was the most frequently isolated MAH genotype, among Belgian porcine samples, as well as the human samples ([Vluggen and others 2016](#)), suggesting a common source of infection ([Vluggen and others 2016](#)). However, in total 11 different MST types were identified among pigs, underlining the large genetic diversity of MAH isolates.

The main limitations of the present report are the small number of *M avium* isolates available in the collection and the lack of information concerning their origin. Although it is known that the strains were isolated from Belgian pigs and analysed in different Belgian laboratories, information about the farms of origin or the reason for sample collection are not available.

In conclusion, the authors report here a temporal difference in the subspecies distribution among porcine *M avium* isolates between the years 1967–1968/1992–1996 and years 2012–2013 in Belgium. MAH seems to have relatively increased at the expense of MAA, and MST types changed over time, although MST12 remained predominant among MAH isolates. The disappearance of MAA isolates could be the result of increased biosafety measures gradually put in place in the pig sector since the 1990s. Analysis of a larger and well characterised sample would be required to confirm the observations reported here.

Acknowledgements

Gene sequencing was performed at the Platform Biotechnology and Molecular Biology at the Scientific Institute of Public Health (WIV-ISP).

Funding This study was funded by the Federal Public Service, Health, Safety of the Food chain and Environment (convention RT 12/5 Lymphindic 1, Contractual Research, FPS Public Health, Security of the Food Chain and Environment). CV is a PhD student funded by the RP/PJ WIV-ISP (TYPTUB). The National Reference Centre for Tuberculosis & Mycobacteria is partially supported by the Belgian Ministry of Social Affairs through a fund within the Health Insurance System. MD is funded by the Belgian Science Policy (Belspo).

References

- ALVAREZ, J., CASTELLANOS, E., ROMERO, B., ARANAZ, A., BEZOS, J., RODRIGUEZ, S., MATEOS, A., DOMINGUEZ, L. & DE, J. L. (2011) Epidemiological investigation of a *Mycobacterium avium* subsp. *hominissuis* outbreak in swine. *Epidemiology and Infection* **139**, 143–148.
- Arrêté royal du 14 juin (1953) déterminant les conditions d'équipement pour la détention des porcs. 2 juillet 1993. Moniteur Belge
- Arrêté royal du 15 mai (2003) relatif à la protection des porcs dans les élevages porcins. 24 juin 2003. Moniteur Belge
- CAYROU, C., TURENNE, C., BEHR, M. A. & DRANCOURT, M. (2010) Genotyping of *Mycobacterium avium* complex organisms using multispacer sequence typing. *Microbiology (Reading, England)* **156**, 687–694.
- EFSA. European Food Safety Authority (2011) Technical specifications on harmonised epidemiological indicators for public health hazards to be covered by meat inspection of swine. *EFSA Journal* **9**, 2371.
- GODFROID, J., DELCORPS, C., IRENGE, L. M., WALRAVENS, K., MARCHE, S. & GALA, J. L. (2005) Definitive differentiation between single and mixed mycobacterial infections in red deer (*Cervus elaphus*) by a combination of duplex amplification of p34 and f57 sequences and Hpy188I enzymatic restriction of duplex amplicons. *Journal of Clinical Microbiology* **43**, 4640–4648.
- GUERRERO, C., BERNASCONI, C., BURKI, D., BODMER, T. & TELENTI, A. (1995) A novel insertion element from *Mycobacterium avium*, IS1245, is a specific target for analysis of strain relatedness. *Journal of Clinical Microbiology* **33**, 304–307.
- ICHIKAWA, K., YAGI, T., MORIYAMA, M., INAGAKI, T., NAKAGAWA, T., UCHIYA, K., NIKAI, T. & OGAWA, K. (2009) Characterization of *Mycobacterium avium* clinical isolates in Japan using subspecies-specific insertion sequences, and identification of a new insertion sequence, ISMav6. *Journal of medical microbiology* **58**, 945–950.
- IWAMOTO, T., NAKAJIMA, C., NISHIUCHI, Y., KATO, T., YOSHIDA, S., NAKANISHI, N., TAMARU, A., TAMURA, Y., SUZUKI, Y. & NASU, M. (2012) Genetic diversity of *Mycobacterium avium* subsp. *hominissuis* strains isolated from humans, pigs, and human living environment. *Infection, Genetics and*

- Evolution : Journal of Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases* **12**, 846–852
- KOMIJN, R. E., WISSELINK, H. J., RIJSMAN, V. M., STOCKHOFF-ZURWIJEDEN, N., BAKKER, D., VAN ZIJDERVELD, F. G., EGER, T., WAGENAAR, J. A., PUTIRULAN, F. F. & URLINGS, B. A. (2007) Granulomatous lesions in lymph nodes of slaughter pigs bacteriologically negative for *Mycobacterium avium* subsp. *avium* and positive for *Rhodococcus equi*. *Veterinary Microbiology* **120**, 352–357
- Méditerranée infection (2015) MST *Mycobacterium avium* Marseille, France. http://iffr48timone.univ-mrs.fr/mst/mycobacterium_avium/groups.html
- MIJS, W., DE HAAS P., ROSSAU, R., VAN DER LAAN, T., RIGOUTS, L., PORTAELS, F. & VAN S OOLINGEN D. (2002) Molecular evidence to support a proposal to reserve the designation *Mycobacterium avium* subsp. *avium* for bird-type isolates and '*M. avium* subsp. *hominissuis*' for the human/porcine type of *M. avium*. *International Journal of Systematic and Evolutionary Microbiology* **52**, 1505–1518
- MUWONGE, A., OLOYA, J., KANKYA, C., NIELSEN, S., GODFROID, J., SKJERVE, E., DJONNE, B. & JOHANSEN, T. B. (2014) Molecular characterization of *Mycobacterium avium* subspecies *hominissuis* isolated from humans, cattle and pigs in the Uganda cattle corridor using VNTR analysis. *Infection, Genetics and Evolution: Journal of Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases* **21**, 184–191
- PATE, M., KUSAR, D., ZOLNIR-DOVC, M. & OCEPEK, M. (2011) MIRU-VNTR typing of *Mycobacterium avium* in animals and humans: heterogeneity of *Mycobacterium avium* subsp. *hominissuis* versus homogeneity of *Mycobacterium avium* subsp. *avium* strains. *Research in Veterinary Science* **91**, 376–381
- RADOMSKI, N., THIBAUT, V. C., KAROU, C., DE CRUZ K., COCHARD, T., GUTIERREZ, C., SUPPLY, P., BIET, F. & BOSCHIROLI, M. L. (2010) Determination of genotypic diversity of *Mycobacterium avium* subspecies from human and animal origins by mycobacterial interspersed repetitive-unit-variable-number tandem-repeat and IS1311 restriction fragment length polymorphism typing methods. *Journal of Clinical Microbiology* **48**, 1026–1034
- THOREL, M. E., KRICHEVSKY, M. & LEVY-FREBAULT, V. V. (1990) Numerical taxonomy of mycobactin-dependent mycobacteria, emended description of *Mycobacterium avium*, and description of *Mycobacterium avium* subsp. *avium* subsp. nov., *Mycobacterium avium* subsp. *paratuberculosis* subsp. nov., and *Mycobacterium avium* subsp. *silvaticum* subsp. nov. *International Journal of Systematic Bacteriology* **40**, 254–260
- TURENNE, C. Y., WALLACE, R. Jr. & BEHR, M. A. (2007) *Mycobacterium avium* in the postgenomic era. *Clinical Microbiology Reviews* **20**, 205–229
- VANTHEMSCHE, P. & SAEGERMAN, C. (1994) Gestion d'un épisode de peste porcine classique en Belgique. *Epidémiologie en santé animale* **26**, 29–41
- VLUGGEN, C., SOETAERT, K., DUYTSCHAEVER, L., DENOEL, J., FAUVILLE-DUFAUX, M., SMEETS, E., BRUFFAERTS, N., HUYGEN, K., FRETIN, D., RIGOUTS, L., SAEGERMAN, C. & MATHYS, V. (2016) Genotyping and strain distribution of *Mycobacterium avium* subspecies *hominissuis* isolated from humans and pigs in Belgium, 2011–2013. *Euro Surveill* **21**, 30111

Trend analysis suggested a change in subspecies among *Mycobacterium avium* isolated from pigs in Belgium, 1967–2013

K. Soetaert, C. Vluggen, L. Duytschaever, J. Denoël, V. Roupie, F. Smeets, N. Bruffaerts, K. Huygen, D. Fretin, M. Diels, L. Rigouts, C. Saegerman and V. Mathys

Veterinary Record 2017 180: 449 originally published online March 10, 2017

doi: 10.1136/vr.103951

Updated information and services can be found at:

<http://veterinaryrecord.bmj.com/content/180/18/449.1>

These include:

References

This article cites 16 articles, 4 of which you can access for free at:

<http://veterinaryrecord.bmj.com/content/180/18/449.1#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>