This item is the archived peer-reviewed author-version of:

In vitro diagnosis of immediate drug hypersensitivity during anesthesia : a review of the literature

Reference:

Ebo Didier, Faber Margaretha, Elst Jessy, Van Gasse Athina, Bridts Christiaan, Mertens Christel, De Clerck Luc S., Hagendorens Margo, Sabato Vito.- In vitro diagnosis of immediate drug hypersensitivity during anesthesia : a review of the literature
The journal of allergy and clinical immunology. In practice - ISSN 2213-2198 - Amsterdam, Elsevier science bv, 6:4(2018), p. 1176-1184
Full text (Publisher's DOI): <https://doi.org/10.1016/J.JAIP.2018.01.004>
To cite this reference: <https://hdl.handle.net/10067/1495010151162165141>

***In vitro* diagnosis of immediate drug hypersensitivity during anaesthesia: a review of the literature**

Ebo Didier G, MD, PhD^{1,*}, Faber Margaretha, MD, PhD¹, Elst Jessy, MSc¹, Van Gasse Athina L, MD^{1,2}, Bridts Chris H, MLT¹, Mertens Christel, MLT¹, De Clerck Luc S, MD, PhD¹, Hagendorens Margo M, MD, PhD^{1,2}, Sabato Vito, MD, PhD¹.

¹ Faculty of Medicine and Health Science, Department of Immunology – Allergology – Rheumatology, University of Antwerp, Antwerp University Hospital.

² Faculty of Medicine and Health Science, Department of Paediatrics, University of Antwerp, Antwerp University Hospital.

* Correspondence to:

Ebo Didier, MD, PhD

Laboratory of Immunology University of Antwerp

CDE T5.95 Universiteitsplein 1

2610 Antwerpen

Belgium

Tel: ++ 32 (0) 3 2652595

Fax: ++ 32 (0) 3 2652655

immuno@uantwerpen.be

Key words: allergy, anaesthesia, basophil activation, drugs, flow cytometry, immediate drug hypersensitivity reaction (IDHR), specific IgE (sIgE), beta-lactam, penicillin, MRGPRX2, neuromuscular blocking agents (NMBA), opiates, tryptase

The authors declare not having a conflict of interest nor financial support.

List of abbreviations:

α -gal: galactose- α (1,3)-galactose

BAT: basophil activation test

DPT: drug provocation test

Fc ϵ RI: high affinity receptor for sIgE

Hev b: *Hevea brasiliensis*

IHR: immediate hypersensitivity reactions

NMBA: neuromuscular blocking agents

PAPPC: p-aminophenyl phosphoryl choline

sIgE: specific IgE antibodies

THIQ: tetrahydroisoquinoline

Abstract

Quantification of specific IgE (sIgE) antibodies constitutes an important measure to document anaesthesia-related immediate hypersensitivity reactions (IHR). However, only a few drug-specific assays are available and their predictive value is not known. In cases of non-IgE-mediated IHR, diagnosis might benefit from cellular tests such as basophil mediator release tests and basophil activation tests (BAT).

To review the potential and limitations of quantification of sIgE, mediator release and BAT in anaesthesia-related IHR a literature search was conducted using the key-words allergy, basophil activation, CD63, CD203c, diagnosis, drugs, hypersensitivity, flow cytometry, MRGPRX2, specific IgE antibodies, leukotrienes, histamine and tryptase; this was complemented by the authors' experience.

The drugs and compounds that have predominantly been studied are neuromuscular blocking agents (NMBA), β -lactams, latex and chlorhexidine. For sIgE, sensitivity and specificity varies between 38.5-92% and 92-100% for NMBA respectively and between 0-85% and 52-100% for

52 β -lactams. Specific IgE to morphine should not be used in isolation to diagnose IHR to NMBA
53 nor opiates. sIgE for latex, and in difficult cases molecular diagnosis with quantification of sIgE
54 to *Hevea* components constitute reliable diagnostics.
55 For drugs, sensitivity of BAT varies between 50 and 60%, specificity reaches 80-90%. Basophil
56 mediator release tests seem to be abandoned and supplanted by BAT.
57

Introduction

The gold standard to ascertain correct diagnosis of immediate hypersensitivity reactions (IHR) to drugs is a controlled drug provocation test (DPT) with the culprit compound(s). However, DPTs entail a risk of severe, life-threatening complications and can be contraindicated (e.g., patients having suffered from life-threatening reactions) or impossible (e.g., full-dose DPT in hypersensitivity to curarizing neuromuscular blocking agents (NMBA)). Moreover, the predictive value of DPTs is not known and DPTs might yield false negative results¹. Therefore, the diagnostic approach of anaesthesia-related IgE-mediated IHR generally starts with history taking, thorough review of the anaesthetic/surgical notes complemented with skin testing and/or *in vitro* quantification of specific IgE (sIgE) antibodies. However, only a few drug-sIgE assays are available, and most of them have not been clinically validated. Furthermore, IHR might not *per se* involve IgE/FcεRI-cross-linking, but may also result from alternative pathways such as an off-target occupation of the Mas-related G-protein receptor MRGPRX2^{2, 3} that cannot be detected by a sIgE antibody assay. The development and validation of cellular tests such as basophil activation tests (BAT) would, to some extent, be promising in such cases.

The objective of this article is to review the literature on the value of serum tryptase, histamine, commercially available drug-sIgE assays and basophil activation tests such as mediator release tests and BAT in the diagnosis of anaesthesia-related IHR. Emphasis is put on some misconceptions, shortcomings, and unmet needs. As with any subject still beset by many questions, alternative interpretations, hypotheses, or explanations expressed here may not find universal acceptance.

Quantification of serum tryptase

Although quantification of peak and baseline serum tryptase do not contribute to the identification of the culprit, serum tryptase has proven to be extremely valuable in diagnosing anaesthesia-related IHR, mainly to confirm mast cell degranulation and/or to rule out or confirm (clonal) mast cell disorders ⁴ and mast cell activation syndromes ⁵. Currently, in the commercially available assay, total tryptase is quantified as the sum of continuously secreted baseline tryptase and β -tryptase released from degranulating mast cells (ImmunoCAP Thermofisher, Uppsala, Sweden). Since relevant increases have been observed way below the traditional decision threshold of 11.4 $\mu\text{g/L}$, it has been suggested to abandon this cut-off ⁶⁻⁸. For example, an incremental threshold of 20% was shown to identify potential mast cell mediator release in an additional 14% of cases with peak tryptase between 5 and 14 $\mu\text{g/L}$ and a further 15% with peak tryptase below 5 $\mu\text{g/L}$. Others have proposed that an increase of tryptase over baseline (24h after the acute event) levels is clinically relevant when it exceeds $2 + 1.2 \times \text{baseline}$ ^{9, 10}. Although in the study by Sprung *et al.*, quantification of peak tryptase was performed between 30 minutes and 4 hours from the event, it is recommend to take the peak sample as close to 60 minutes after the reaction as possible, and if not possible later samples should still be taken and compared with a baseline taken at a later date ¹¹. Alternatively, by comparing the two measurements, anaphylaxis could be ruled out even for acute tryptase values $> 11.4 \mu\text{g/L}$ in cases of baseline hypertryptasemia ⁹. Quantifying baseline tryptase has another additional purpose, as elevated baseline levels might be indicative for underlying (clonal) mast cell disorders ⁴ that might underlie severe IHR, particularly in men who do not demonstrate urticaria/angioedema ¹². Levels of β -tryptase $> 1 \mu\text{g/L}$ indicate mast cell degranulation. However, this test is not commercially available. Quantification of plasma histamine, although highly sensitive, was inferior to quantification of serum tryptase for

discrimination between IgE-dependent and IgE-independent anaesthesia-related IHR ¹³. Resuscitation manoeuvres by themselves appear not to modify mediator concentrations ¹⁴. Alternatively, it is important to stress that an elevated peak tryptase measurement does not necessarily indicate mast cell activation ^{9, 15}. In chronic renal failure elevated “peak” serum tryptase ¹⁵ might result from mast cell hyperplasia due to slow elimination of stem cell factor ¹⁶. Note that tryptase is not cleared by the kidneys ¹⁷. Tryptase can also be elevated in critically ill patients without anaphylaxis ¹⁸ and victims of trauma ¹⁹. False negative results mainly result from incorrect sampling time (ideally 60-90 minutes after onset of symptoms).

Principles of quantification of drug specific IgE (sIgE) antibodies and BAT

Like tissue resident mast cells, basophils can be triggered in IgE-dependent and various IgE-independent ways. Cross-linking of the surface-bound high-affinity IgE receptor (FcεRI) generally occurs through (glyco)proteins, chemical allergens or auto-antibodies directed against the FcεRI receptor or membrane-bound IgE antibodies. Quantification of sIgE antibodies predominantly relies upon quantification of a drug-(hapten)-carrier antibody complex in which the secondary antihuman IgE is conjugated to an enzyme with colorimetric reading in the enzyme linked immunosorbent test (ELISA) or with a fluorescence reading in the fluorescent enzyme immunoassay (FEIA) ²⁰. However, only a limited number of drug-specific sIgE immunoassays are available and most of these assays have not been thoroughly validated, mainly as a result of the unavailability of sufficient numbers of patients and exposed or challenged control individuals.

An IgE-independent, activation will mainly result from coupling of surface receptors with endogenous (*e.g.* cytokines, anaphylatoxins, chemokines, IgG, neuropeptides) or exogenous (*e.g.* pathogen associated molecular patterns) elements. Amongst these receptors is the Mas-

related G protein receptor MRGPRX2 that can lead to a quick but rather transient mast cell degranulation ²¹ and appears to be involved in different mast cell-associated conditions including non-immune immediate drug hypersensitivity reactions ^{22,23}. Recently, McNeil *et al.* ² described the potential of MRGPRX2-related mast cell activation by various drugs containing a tetrahydroisoquinoline (THIQ) motif such as some fluoroquinolones and various NMBA. The MRGPRX2 receptor has subsequently also been incriminated in reaction towards opioids ²⁴ and vancomycin ²⁵. Alternatively, other largely unknown pathways might also induce degranulation.

The foundations of current flow-assisted BAT were laid 25 years ago ²⁶ and in the meantime the technique has largely supplanted older mediator release assays that rely upon difficult quantification of mediators released in the supernatant. Actually, the last reviews on mediator release tests date back to 2003 ^{27, 28}. To our knowledge, since then no large case-control studies including more than 15 patients and significant numbers of (exposed) control individuals on the application of the various mediator release tests have been published, except one report including patients who suffered from peri-operative hypersensitivity reactions resulting from various causes ²⁹.

Traditional BAT relies upon a flow cytometric analysis of various activation and degranulation markers on the surface membrane. These changes can be detected and quantified on a single-cell level using specific monoclonal antibodies conjugated with different LASER-excitabile fluorochromes. The technical principles and requirements of BAT have been detailed elsewhere ³⁰. Basophils are traditionally identified by markers such as CCR3 (CD193)/CD3, CD123/HLA-DR or IgE/CD203c. Of these markers, only CD203c, is lineage specific. After activation, the appearance and/or up-regulation of surface activation and/or degranulation markers, such as CD203c and/or CD63 is quantified. For a review on the applications and

limitations of the BAT in drug IHR see Mangoldt et al ³¹. Histamine release can also be quantified by flow cytometry ³² and the technique is applicable in IHR to drugs ³³.

β-lactams

The most studied sIgE assays are those for β-lactams, especially amoxicillin and benzyl penicilloyl. Although, several cases of positive sIgE results in IHR with negative skin tests have been described ³⁴⁻³⁸, sIgE assays for β-lactams, as shown in table 1, generally exhibit a poor sensitivity that decreases over time ³⁹. Besides these disappointing sensitivity data, there is increasing evidence supporting low specificity of the tests ^{35, 37, 40-43}. In some studies false positivity could have resulted from nonspecific binding in the solid phase assay as a result of elevated total IgE titres ⁴¹⁻⁴⁴. An alternative explanation for false-positive sIgE penicillin seems to be sIgE antibodies to phenylethylamine that test negative in a BAT ⁴⁴. In summary, sIgE antibodies to β-lactams seem of restricted value and should ideally not be used in isolation to exclude or confirm IHR to these antibiotics. In order to avoid misdiagnosis, these assays should be complemented with BAT, skin testing and, where appropriate a DPT ^{45, 46}. Table 2 summarizes the data of BAT in IHR to β-lactams. Up to now, 10 studies have investigated the BAT as a diagnostic in IHR to β-lactams, mainly to amoxicillin. Compared with the quantification of sIgE antibodies, BAT shows a higher sensitivity (about 50%) and specificity (approximately 90%). As for specific IgE, sensitivity of BAT to β-lactams is rather low and decreases over time but both tests can remain positive for years ^{39, 47}. Therefore, we cannot adhere to the recommendation by ENDA/EAACI drug allergy interest group not to perform drug-sIgE tests after three years ⁴⁸, particularly as also skin test responsiveness decreases over time ^{39, 49}. Finally, it should be kept in mind that sensitisation to amoxicillin/clavulanate can also result from sensitization to clavulanic acid ⁵⁰, which needs specific testing ⁵¹. For a diagnostic algorithm for IHR to β-lactam antibiotic the reader is referred elsewhere ⁵².

Neuromuscular blocking agents (NMBA)

In many countries, curarizing neuromuscular blocking agents (NMBA) represent a significant cause of anaesthesia-related anaphylaxis⁵³⁻⁵⁷. As indicated in our diagnostic algorithm for NMBA (figure 1), skin tests are the primary instrument to confirm IHR to NMBA and cannot be substituted by BAT or quantification of sIgE⁵⁸. However, the predictive value of skin testing is not absolute thereby leaving room for additional *in vitro* tests. In the absence of readily available assays, for about 2 decades, several groups have tried to define the accuracy of various home-made NMBA-sIgE assays (table 3)⁵⁹⁻⁶². At present, IHR to NMBA are serologically assessed indirectly through assays measuring IgE reactivity to tertiary and quaternary substituted ammonium structures that have been shown to be the major epitopes of NMBA^{63, 64}. Most frequently applied methods are a choline chloride^{59, 60, 65-70}, a p-aminophenyl phosphoryl choline (PAPPC)^{59, 65, 66, 71} and/or morphine-based assays^{59-61, 71-76}. With respect to the ImmunoCAP FEIA for suxamethonium, rocuronium, atracurium and morphine the sensitivity and specificity for the individual NMBA-specific varies between 38.5-92% and 85.7-100%, respectively. Furthermore, it has been demonstrated that a morphine-based immuno assay is a valuable test to detect suxamethonium and rocuronium-reactive antibodies but not to depict atracurium-reactive antibodies^{72, 74}. Quantifying IgE reactivity to tertiary and quaternary substituted ammonium structures to identify patients at risk or to document NMBA hypersensitivity^{77, 78} might cause a large number of false positive results as they are prevalent in the general population^{73, 74, 76}. Therefore, these assays cannot be recommended as a screening tool to identify patients at risk or to document NMBA hypersensitivity^{77, 78}. For example, Leysen *et al.*⁷⁸, showed that a positive sIgE to morphine is not a contraindication for administration of atracurium and cisatracurium in patients with negative skin tests to these benzyliisoquinolines. Explanations for these false-positive sIgE results are an elevated total IgE

⁷⁴ and intake of the opiate antitussive pholcodine ⁷⁹. Specific IgE antibodies to tertiary and quaternary substituted ammonium epitopes can remain positive for several years after the acute reaction ^{77, 80}. Alternatively, as recently stressed by Spoerl *et al.* ³, IHR to NMBA such as rocuronium might occur independently from IgE/FcεRI cross-linking and relate to MRGPRX2-mediated activation of mast cells ² and, therefore, not be detected by sIgE assays.

Table 4 displays the data about BAT in IHR to NMBA. In general, sensitivity of the assay varies between 36 and 92%, whereas specificity reaches 95%. Importantly, BAT not only enables identification of the culprit drug but also provides the opportunity to study cross-reactivity and identify safe alternatives for future anaesthesia ^{81, 82}. Alternatively, as resting basophils barely express MRGPRX2, it is unlikely that traditional BAT using cells in steady state be of value in reactions occurring from off-target occupation of this receptor ⁸³. Our current policy is to offer BAT and sIgE to patients with negative or equivocal skin tests who suffered from severe anaphylaxis and in whom no alternative cause was demonstrable.

Opiates and (semi-)synthetic opioids

Genuine IgE-mediated allergies to opiates (morphine, codeine) and semi-synthetic opioid pholcodine remain rare notwithstanding their frequent and universal use. Additionally, correct diagnosis is not straightforward, mainly because of uncertainties associated with measurement of drug-specific IgE antibodies and skin testing ⁸⁴. Recently, it has been suggested that the two commercially available sIgE assays for a *Papaver somniferum* (poppy seed) extract and morphine can add to the diagnosis of IgE-mediated opiate allergy ^{85, 86}.

However, using DPT we were unable to confirm these data ⁸⁷, mainly because of the high prevalence of sIgE antibodies to these compounds in an allergic population. This observation is highly relevant when facing patients for whom correct identification of the causative compound(s) is impeded because of simultaneous intake or administration of different agents,

e.g., during general anaesthesia. Erroneous opiate allergy diagnosis might not only entail unnecessary avoidance measures but also, most importantly, ultimately put patients at risk by overlooking alternative diagnoses such as an allergy to rocuronium or suxamethonium. For the time being the sole *in vitro* method to document opiate allergy is the basophil activation test, as basophils from opiate tolerant individuals, unlike their cutaneous mast cells, are unresponsive to opiates^{87, 88}. Moreover, negative BAT, along with negative skin testing for different NMBA and negative provocation tests for the structurally almost similar opiates suggest that these drugs are probably safe in pholcodine hypersensitivity⁸⁸. Therefore, for opiates and semi-synthetic opioids we recommend to start with BAT, and, if negative, a DPT. In contrast, for synthetic opioids such as fentanyl, sufentanyl and remifentanyl diagnosis can start with skin testing eventually complemented with BAT and/or DPT for difficult cases^{84, 89}.

Chlorhexidine

Chlorhexidine, a cationic bisguanide antiseptic and disinfectant, is used as the (di)acetate or (di)glucuronide salt. These chlorhexidine salts can trigger irritant dermatitis, allergic contact dermatitis⁹⁰, IDHR (including life-threatening anaphylaxis)⁹¹⁻⁹⁴ and even a combination of both, contact dermatitis and IDHR⁹⁵. For a traditional arbitrarily chosen decision threshold of 0.35 kUA/L, the sensitivity of sIgE chlorhexidine varied between 84.2-91.6% and specificity between 93.7-100%. For a ROC-generated threshold of 0.20 kUA/L sensitivity was 94.1% and specificity 90.7%^{93, 94}. As for β -lactam⁴²⁻⁴⁴ and NMBA⁷⁴, raised total IgE levels were shown to have an impact on chlorhexidine sIgE measurement at levels higher than 500 kU/L and more particularly at levels higher than 2,000 kU/L⁹⁴. Recently, it was demonstrated the optimal sampling time for sIgE chlorhexidine was between 1 and 4 months⁹⁶, but sIgE might persist for years⁵⁵.

Latex

Another significant cause of anaesthesia-related anaphylaxis is *Hevea latex* (*Hevea brasiliensis*; *Hev b*). Diagnosis of latex allergy mainly relies upon skin testing and/or various in vitro tests. However, correct diagnosis of IgE-mediated allergy to natural latex is not always straightforward, mainly because of the false-positive sIgE results^{97, 98}, especially in patients suffering from grass and weed pollen allergy who are sensitized to cross-reactive carbohydrate determinants and/or profilin *Hev b* 8⁹⁹⁻¹⁰³. Therefore, in a significant number of patients additional tests such as skin tests, component resolved diagnosis¹⁰⁰⁻¹⁰³ and eventually BAT^{98, 104-107} might be required to establish correct diagnosis. For a review on component resolved diagnosis the reader is referred elsewhere¹⁰⁸.

Other agents

Bovine gelatine constitutes the active component in certain plasma substitutes, haemostatic sponges and can be present in various other drugs such as vaccines. Since the first descriptions of the allergenicity of gelatine¹⁰⁹, IgE-mediated IHR to this compound, including fatal anaphylaxis, have been increasingly reported. Today, 2 distinct types of IgE-mediated bovine gelatine allergy are recognized. First, genuine gelatine allergy that results from sensitization to the protein part of the molecule. Second, gelatine allergy resulting from a sensitization to a glycan moiety of the molecule, *i.e.* galactose- α (1,3)-galactose (α -gal)¹¹⁰⁻¹¹², as first described by Chung *et al.*¹¹³ and Commins *et al.*¹¹⁴. To our knowledge, there are no studies that have determined the diagnostic accuracy of sIgE gelatine. However, it is of note that patients with life-threatening anaphylaxis to gelatine as a result of α -gal sensitization are generally overlooked by traditional gelatine-sIgE assay and need additional testing including quantification of α -gal specific IgE antibodies and gelatine skin testing¹¹⁰⁻¹¹².

Discussion

270 Correct management of anaphylaxis during anaesthesia requires a multidisciplinary approach
271 with prompt recognition and treatment of the attending anaesthesiologist, quantification of
272 peak tryptase, and subsequent determination of the responsible agent(s) with strict avoidance
273 of all incriminated and cross-reactive compounds. From this review it emerges that diagnosis
274 of anaphylaxis during anaesthesia is not always straight forward. It can be hindered as a broad
275 spectrum of different drugs can elicit heterogeneous immune and non-immune
276 hypersensitivity reactions. Problems are certainly compounded as multiple drugs are
277 administered simultaneously and non-anaesthesia related drugs or compounds (e.g.
278 premedication, disinfection, antibiotic rinsing by surgeon, lymph node mapping) can also be
279 the cause of an IHR. Nevertheless, diagnostic work-up should be offered to all patients with a
280 clinical suspicion of anaesthesia-related IHR, irrespective of the grade of severity. Evaluation
281 should comprise all agents the patient was exposed to and it should be kept in mind that a
282 patient might demonstrate multiple sensitizations. In an own survey of over 650 patients
283 double sensitization occurs in approximately 8% of the cases (unpublished data).

284 From this review it appears that drug, chlorhexidine and latex-sIgE antibody testing can
285 provide useful information but can rarely be applied as solitary diagnostic tests to exclude or
286 document IHR, as they lack 100 percent predictive values. For β -lactam determinants the main
287 issue is poor sensitivity, which could not be increased without significant loss of specificity ⁴².
288 For NMBA, drug-specific IgE tests seem to attain acceptable sensitivity and specificity,
289 provided the application of drug-specific cut-offs ^{74, 94}. Although quantification of sIgE to
290 morphine appears to be a reliable biomarker of sensitization to tertiary and quaternary
291 ammonium structures, IgE reactivity to this compound in general and allergic population is as
292 high as 5-10%. Therefore, the test should not be applied in isolation to diagnose IHR to NMBA
293 or opiates. With respect to the unsatisfactory sensitivity of some tests it has been argued that

294 this observation relates to the time-interval elapsed between the acute reaction and testing.
295 Although we agree that late testing can result in lower sensitivity we do not adhere to the
296 recommendation of the ENDA/EAACI Drug Allergy Interest Group. Based upon a single
297 publication about negativation of sIgE to β -lactams ³⁹, further use of drug-sIgE is dissuaded
298 when the time-interval exceeds 3 years in their Position Paper ⁴⁸. However, this is not our
299 experience ⁷⁷ and drug-sIgE may persist as long as 5-30 years ^{49, 80}. With respect to the low
300 specificity of some tests it is reemphasized that correct interpretation of sIgE results requires
301 taking into account total IgE values ^{42, 74, 94}. Whether the introduction of sIgE/total IgE ratio's
302 increases specificity ⁴² remains to be confirmed. We currently do not apply these ratio's.

303 Since the earliest days of the BAT it was obvious that this technique would become an asset
304 in the allergological diagnostic instrumentation to document IHR, particularly when diagnosis
305 cannot be established by other means. Moreover, it is anticipated that BAT using
306 "conditioned" basophils expressing the MRGPRX2 receptor might become an easy accessible
307 instrument to study and diagnose IgE/Fc ϵ RI-independent IDHR that result from off-target
308 occupation of this receptor. However, additional collaborative large-scale studies are needed
309 to verify whether the BAT fulfils this promise, to optimize and harmonize the protocols, to
310 avoid instigation of cynicism and scepticism, and to enable and justify its entrance in routine
311 diagnostic application.

313 **Acknowledgement**

314 EDG is a Senior Clinical Researcher of the FWO (1800614N), SV is a Senior Clinical Researcher
315 of the FWO (1804518N), VGAL is a fellow of the FWO (1113617N).

317 **Conflict of interest**

318 The authors declare no conflict of interest related to this publication

Table 1: Specific IgE to β -lactams						
Compound	Reference test	Assay	Sensitivity	Specificity	N	Reference
Various β -lactams	H + ST	CAP-FEIA	BPO + AXO + peni G + AMP: 31.8%	BPO + AXO + peni G + AMP: 88.6%	58	¹¹⁵
Various β -lactams	H $\pm$ ST $\pm$ DPT	CAP-FEIA	BPO: 32% AXO: 43% BPO+AXO: 50%	BPO: 98% AXO: 98% BPO+AXO: 96%	129	¹¹⁶
Various β -lactams	H $\pm$ ST $\pm$ DPT	CAP-FEIA	BPO: 10-68% AXO: 41-53%	BPO: 98% AXO: 95%	410	³⁸
Various β -lactams	H	CAP-FEIA	37.9%	86.7%	58	¹¹⁷
Various β -lactams ¹	H $\pm$ ST $\pm$ DPT	CAP-FEIA RAST ²	0-25% ² 42.9-75% ²	83.3-100% ² 66.7-83.3% ²	45	³⁴
Various β -lactams	H $\pm$ ST	CAP-FEIA CAP-FEIA	85% ³ 44% ⁴	54% ³ 80% ⁴	176	⁴²
Various β -lactams	H $\pm$ ST	CAP-FEIA	66%	52%	293	⁴³
Amoxicillin	H + ST + DPT	CAP-FEIA	19%	NA	57	⁵¹
¹ home-made assay, ² sensitivity and specificity vary according to clinical manifestations, ³ for a threshold of 0.10 kUA/L, ⁴ for a threshold of 0.35 kUA/L. H: history, ST: skin test, DPT: drug provocation test, N: number CAP-FEIA: fluorescence enzyme immunoassay available from Phadia Thermofisher. RAST: radio allerge sorbent test. Peni G: penicillin G, AMP: ampicillin, BPO: benzyl penicilloyl, AXO: amoxicillin Note that there is no IgE for clavulanic acid available						

Table 2: BAT in immediate β-lactam hypersensitivity						
Stimulus	Reference test	Activation marker	Sensitivity (%)	Specificity (%)	Number of patients and controls	Ref.
β-lactam	H	CD63	50	93	88	117
β-lactam	H + DPT ¹	CD63	39	93	53	118
β-lactam	H ± ST ± IgE ± DPT	CD63	49	91	110	119
Amoxicillin	H ± ST	CD203c	52	100	41	120
		CD63	22	79		
β-lactam	H	CD63	50	89-97	262	121
β-lactam	H ± ST ± IgE	CD63-CCR3	55	100	39	122
		CD63-IgE	53			
Amoxicillin	H	CD63	29	/	14 patients, no controls	123
Amoxicillin	H ± ST ± DPT	CD63	50	/	61 patients, number of controls not mentioned	124
Amoxicillin	H ± ST	CD63	50	/	30 patients	125
Amoxicillin Clavulanic acid	H ± ST ± DPT	CD63	47	93	57	51
			62	89	58	
Cefazolin	H + ST	CD63	33	94	16 patients,	47
		CD203c	67	94	17 controls	
For legend see table 1.						

Table 3: Specific IgE to NMBA and substituted ammonium structures						
Compound	Reference test	Assay	Sensitivity	Specificity	N	Reference
Various NMBA	H + ST	RIA RIA RIA	PAPPC: 97% MOR: 83% QAS: 86%	PAPPC: 97% NA NA	75	⁵⁹
Various NMBA	H + ST	RIA RAST RAST	QAS: 87.9% SUC: 66.7% Alcuronium: 40.7%	NA	83	⁶⁰
Various NMBA	H + ST	RIA RIA	MOR: 85% NMBA-specific: 52%	98%	118	⁷²
Various NMBA	H + ST	CAP-FEIA CAP-FEIA	SUXA: 38.5% MOR: 67.7%	SUX: 96.3-99.6% MOR: 90-95%	866	⁷³
Rocuronium ¹	H + ST	CAP-FEIA	SUXA: 72% ² SUXA: 60% ³ ROCU: 92% ² ROCU: 68% ³ MOR: 88% PHOL: 86%	SUXA: 100% ² SUXA: 100% ³ ROCU: 93% ² ROCU: 93% ³ MOR: 100% PHOL: 100%	82	⁷⁴
Rocuronium	H + 2 tests*	CAP-FEIA	ROCU: 83%	ROCU: 72%	66	⁷⁷
Various NMBA ¹	H + ST	CAP-FEIA	QAM ⁴ : 87.7%	QAM ⁴ : 90.7%	168	⁷⁶
Atracurium ¹	H + ST	CAP-FEIA	SUXA: 28.6% ATRA: 57.1% MOR: 14.2%	SUXA: 85.7% ATRA: 100% MOR: 85.7%	78	⁷⁵

* For validation purposes, diagnosis of rocuronium allergy was considered definite when at least 2 out of 3 tests (skin test, BAT, sIgE) were positive, as rocuronium challenges at full dose are not possible for obvious reasons.

¹ applying ROC-generated drug-specific thresholds, ² for a ROC-generated threshold of 0.11 kUA/L for suxamethonium and 0.13 kUA/L for rocuronium, ³ for a traditional threshold of 0.35 kUA/L.

⁴ "optimized" morphine-based assay

H: history, ST: skin tests, RIA: radio immunoassay, RAST: radio allergosorbent test, CAP-FEIA: fluorescence enzyme immunoassay available from Phadia Thermofisher, PAPPC: p-aminophenyl phosphoryl choline, MOR: morphine, QAS: quaternary ammonium structure, SUC: succinyl choline, SUX: suxamethonium, ROCU: rocuronium, QAM: quaternary ammonium morphine, ATRA: atracurium. N: number. NA: not available.

Table 4: BAT in immediate neuromuscular blocking agent (NMBA) hypersensitivity						
Stimulus	Reference Test	Activation marker	Sensitivity (%)	Specificity (%)	N	Ref.
Various NMBA	H	CD63	64	81	26	126
		CD45	43	96		
Various NMBA	H ± ST	CD63	54	100	56	127
Various NMBA	H	CD63	79	100	31	128
		CD203c	36	100		
Various NMBA	H ± ST	CD63	36-86 ¹	93	92	129
Rocuronium	H ± ST	CD63	92 ²	100	22	81
Various NMBA	H ± ST ± IgE	CD63	60	100	49	130
Rocuronium	H + 2 tests*	CD63	80	96	104	77
Various NMBA	H+ST	CD63	68	100	56	131
Atracurium	H ± ST	CD63	71 ³	100	75	82
<p>* For validation purposes, diagnosis of rocuronium allergy was considered definite when at least 2 out of 3 tests (skin test, BAT, sIgE) were positive, as rocuronium challenges at full dose are not possible for obvious reasons.</p> <p>NMBA: neuromuscular blocking agent, H: history, ST: skin test, N: number of patients and control individuals</p> <p>¹ Increasing sensitivity when only the reactions that occurred during the 3 years were taken into account,</p> <p>² taking into account the non-responders sensitivity is 76%,</p> <p>³ taking into account the non-responders sensitivity is 63%.</p>						

Figure 1: diagnostic algorithm NMBA

Legend of the figure:

NMBA: neuromuscular blocking agent

BAT: basophil activation test, sIgE: specific IgE

PPV: positive predictive value, NPV: negative predictive. Values are provided for rocuronium ⁷⁷ using a ROC-calculated drug-specific decision threshold of 0.13 kUA/L ⁷⁴.

¹ drug specific IgE available for suxamethonium, rocuronium and atracurium and should be using drug-specific thresholds. Specific IgE morphine is applied as a biomarker for sensitization to aminosteroids and suxamethonium (not useful for benzylisoquinolines).

² challenging at maximum 1/10th of therapeutic dose.

References

1. Demoly P, Romano A, Botelho C, Bousquet-Rouanet L, Gaeta F, Silva R, et al. Determining the negative predictive value of provocation tests with beta-lactams. *Allergy* 2010; 65:327-32.
2. McNeil BD, Pundir P, Meeker S, Han L, Undem BJ, Kulka M, et al. Identification of a mast-cell-specific receptor crucial for pseudo-allergic drug reactions. *Nature* 2015; 519:237-41.
3. Spoerl D, D'Incau S, Roux-Lombard P, Harr T, Czarnetzki C. Non-IgE-Dependent Hypersensitivity to Rocuronium Reversed by Sugammadex: Report of Three Cases and Hypothesis on the Underlying Mechanism. *Int Arch Allergy Immunol* 2016; 169:256-62.
4. Bonadonna P, Pagani M, Aberer W, Bilo MB, Brockow K, Oude Elberink H, et al. Drug hypersensitivity in clonal mast cell disorders: ENDA/EAACI position paper. *Allergy* 2015; 70:755-63.
5. Akin C. Mast cell activation syndromes presenting as anaphylaxis. *Immunol Allergy Clin North Am* 2015; 35:277-85.
6. Borer-Reinhold M, Haeberli G, Bitzenhofer M, Jandus P, Hausmann O, Fricker M, et al. An increase in serum tryptase even below 11.4 ng/mL may indicate a mast cell-mediated hypersensitivity reaction: a prospective study in Hymenoptera venom allergic patients. *Clin Exp Allergy* 2011; 41:1777-83.
7. Garvey LH, Bech B, Mosbech H, Kroigaard M, Belhage B, Husum B, et al. Effect of general anesthesia and orthopedic surgery on serum tryptase. *Anesthesiology* 2010; 112:1184-9.
8. Egner W, Sargur R, Shrimpton A, York M, Green K. A 17-year experience in perioperative anaphylaxis 1998-2015: harmonizing optimal detection of mast cell mediator release. *Clin Exp Allergy* 2016.
9. Sprung J, Weingarten TN, Schwartz LB. Presence or absence of elevated acute total serum tryptase by itself is not a definitive marker for an allergic reaction. *Anesthesiology* 2015; 122:713-4.
10. Baretto RL, Beck S, Heslegrave J, Melchior C, Mohamed O, Ekbote A, et al. Validation of international consensus equation for acute serum total tryptase in mast cell activation: A perioperative perspective. *Allergy* 2017.
11. Schwartz LB. Diagnostic value of tryptase in anaphylaxis and mastocytosis. *Immunol Allergy Clin North Am* 2006; 26:451-63.
12. Alvarez-Twose I, Gonzalez-de-Olano D, Sanchez-Munoz L, Matito A, Jara-Acevedo M, Teodosio C, et al. Validation of the REMA score for predicting mast cell clonality and systemic mastocytosis in patients with systemic mast cell activation symptoms. *Int Arch Allergy Immunol* 2012; 157:275-80.
13. Berroa F, Lafuente A, Javaloyes G, Ferrer M, Moncada R, Goikoetxea MJ, et al. The usefulness of plasma histamine and different tryptase cut-off points in the diagnosis of perianaesthetic hypersensitivity reactions. *Clin Exp Allergy* 2014; 44:270-7.
14. Laroche D, Gomis P, Gallimidi E, Malinovsky JM, Mertes PM. Diagnostic value of histamine and tryptase concentrations in severe anaphylaxis with shock or cardiac arrest during anesthesia. *Anesthesiology* 2014; 121:272-9.
15. Sprung J, Larson KJ, Divekar RD, Butterfield JH, Schwartz LB, Weingarten TN. Refractory intraoperative hypotension with elevated serum tryptase. *Asia Pac Allergy* 2015; 5:47-50.
16. Kitoh T, Ishikawa H, Ishii T, Nakagawa S. Elevated SCF levels in the serum of patients with chronic renal failure. *Br J Haematol* 1998; 102:1151-6.
17. Simon MR, Jan M, Yee J, Nori US, Hu J, Akin C, et al. Tryptase is not cleared by the kidneys into the urine. *Int Arch Allergy Immunol* 2010; 152:28-31.
18. Francis A, Fatovich DM, Arendts G, Macdonald SP, Bosio E, Nagree Y, et al. Serum mast cell tryptase measurements: Sensitivity and specificity for a diagnosis of anaphylaxis in emergency department patients with shock or hypoxaemia. *Emerg Med Australas* 2017.

19. Edston E, van Hage-Hamsten M. Mast cell tryptase and hemolysis after trauma. *Forensic Sci Int* 2003; 131:8-13.
20. Hamilton RG. Clinical laboratory assessment of immediate-type hypersensitivity. *J Allergy Clin Immunol* 2010; 125:S284-96.
21. Gaudenzio N, Sibilano R, Marichal T, Starkl P, Reber LL, Cenac N, et al. Different activation signals induce distinct mast cell degranulation strategies. *J Clin Invest* 2016; 126:3981-98.
22. Ali H. Mas-related G protein coupled receptor-X2: A potential new target for modulating mast cell-mediated allergic and inflammatory diseases. *J Immunobiol* 2016; 1.
23. Subramanian H, Gupta K, Ali H. Roles of Mas-related G protein-coupled receptor X2 on mast cell-mediated host defense, pseudoallergic drug reactions, and chronic inflammatory diseases. *J Allergy Clin Immunol* 2016; 138:700-10.
24. Lansu K, Karpiak J, Liu J, Huang XP, McCorvy JD, Kroeze WK, et al. In silico design of novel probes for the atypical opioid receptor MRGPRX2. *Nat Chem Biol* 2017; 13:529-36.
25. Azimi E, Reddy VB, Lerner EA. Brief communication: MRGPRX2, atopic dermatitis and red man syndrome. *Itch (Phila)* 2017; 2.
26. Knol EF, Mul FP, Jansen H, Calafat J, Roos D. Monitoring human basophil activation via CD63 monoclonal antibody 435. *J Allergy Clin Immunol* 1991; 88:328-38.
27. Demoly P, Lebel B, Arnoux B. Allergen-induced mediator release tests. *Allergy* 2003; 58:553-8.
28. de Weck AL, Sanz ML. Cellular allergen stimulation test (CAST) 2003, a review. *J Investig Allergol Clin Immunol* 2004; 14:253-73.
29. Xin X, Zou Y, Xing L, Yin J, Gu J, Wang Z, et al. Investigation of drugs responsible for perioperative anaphylactic reactions using cellular allergen stimulation test. *Chin Med J (Engl)* 2014; 127:3738-43.
30. Bridts CH, Sabato V, Mertens C, Hagendorens MM, De Clerck LS, Ebo DG. Flow cytometric allergy diagnosis: basophil activation techniques. *Methods Mol Biol* 2014; 1192:147-59.
31. Mangodt EA, Van Gasse AL, Decuyper I, Uyttebroek A, Faber MA, Sabato V, et al. In vitro Diagnosis of Immediate Drug Hypersensitivity: Should We Go with the Flow. *Int Arch Allergy Immunol* 2015; 168:3-12.
32. Ebo DG, Bridts CH, Mertens CH, Hagendorens MM, Stevens WJ, De Clerck LS. Analyzing histamine release by flow cytometry (HistaFlow): a novel instrument to study the degranulation patterns of basophils. *J Immunol Methods* 2012; 375:30-8.
33. Cop N, Uyttebroek AP, Sabato V, Bridts CH, De Clerck LS, Ebo DG. Flow cytometric analysis of drug-Induced basophil histamine release. *Cytometry B Clin Cytom* 2015.
34. Fontaine C, Mayorga C, Bousquet PJ, Arnoux B, Torres MJ, Blanca M, et al. Relevance of the determination of serum-specific IgE antibodies in the diagnosis of immediate beta-lactam allergy. *Allergy* 2007; 62:47-52.
35. Silva R, Cruz L, Botelho C, Castro E, Cadinha S, Castel-Branco MG, et al. Immediate hypersensitivity to penicillins with negative skin tests--the value of specific IgE. *Eur Ann Allergy Clin Immunol* 2009; 41:117-9.
36. Qiao HL, Li Z, Yang J, Tian X, Gao N, Jia LJ. Hypersensitivity reactions to penicillins: studies in a group of patients with negative benzylpenicillin G skin test. *J Clin Pharm Ther* 2009; 34:249-54.
37. Hjortlund J, Mortz CG, Skov PS, Bindslev-Jensen C. Diagnosis of penicillin allergy revisited: the value of case history, skin testing, specific IgE and prolonged challenge. *Allergy* 2013; 68:1057-64.
38. Torres MJ, Romano A, Mayorga C, Moya MC, Guzman AE, Reche M, et al. Diagnostic evaluation of a large group of patients with immediate allergy to penicillins: the role of skin testing. *Allergy* 2001; 56:850-6.
39. Fernandez TD, Torres MJ, Blanca-Lopez N, Rodriguez-Bada JL, Gomez E, Canto G, et al. Negativization rates of IgE radioimmunoassay and basophil activation test in immediate reactions to penicillins. *Allergy* 2009; 64:242-8.

40. Macy E, Goldberg B, Poon KY. Use of commercial anti-penicillin IgE fluorometric enzyme immunoassays to diagnose penicillin allergy. *Ann Allergy Asthma Immunol* 2010; 105:136-41.
41. Aberer W, Zidarn M, Kranke B. IgE antibodies to penicillin are indicative for but not conclusive proof of penicillin allergy. *Br J Dermatol* 2006; 154:1209-10.
42. Vultaggio A, Matucci A, Virgili G, Rossi O, Fili L, Parronchi P, et al. Influence of total serum IgE levels on the in vitro detection of beta-lactams-specific IgE antibodies. *Clin Exp Allergy* 2009; 39:838-44.
43. Vultaggio A, Virgili G, Gaeta F, Romano A, Maggi E, Matucci A. High serum beta-lactams specific/total IgE ratio is associated with immediate reactions to beta-lactams antibiotics. *PLoS One* 2015; 10:e0121857.
44. Johansson SG, Adedoyin J, van Hage M, Gronneberg R, Nopp A. False-positive penicillin immunoassay: an unnoticed common problem. *J Allergy Clin Immunol* 2013; 132:235-7.
45. Macy E. Penicillin allergy: optimizing diagnostic protocols, public health implications, and future research needs. *Curr Opin Allergy Clin Immunol* 2015; 15:308-13.
46. Mirakian R, Leech SC, Krishna MT, Richter AG, Huber PA, Farooque S, et al. Management of allergy to penicillins and other beta-lactams. *Clin Exp Allergy* 2015; 45:300-27.
47. Uyttebroek AP, Sabato V, Cop N, Decuyper, II, Faber MA, Bridts CH, et al. Diagnosing cefazolin hypersensitivity: Lessons from dual-labeling flow cytometry. *J Allergy Clin Immunol Pract* 2016; 4:1243-5.
48. Mayorga C, Celik G, Rouzaire P, Whitaker P, Bonadonna P, Cernadas JR, et al. In vitro tests for Drug Hypersensitivity Reactions. An ENDA/EAACI Drug Allergy Interest Group Position Paper. *Allergy* 2016.
49. Romano A, Gaeta F, Valluzzi RL, Zaffiro A, Caruso C, Quarantino D. Natural evolution of skin-test sensitivity in patients with IgE-mediated hypersensitivity to cephalosporins. *Allergy* 2014; 69:806-9.
50. Salas M, Laguna JJ, Dona I, Barrionuevo E, Fernandez-Santamaria R, Ariza A, et al. Patients Taking Amoxicillin-Clavulanic Can Become Simultaneously Sensitized to Both Drugs. *J Allergy Clin Immunol Pract* 2017; 5:694-702.e3.
51. Salas M, Fernandez-Santamaria R, Mayorga C, Barrionuevo E, Ariza A, Posadas T, et al. Use of the Basophil Activation Test May Reduce the Need for Drug Provocation in Amoxicillin-Clavulanic Allergy. *J Allergy Clin Immunol Pract* 2017.
52. Torres MJ, Blanca M, Fernandez J, Romano A, Weck A, Aberer W, et al. Diagnosis of immediate allergic reactions to beta-lactam antibiotics. *Allergy* 2003; 58:961-72.
53. Mertes PM, Alla F, Trechot P, Auroy Y, Jouglu E. Anaphylaxis during anesthesia in France: an 8-year national survey. *J Allergy Clin Immunol* 2011; 128:366-73.
54. Dong SW, Mertes PM, Petitpain N, Hasdenteufel F, Malinovsky JM. Hypersensitivity reactions during anesthesia. Results from the ninth French survey (2005-2007). *Minerva Anesthesiol* 2012; 78:868-78.
55. Leysen J, De Witte, L., Bridts, CH., Ebo, DG. Anaphylaxis during general anaesthesia: a 10-year survey at the University Hospital Antwerp P Belg Roy Acad Med 2013; 2:88-100.
56. Antunes J, Kochuyt AM, Ceuppens JL. Perioperative allergic reactions: experience in a Flemish referral centre. *Allergol Immunopathol (Madr)* 2014; 42:348-54.
57. Low AE, McEwan JC, Karanam S, North J, Kong KL. Anaesthesia-associated hypersensitivity reactions: seven years' data from a British bi-specialty clinic. *Anaesthesia* 2016; 71:76-84.
58. Mertes PM, Malinovsky JM, Jouffroy L, Aberer W, Terreehorst I, Brockow K, et al. Reducing the risk of anaphylaxis during anesthesia: 2011 updated guidelines for clinical practice. *J Investig Allergol Clin Immunol* 2011; 21:442-53.
59. Guilloux L, Ricard-Blum S, Ville G, Motin J. A new radioimmunoassay using a commercially available solid support for the detection of IgE antibodies against muscle relaxants. *J Allergy Clin Immunol* 1992; 90:153-9.

60. Gueant JL, Mata E, Monin B, Moneret-Vautrin DA, Kamel L, Nicolas JP, et al. Evaluation of a new reactive solid phase for radioimmunoassay of serum specific IgE against muscle relaxant drugs. *Allergy* 1991; 46:452-8.
61. Harle DG, Baldo BA, Fisher MM. Immunoassays employing substituted ammonium compounds other than neuromuscular blocking drugs to increase the detection of IgE antibodies to these drugs. *Mol Immunol* 1990; 27:1039-45.
62. Baldo BA, Fisher MM. Diagnosis of IgE-dependent anaphylaxis to neuromuscular blocking drugs, thiopentone and opioids. *Ann Fr Anesth Reanim* 1993; 12:173-81.
63. Baldo BA, Fisher MM. Substituted ammonium ions as allergenic determinants in drug allergy. *Nature* 1983; 306:262-4.
64. Didier A, Cador D, Bongrand P, Furstoss R, Fourneron P, Senft M, et al. Role of the quaternary ammonium ion determinants in allergy to muscle relaxants. *J Allergy Clin Immunol* 1987; 79:578-84.
65. Laxenaire MC, Mertes PM. Anaphylaxis during anaesthesia. Results of a two-year survey in France. *Br J Anaesth* 2001; 87:549-58.
66. Mertes PM, Laxenaire MC, Alla F. Anaphylactic and anaphylactoid reactions occurring during anesthesia in France in 1999-2000. *Anesthesiology* 2003; 99:536-45.
67. Mertes PM, Laxenaire MC. Adverse reactions to neuromuscular blocking agents. *Curr Allergy Asthma Rep* 2004; 4:7-16.
68. Mertes PM, Laxenaire MC, Lienhart A, Aberer W, Ring J, Pichler WJ, et al. Reducing the risk of anaphylaxis during anaesthesia: guidelines for clinical practice. *J Investig Allergol Clin Immunol* 2005; 15:91-101.
69. Harle DG, Baldo BA, Fisher MM. Detection of IgE antibodies to suxamethonium after anaphylactoid reactions during anaesthesia. *Lancet* 1984; 1:930-2.
70. Harle DG, Baldo BA, Fisher MM. Assays for, and cross-reactivities of, IgE antibodies to the muscle relaxants gallamine, decamethonium and succinylcholine (suxamethonium). *J Immunol Methods* 1985; 78:293-305.
71. Harboe T, Guttormsen AB, Irgens A, Dybendal T, Florvaag E. Anaphylaxis during anesthesia in Norway: a 6-year single-center follow-up study. *Anesthesiology* 2005; 102:897-903.
72. Fisher MM, Baldo BA. Immunoassays in the diagnosis of anaphylaxis to neuromuscular blocking drugs: the value of morphine for the detection of IgE antibodies in allergic subjects. *Anaesth Intensive Care* 2000; 28:167-70.
73. Florvaag E, Johansson SG, Oman H, Venemalm L, Degerbeck F, Dybendal T, et al. Prevalence of IgE antibodies to morphine. Relation to the high and low incidences of NMBA anaphylaxis in Norway and Sweden, respectively. *Acta Anaesthesiol Scand* 2005; 49:437-44.
74. Ebo DG, Venemalm L, Bridts CH, Degerbeck F, Hagberg H, De Clerck LS, et al. Immunoglobulin E antibodies to rocuronium: a new diagnostic tool. *Anesthesiology* 2007; 107:253-9.
75. Uyttebroek AP, Sabato V, Bridts CH, De Clerck LS, Ebo DG. Immunoglobulin E antibodies to atracurium: a new diagnostic tool? *Clin Exp Allergy* 2015; 45:485-7.
76. Laroche D, Chollet-Martin S, Leturgie P, Malzac L, Vergnaud MC, Neukirch C, et al. Evaluation of a new routine diagnostic test for immunoglobulin E sensitization to neuromuscular blocking agents. *Anesthesiology* 2011; 114:91-7.
77. Leysen J, Bridts CH, De Clerck LS, Vercauteren M, Lambert J, Weyler JJ, et al. Allergy to rocuronium: from clinical suspicion to correct diagnosis. *Allergy* 2011; 66:1014-9.
78. Leysen J, Uyttebroek A, Sabato V, Bridts CH, De Clerck LS, Ebo DG. Predictive value of allergy tests for neuromuscular blocking agents: tackling an unmet need. *Clin Exp Allergy* 2014; 44:1069-75.
79. Florvaag E, Johansson SG, Oman H, Harboe T, Nopp A. Pholcodine stimulates a dramatic increase of IgE in IgE-sensitized individuals. A pilot study. *Allergy* 2006; 61:49-55.
80. Fisher MM, Baldo BA. Persistence of allergy to anaesthetic drugs. *Anaesth Intensive Care* 1992; 20:143-6.

81. Ebo DG, Bridts CH, Hagendorens MM, Mertens CH, De Clerck LS, Stevens WJ. Flow-assisted diagnostic management of anaphylaxis from rocuronium bromide. *Allergy* 2006; 61:935-9.
82. Uyttebroek AP, Sabato V, Leysen J, Bridts CH, De Clerck LS, Ebo DG. Flowcytometric diagnosis of atracurium-induced anaphylaxis. *Allergy* 2014; 69:1324-32.
83. Sabato V, Van Gasse A, Cop N, Claesen K, Decuyper I, Faber M, et al. The Mas-Related G Protein-Coupled Receptor MRGPRX2 Is Expressed on Human Basophils and up-Regulated upon Activation. *J All Clin Imm* 2017.
84. Baldo BA, Pham NH. Histamine-releasing and allergenic properties of opioid analgesic drugs: resolving the two. *Anaesth Intensive Care* 2012; 40:216-35.
85. Armentia A, Ruiz-Munoz P, Quesada JM, Postigo I, Herrero M, Martin-Gil FJ, et al. Clinical value of morphine, pholcodine and poppy seed IgE assays in drug-abusers and allergic people. *Allergol Immunopathol (Madr)* 2013; 41:37-44.
86. Armentia A, Pineda F, Palacios R, Martin-Gil FJ, Miguel AS, Arenal JJ, et al. Utility of opium seed extract tests in preventing hypersensitivity reactions during surgery. *Allergol Immunopathol (Madr)* 2014; 42:56-63.
87. Van Gasse AL, Hagendorens MM, Sabato V, Bridts CH, De Clerck LS, Ebo DG. IgE to Poppy Seed and Morphine Are Not Useful Tools to Diagnose Opiate Allergy. *J Allergy Clin Immunol Pract* 2015; 3:396-9.
88. Leysen J, De Witte L, Sabato V, Faber M, Hagendorens M, Bridts C, et al. IgE-mediated allergy to pholcodine and cross-reactivity to neuromuscular blocking agents: Lessons from flow cytometry. *Cytometry B Clin Cytom* 2013; 84:65-70.
89. Swerts S, Van Gasse A, Leysen J, Faber M, Sabato V, Bridts CH, et al. Allergy to illicit drugs and narcotics. *Clin Exp Allergy* 2014; 44:307-18.
90. Opstrup MS, Johansen JD, Zachariae C, Garvey LH. Contact allergy to chlorhexidine in a tertiary dermatology clinic in Denmark. *Contact Dermatitis* 2016; 74:29-36.
91. Ebo DG, Bridts CH, Stevens WJ. IgE-mediated anaphylaxis from chlorhexidine: diagnostic possibilities. *Contact Dermatitis* 2006; 55:301-2.
92. Faber M, Leysen J, Bridts C, Sabato V, De Clerck LS, Ebo DG. Allergy to chlorhexidine: beware of the central venous catheter. *Acta Anaesthesiol Belg* 2012; 63:191-4.
93. Opstrup MS, Malling HJ, Kroigaard M, Mosbech H, Skov PS, Poulsen LK, et al. Standardized testing with chlorhexidine in perioperative allergy--a large single-centre evaluation. *Allergy* 2014; 69:1390-6.
94. Anderson J, Rose M, Green S, Fernando SL. The utility of specific IgE testing to chlorhexidine in the investigation of perioperative adverse reactions. *Ann Allergy Asthma Immunol* 2015; 114:425-6.e1.
95. Ebo DG, Stevens WJ, Bridts CH, Matthieu L. Contact allergic dermatitis and life-threatening anaphylaxis to chlorhexidine. *J Allergy Clin Immunol* 1998; 101:128-9.
96. Opstrup MS, Poulsen LK, Malling HJ, Jensen BM, Garvey LH. Dynamics of plasma levels of specific IgE in chlorhexidine allergic patients with and without accidental re-exposure. *Clin Exp Allergy* 2016.
97. Ebo DG, Stevens WJ, Bridts CH, De Clerck LS. Latex-specific IgE, skin testing, and lymphocyte transformation to latex in latex allergy. *J Allergy Clin Immunol* 1997; 100:618-23.
98. Ebo DG, Lechkar B, Schuerwegh AJ, Bridts CH, De Clerck LS, Stevens WJ. Validation of a two-color flow cytometric assay detecting in vitro basophil activation for the diagnosis of IgE-mediated natural rubber latex allergy. *Allergy* 2002; 57:706-12.
99. Ebo DG, Hagendorens MM, Bridts CH, De Clerck LS, Stevens WJ. Sensitization to cross-reactive carbohydrate determinants and the ubiquitous protein profilin: mimickers of allergy. *Clin Exp Allergy* 2004; 34:137-44.
100. Quercia O, Stefanini GF, Scardovi A, Asero R. Patients monosensitized to Hev b 8 (Hevea brasiliensis latex profilin) may safely undergo major surgery in a normal (non-latex safe) environment. *Eur Ann Allergy Clin Immunol* 2009; 41:112-6.

101. Garnier L, Selman L, Rouzaire P, Bouvier M, Roberts O, Berard F, et al. Molecular allergens in the diagnosis of latex allergy. *Eur Ann Allergy Clin Immunol* 2012; 44:73-9.
102. Ebo DG, Hagendorens MM, De Knop KJ, Verweij MM, Bridts CH, De Clerck LS, et al. Component-resolved diagnosis from latex allergy by microarray. *Clin Exp Allergy* 2010; 40:348-58.
103. Schuler S, Ferrari G, Schmid-Grendelmeier P, Harr T. Microarray-based component-resolved diagnosis of latex allergy: isolated IgE-mediated sensitization to latexprofilin Hev b8 may act as confounder. *Clin Transl Allergy* 2013; 3:11.
104. Sanz ML, Gamboa PM, Garcia-Aviles C, Vila L, Dieguez I, Antepara I, et al. Flow-cytometric cellular allergen stimulation test in latex allergy. *Int Arch Allergy Immunol* 2003; 130:33-9.
105. Netti E, Colanardi MC, Dambra PP, Capuzzimati L, Loria MP, Ferrannini A, et al. Flow cytometric basophil activation test: detection of CD63 expression as a useful aid to diagnosis of latex allergy. *Ann Allergy Asthma Immunol* 2006; 97:715-6.
106. Sanz ML, Garcia-Aviles MC, Tabar AI, Anda M, Garcia BE, Barber D, et al. Basophil Activation Test and specific IgE measurements using a panel of recombinant natural rubber latex allergens to determine the latex allergen sensitization profile in children. *Pediatr Allergy Immunol* 2006; 17:148-56.
107. Faber MA, Sabato V, Bridts CH, Nayak A, Beezhold DH, Ebo DG. Clinical relevance of the Hevea brasiliensis lipid transfer protein Hev b 12. *J Allergy Clin Immunol* 2015; 135:1645-8.
108. Van Gasse AL, Mangoldt EA, Faber M, Sabato V, Bridts CH, Ebo DG. Molecular allergy diagnosis: status anno 2015. *Clin Chim Acta* 2015; 444:54-61.
109. Ratner B, Crawford LV. The allergenicity of gelatin. *Int Arch Allergy Appl Immunol* 1955; 6:370-1.
110. Mullins RJ, James H, Platts-Mills TA, Commins S. Relationship between red meat allergy and sensitization to gelatin and galactose-alpha-1,3-galactose. *J Allergy Clin Immunol* 2012; 129:1334-42.e1.
111. Ebo DG, Faber M, Sabato V, Leysen J, Gadisseur A, Bridts CH, et al. Sensitization to the mammalian oligosaccharide galactose-alpha-1,3-galactose (alpha-gal): experience in a Flemish case series. *Acta Clin Belg* 2013; 68:206-9.
112. Uyttebroek A, Sabato V, Bridts CH, De Clerck LS, Ebo DG. Anaphylaxis to succinylated gelatin in a patient with a meat allergy: galactose-alpha(1, 3)-galactose (alpha-gal) as antigenic determinant. *J Clin Anesth* 2014; 26:574-6.
113. Chung CH, Mirakhur B, Chan E, Le QT, Berlin J, Morse M, et al. Cetuximab-induced anaphylaxis and IgE specific for galactose-alpha-1,3-galactose. *N Engl J Med* 2008; 358:1109-17.
114. Commins SP, Satinover SM, Hosen J, Mozena J, Borish L, Lewis BD, et al. Delayed anaphylaxis, angioedema, or urticaria after consumption of red meat in patients with IgE antibodies specific for galactose-alpha-1,3-galactose. *J Allergy Clin Immunol* 2009; 123:426-33.
115. Sanz ML, Garcia BE, Prieto I, Tabar A, Oehling A. Specific IgE determination in the diagnosis of beta-lactam allergy. *J Investig Allergol Clin Immunol* 1996; 6:89-93.
116. Blanca M, Mayorga C, Torres MJ, Reche M, Moya MC, Rodriguez JL, et al. Clinical evaluation of Pharmacia CAP System RAST FEIA amoxicilloyl and benzylpenicilloyl in patients with penicillin allergy. *Allergy* 2001; 56:862-70.
117. Sanz ML, Gamboa PM, Antepara I, Uasuf C, Vila L, Garcia-Aviles C, et al. Flow cytometric basophil activation test by detection of CD63 expression in patients with immediate-type reactions to betalactam antibiotics. *Clin Exp Allergy* 2002; 32:277-86.
118. Gamboa PM, Garcia-Aviles MC, Urrutia I, Antepara I, Esparza R, Sanz ML. Basophil activation and sulfidoleukotriene production in patients with immediate allergy to betalactam antibiotics and negative skin tests. *J Investig Allergol Clin Immunol* 2004; 14:278-83.
119. Torres MJ, Padial A, Mayorga C, Fernandez T, Sanchez-Sabate E, Cornejo-Garcia JA, et al. The diagnostic interpretation of basophil activation test in immediate allergic reactions to betalactams. *Clin Exp Allergy* 2004; 34:1768-75.

120. Abuaf N, Rostane H, Rajoely B, Gaouar H, Autegarden JE, Leynadier F, et al. Comparison of two basophil activation markers CD63 and CD203c in the diagnosis of amoxicillin allergy. *Clin Exp Allergy* 2008; 38:921-8.
121. De Week AL, Sanz ML, Gamboa PM, Aberer W, Sturm G, Bilo MB, et al. Diagnosis of immediate-type beta-lactam allergy in vitro by flow-cytometric basophil activation test and sulfidoleukotriene production: a multicenter study. *J Investig Allergol Clin Immunol* 2009; 19:91-109.
122. Eberlein B, Leon Suarez I, Darsow U, Rueff F, Behrendt H, Ring J. A new basophil activation test using CD63 and CCR3 in allergy to antibiotics. *Clin Exp Allergy* 2010; 40:411-8.
123. Garcia-Ortega P, Marin A. Usefulness of the basophil activation test (BAT) in the diagnosis of life-threatening drug anaphylaxis. *Allergy* 2010; 65:1204.
124. Torres MJ, Ariza A, Fernandez J, Moreno E, Laguna JJ, Montanez MI, et al. Role of minor determinants of amoxicillin in the diagnosis of immediate allergic reactions to amoxicillin. *Allergy* 2010; 65:590-6.
125. Torres MJ, Romano A, Blanca-Lopez N, Dona I, Canto G, Ariza A, et al. Immunoglobulin E-mediated hypersensitivity to amoxicillin: in vivo and in vitro comparative studies between an injectable therapeutic compound and a new commercial compound. *Clin Exp Allergy* 2011; 41:1595-601.
126. Abuaf N, Rajoely B, Ghazouani E, Levy DA, Pecquet C, Chabane H, et al. Validation of a flow cytometric assay detecting in vitro basophil activation for the diagnosis of muscle relaxant allergy. *J Allergy Clin Immunol* 1999; 104:411-8.
127. Monneret G, Benoit Y, Debard AL, Gutowski MC, Topenot I, Bienvenu J. Monitoring of basophil activation using CD63 and CCR3 in allergy to muscle relaxant drugs. *Clin Immunol* 2002; 102:192-9.
128. Sudheer PS, Hall JE, Read GF, Rowbottom AW, Williams PE. Flow cytometric investigation of peri-anaesthetic anaphylaxis using CD63 and CD203c. *Anaesthesia* 2005; 60:251-6.
129. Kvedariene V, Kamey S, Ryckwaert Y, Rongier M, Bousquet J, Demoly P, et al. Diagnosis of neuromuscular blocking agent hypersensitivity reactions using cytofluorimetric analysis of basophils. *Allergy* 2006; 61:311-5.
130. Sainte-Laudy J, Orsel I. Interest of a new flow cytometric protocol applied to diagnosis and prevention of per anaesthetic accidents induced by neuromuscular blockers. *Rev. Fr. Allergol.* 2008; 48:5.
131. Hagau N, Gherman-Ionica N, Sfichi M, Petrisor C. Threshold for basophil activation test positivity in neuromuscular blocking agents hypersensitivity reactions. *Allergy Asthma Clin Immunol* 2013; 9:42.