This item is the archived peer-reviewed author-version of:

Single-layer graphene modulates neuronal communication and augments membrane ion currents

Reference:

Pampaloni Niccolò Paolo, Lottner Martin, Giugliano Michele, Matruggio Alessia, D' Amico Francesco, Prato Maurizio, Garrido Josè Antonio, Ballerini Laura, Scaini Denis.-
Single-layer graphene modulates neuronal communication and augments membrane ion currents
Nature nanotechnology - ISSN 1748-3387 - 13:8(2018), p. 755-764
Full text (Publisher's DOI): <https://doi.org/10.1038/S41565-018-0163-6>
To cite this reference: <https://hdl.handle.net/10067/1521530151162165141>

Single-layer graphene modulates neuronal communication and augments membrane ion currents

Niccolò Paolo Pampaloni^{1a}, Martin Lottner^{2a}, Michele Giugliano^{3,4,5}, Alessia Matruglio^{6b}, Francesco D'Amico⁷, Maurizio Prato^{8,9,10}, Josè Antonio Garrido^{11,12*}, Laura Ballerini^{1*}, Denis Scaini^{1,7*}

¹ International School for Advanced Studies (SISSA), Trieste, Italy

² Walter Schottky Institut and Physik-Department, Technische Universität München, Am Coulombwall, Garching, Germany

³ Molecular, Cellular, and Network Excitability, Department of Biomedical Sciences, Universiteit Antwerpen, Antwerpen, Belgium

⁴ Department of Computer Science, University of Sheffield, S1 4DP Sheffield, UK

⁵ Lab of Neural Microcircuitry, Brain Mind Institute, EPFL, CH-1015 Lausanne, Switzerland

⁶ CNR-IOM - Istituto Officina dei Materiali, Area Science Park – Basovizza, S.S. 14 km 163,5 Trieste - Italy

⁷ Elettra Sincrotrone Trieste S.C.p.A., S.S. 14 Km 163.5 in Area Science Park, I-34149 Trieste, Italy

⁸ Department of Chemical and Pharmaceutical Sciences, University of Trieste, Trieste, Italy

⁹ Nanobiotechnology Laboratory, CIC biomaGUNE, San Sebastián, Spain

¹⁰ Ikerbasque, Basque Foundation for Science, Bilbao, Spain

¹¹ Catalan Institute of Nanoscience and Nanotechnology (ICN2), CSIC and The Barcelona Institute of Science and Technology, Campus UAB, Bellaterra, 08193 Barcelona, Spain

¹² ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain

^a these authors equally contributed to the work

^b Present address: CERIC-ERIC (Central European Research Infrastructure Consortium), Area Science Park, Basovizza, S.S. 14 km 163,5 Trieste, Italy

Abstract

The use of graphene-based materials to engineer sophisticated bio-sensing interfaces adaptable to the central nervous system, requires a detailed comprehension of the behaviour of such materials in a biological context. Graphene peculiar properties may cause various cellular changes, but the underlying mechanisms remain unclear. Here, we show that single-layer graphene increases neuronal firing *via* altering membrane-associated functions in cultured cells. Graphene tunes the extracellular ions distribution at the interface with neurons, a key regulator of neuronal excitability. The resulting membrane biophysical changes include stronger potassium ion currents, with a significant shift in the fraction of neuronal firing phenotypes from *adapting* to *tonically firing*. By experimental and theoretical approaches we hypothesize that crucial to these effects are the graphene-ion interactions that are maximized when single layer graphene is deposited on electrically insulating substrates.

Graphene is a highly versatile two-dimensional nanomaterial widely adopted in many domains of science and technology, including advanced biomedical applications, due to its important physical properties [1,2,3]. Its high carrier mobility and optical transparency enable, for example, the design of novel transparent electrodes in optoelectronics [4]. Combining these features and, in particular, its remarkable electro-conductivity, makes graphene extremely appealing in neuroengineering, with reference to invasive implant technologies for brain biosensors and electrodes [5,6,7]. Despite the great interest and hopes raised by late development in graphene applications, the understanding of its functional interactions with the brain tissue is

55 still limited, particularly concerning the close proximity of a single plain layer of
56 carbon atoms and the neuronal membrane ion fluxes in a biological milieu. So far,
57 reports have shown that graphene-based materials can be safely interfaced with
58 active neuronal cells [8,9,10], however an in-depth study on the influence of single-
59 layer graphene (SLG) on the biophysics of neurons and *ex vivo* neuronal
60 microcircuits upon is missing. In numerous electroceuticals applications [11,12],
61 graphene is in contact with the extracellular environment that surrounds the excitable
62 cell membranes. We thus asked: can SLG directly or indirectly alter neuronal
63 activity? Which manipulations of graphene might be adopted to regulate *ad hoc*
64 these interactions? Answering to these questions appears pivotal for future research
65 in bio-hybrid electronic devices and, more in general, for providing insights on the
66 deep interactions of technology with nature. Here, SLG obtained by chemical vapour
67 deposition (CVD) was used to interface mammalian neurons, dissociated from the
68 rat hippocampi, as a culture substrate. We employed different architectures of a
69 single-atomic layer graphene: (i) in contact with electrically-insulating/conductive
70 substrates or (ii) suspended. By these arrangements, we studied the collective
71 electrical activity of neuronal networks coupled on graphene and demonstrated that,
72 when suspended, SLG increased neuronal excitability *via* inducing specific changes
73 in membrane biophysics. These consist in a significant shift of the fraction of
74 neuronal firing phenotypes from *adapting* to *tonically* firing. We then propose that
75 graphene selectively modifies membrane-associated neuronal functions and
76 hypothesize a specific interaction between graphene and cations, in particular
77 potassium, in the extracellular solution crucially regulating cell excitability.

78

79 **SLG potentiates cell signalling in neuronal networks**

80 Large films of SLG and multi-layer graphene (MLG) were characterized by atomic
 81 force microscopy (AFM) and compared to glass pristine (Control) and gold-metalized
 82 (Au) glass samples (Fig. 1a). The quality of CVD-grown SLG and MLG samples was
 83 assessed by Raman and X-Ray photo-electron analysis. The recorded Raman
 84 spectra (Fig. 1b) supports the high quality of the SLG and MLG, by the low I_D/I_G
 85 ratios indicating a small amount of sp^3 hybridized carbon atoms at grain boundaries
 86 or binding surface moieties [13]. X-Ray photo-electron spectroscopy (XPS) analysis
 87 reveals that both SLG and MLG samples contain a low degree of metal
 88 contamination (Fig. 1c).

89 To probe the electrical behaviour of excitable biological cells, we plated hippocampal
 90 neurons directly on graphene- and Au-coated coverslips. Recently, several reports
 91 described the successful growth of different cell types on graphene and graphene-
 92 based materials [9,14,15], but rarely investigating the *ex vivo* development and
 93 functional analysis of primary mammalian cells and neuronal microcircuits on
 94 uncoated monolayer of graphene. Neurons plated on glass coverslips were instead
 95 used as Control cultures [16,17,18]. SLG, MLG and Au substrates allow the growth
 96 of cells, with comparable mature morphology to Control (Fig. 2a). We further probed
 97 neuronal networks viability by quantifying network size and the ratio between
 98 neuronal and glial cells after 8-10 *days in vitro* (DIV), using immunofluorescence
 99 markers for neurons (class III β -tubulin) and astrocytes (GFAP). No differences were
 100 detected across all groups (Fig. 2b).

101 In culture, neurons develop functional synapses and display spontaneous collective
 102 electrical activity as a result of recurrent connections. We recorded spontaneous
 103 synaptic activity after 8÷10 DIV. This is indirectly informative of the combined effect
 104 of existence, number, transfer gain of neuronal connections, and intensity of

collective neuronal interactions. Heterogeneous post-synaptic currents (PSC, Supplementary Fig. 1b) were detected as inward currents of variable amplitudes [16] in all conditions, as shown in Figure 2c. While PSCs amplitude recorded from neurons growing on SLG, MLG and Au were similar to Control (top box plots in Fig. 2c) the PSCs frequency (bottom) was significantly higher in SLG than in other conditions. Instead, both MLG (chemically similar to SLG), and Au (chemically different but characterized by high electrical conductivity), did not affect the frequency of the synaptic events. The observed effect on PSCs is independent of the SLG film transfer-method used (Supplementary Fig. 1a). This suggests the mechanistic involvement of specific properties of the plain sheet of carbon atoms, in the modulation of PSCs frequency, but only when assembled as a monolayer. Miniature synaptic currents (mPSCs; Fig. 3a) were then recorded in a subset of Control and SLG neurons by further application of the fast-inactivating voltage-gated sodium channel blocker, tetrodotoxin (TTX, 1 μ M), which impairs the action potentials (APs) and thus blocks network activity. Studying mPSCs allows disambiguating dynamical from structural components of the emerging network activity. In particular, mPSCs reflect the stochastic release of vesicles from the presynaptic terminals at individual synapses impinging onto the recorded neuron: their frequency depends on the pre-synaptic release probability and on the number of synaptic contacts, while their amplitude depends on postsynaptic receptor sensitivity [19]. As pointed out by the box plots in Figure 3a, we found significant difference neither in the frequency nor the amplitude of mPSCs, recorded in Control or SLG conditions. This suggests that the increased PSCs activity described earlier in SLG does not involve structural changes in the number or properties of synaptic connections. This is further supported by immune-labelling experiments, where the

130 number of VGlut1-positive *puncta*, used to label and identify glutamatergic
131 presynaptic terminals [20], was not altered by the presence of SLG (Fig. 3b).
132 To rule out that SLG could interfere with the network composition or the maturation
133 of inhibitory neurons, we carefully considered two alternative hypotheses. In the first
134 one, the fraction of excitatory to inhibitory neurons is altered by SLG (e.g. in favour of
135 the former), thus biasing the spontaneous network electrical activity detected as
136 PSCs. We thus performed co-immunostaining with antibodies anti-class III β -tubulin
137 and anti-GABA (Fig. 3c) and quantified the percentage of double-positive cells,
138 indicating exclusively inhibitory neurons. In Control and SLG, we detected a
139 comparable probability of finding double-positive cells (plot in Fig. 3c), thus ruling out
140 SLG-induced alterations in the excitatory/inhibitory balance.
141 In the second hypothesis, SLG slows down the maturation of chloride ion fluxes
142 through GABA_A receptors. In neurons, the intracellular chloride concentration
143 determines the amplitude of the inhibitory currents and, across successive
144 developmental stages, shifts from higher to lower values, compared to the
145 extracellular milieu [21]. Correspondingly, the activation of GABA_A receptors results
146 in a depolarizing (hyperpolarizing) drive of immature (mature) neurons [22]. We
147 performed chloride imaging in living cells, using a quinoline-based Cl⁻ indicator dye:
148 MQAE (N-[6-methoxyquinolyl] acetoethyl ester) [23]. As reported in Supplementary
149 Fig. 2a, SLG had no impact in the GABAergic system maturation *in vitro*, a notion
150 further supported by NKCC1 quantification (Supplementary Fig. 2b).
151 Thus, the SLG-mediated increase in neuronal signalling involve neither major
152 network synaptic rearrangements, such as increased synaptogenesis, nor alterations
153 in network composition or maturation of network inhibition.

To ultimately clarify the biophysical mechanisms leading to the boost in neuronal activity exhibited only by SLG, we examined single-cell excitability by current-clamp recordings. When Control and SLG neurons were held at -60 mV in standard extracellular solution, an unbiased comparison of the basal AP frequency (Fig. 4a) could be obtained. Consistent with the PSC observations, we detected a significantly higher AP frequency in SLG neurons than in Control ones. The subsequent bath addition of antagonists, selective for excitatory and inhibitory synaptic receptors, such as Gabazine ($5\text{ }\mu\text{M}$), CNQX ($20\text{ }\mu\text{M}$) and APV ($50\text{ }\mu\text{M}$), was employed to functionally decouple the recorded neurons from the synaptic network. Under these conditions, intrinsic neuronal active membrane properties were evaluated evoking AP responses by positive current pulses, delivered from the same resting potential (-60 mV) [24,25]. When comparing the AP overshoot amplitude, half-amplitude width, threshold and maximal rising slope [25,26,27] in Control and SLG conditions no significant difference was found in the two groups (see Methods). All these considerations taken together suggest no major involvement of voltage-gated fast-inactivating Na^+ channel [27,28,29] in explaining the SLG-induced effects.

In addition, at -60 mV resting potential, under the cocktail of synaptic blockers, brief and sufficiently strong depolarizing pulses ($2\div 4$ ms; 1 nA) easily evoked single APs in SLG and Control. The voltage trajectory of evoked single APs was followed by a transient after-hyperpolarizing (AHP) in SLG neurons, while only by a small after depolarization (ADP) in Control (Fig. 4b). When quantified in terms of the area below such trajectories, referred to the resting potential as baseline (histogram in Fig. 4b), the AHP in SLG neurons was significantly different than the ADP detected in Control. Figure 4b shows the sensitivity of the AHP to various K^+ channel-blockers. All these observations demonstrate that the AHP detected in SLG

neurons was likely mediated by mixed K^+ conductances, including those activated by intracellular accumulation of free Ca^{++} [30,31]. The expression of these membrane channels is functionally related to spike-frequency adaptation, where sustained APs progressively slow-down over time. We then further examined the sustained discharge patterns of Control and SLG neurons, by injecting longer (1 s; 200 pA) depolarizing pulses (Fig. 4c and 4e). In the majority (81.8%; Fig. 4d) of Control neurons, sustained AP firing was dominated by spike frequency adaptation, which we named *adapting* discharge phenotype. This often resulted in an early burst of closely spaced APs, followed by a progressive decay of AP amplitudes, leading to adaptation. On the contrary, SLG neurons (83.3%; Fig. 4d) showed no APs adaptation, which we named *tonic* discharge phenotype, where cells fired continuously without apparent accommodation [32,33]. Taken together, these data hint at a complex homeostasis in the K^+ currents expressed by neurons when coupled to SLG substrates. This hypothesis was reinforced by results obtained under voltage-clamp in control and SLG neurons, where depolarizing voltage pulses starting from a -60 mV holding potential baseline evoked an outward current (Fig. 4f), presumably due to the activation of a mixed population of K^+ channels. When examined under these conditions, SLG neurons were characterized by a significantly larger outward current at positive potentials, shown in the steady-state current/voltage (I/V) plot of Figure 4f, likely consequence of an up-regulation of mixed K^+ currents. This evidence taken together strongly indicates that SLG substrates induce active changes in the electrical properties of growing neurons, presumably related to altered homeostasis of K^+ membrane currents and leading to a modulation of the single-cell firing phenotypes and ultimately to an increased network activity. Importantly, SLG neurons generated more APs when compared to

Controls, even in response to milder and shorter stimuli (Supplementary Fig. 3a and b).

The observed correlations between single-cell properties, resulting phenotype, and network effects were further investigated by mathematical modelling. We addressed the causality between neuronal firing patterns and network activity, by examining a spike-rate model of the electrical activity emerging in populations of cultured neurons with recurrent synaptic connections (Fig. 5a).

We simulated a network of 1600 neurons with a fraction of excitatory to inhibitory neurons equal to 80/20 [34]. Since inhibitory cells usually display a tonic electrical phenotype only, we hypothesized that the change in the ratio between adapting and tonically firing neurons observed in our experiments, occurred in excitatory cells only. We therefore modelled two subpopulations of excitatory neurons: one with *adapting* and the other with *tonic* phenotypes. We found that the higher the relative fraction of *tonic* firing neurons, the higher the rate of occurrence of synchronized bursts (Fig. 5b and 5c). This supports the conclusion that the observed increase in the frequency of spontaneous (PSCs/APs) activity (Fig. 2c, SLG) is caused by the different ratio of cells with *adapting/tonic* phenotypes.

The potassium ions hypothesis

By a biophysical model, we tested *in silico* two mutually non-exclusive hypotheses for the observed changes in single-cell firing phenotype: (i) an increase in the total outward ionic conductance or, alternatively, (ii) a modest depletion in the extracellular concentration of K^+ ions. The rationale behind these single-cell simulations is that either SLG induced a chronic increase in ion currents involved in firing regulation, or SLG acutely altered cell firing by changing ion mobility.

In our model, by stripping down cell excitability to its bare essential, we explored whether stronger outward potassium currents may favour excitability. We found that the progressive sodium current inactivation (Fig. 5d, lower left, green traces) – occurring for simplicity in the model only over *fast* time scales – could be counterbalanced by strong K^+ currents (Fig. 5d, lower right, red traces) resulting in a sustained, tonic, response to an external current stimulus (i.e. compare Fig. 5d to 4c), instead of a progressive firing inactivation. While this effect is reversed by simulating an overexpression of Na^+ channels (e.g. at the axon initial segment), it serves us here as a proof of concept of a counter-intuitive phenomenon: increasing outward currents increases cellular excitability, by removing sodium current inactivation. Of course, an *ad hoc* increase in the maximal conductance of Na^+ channels also increases excitability, although – in the model – with distinct features in the type of transition associated to the *limit cycle* to sustained AP firing (Supplementary Fig. 4). Therefore, in the model the more K^+ channels the higher the excitability, in those regimes where progressive sodium inactivation affects firing (Fig. 5e).

Outward K^+ currents also depend on the ionic driving force beyond on the maximal conductance (i.e. $I_K \sim G_K (E_K - V)$ - see Methods), so that a change in the local ionic composition might reverse sodium inactivation too. In fact, the Nernst equilibrium potential E_K depends on the K^+ concentrations outside and inside the membranes [35]. Should SLG interfere extracellularly with K^+ bulk diffusion in its proximity (see below) then a depletion of K^+ (e.g. 10-20%, as $[K^+]_{EX} \rightarrow \delta \cdot [K^+]_{EX}$, $\delta = 0.8 - 0.9$), would lead to an increase in the ionic driving force, as E_K would decrease accordingly (i.e. $\sim 2-5$ mV). By simulations (Fig. 5f), we found that a modest decrease in E_K (e.g. from -75 to -77.6 mV) could indeed counterbalance sodium inactivation,

at least for an intermediate external stimulus intensity and without altering significantly the resting membrane potential.

We thus speculate that changes in excitability of cells coupled to SLG might be caused by an extracellular reduced mobility of K^+ at the interface between the nanomaterial and the solution, leading to a K^+ depletion at the neuronal membranes.

Localized potassium ions depletion in cell-substrate cleft

We hypothesize that at the core of SLG ability to alter neuronal excitability is the ion adsorption on graphene surfaces. This may result in a modification of ion mobility, in particular K^+ , at the neuronal/graphene interface.

Within this proposed mechanism, it is still unclear how SLG might modify K^+ ion mobility while MLG (or Au) do not. It is well known that carbon-based π electron-rich surfaces show a significant surface enrichment of cations in ionic solutions, due to specific cation- π interactions [36,37,38,39]. Because of its size, in solution, K^+ ions are weaker solvated by water, when compared to other species (e.g. Li^+ or Na^+ ions) but are still good π binders. This feature makes K^+ the best alkali metal binder to carbon-based surfaces in aqueous solutions [38,39]. This implies that, in nanoscale-confined systems, cation trapping occurring at the carbon surface level may lead to a significant local depletion of ions, in particular potassium, at the neuronal membrane/surface level.

Raman spectroscopy (sketched in Supplementary Fig. 5a) was performed on supported SLG and MLG samples in liquid condition without and in the presence of KCl and NaCl at physiological concentrations (4 mM and 150 mM, respectively) in the solution. SLG Raman G-peak exhibited a change in shape and position in the presence of NaCl and KCl D_2O solutions (Fig. 6a, left, inset) while in MLG it did not

change (Fig. 6a, right, inset). The mechanisms responsible for the observed shift in the G Raman band of graphene with, more importantly, the narrowing of the G band (FWHM) detected in SLG when samples are immersed in a KCl solution could result from charge doping [40] or internal strain [41]. More specifically, the shifts in G-band position exhibited by SLG in the presence of salt solutions could be indicative of a specific cation interaction, not measurable in MLG. Notably, the larger G-band Raman shift in KCl-treated samples ($3 \pm 0.5 \text{ cm}^{-1}$) than in the presence of NaCl ($1 \pm 0.5 \text{ cm}^{-1}$) correlates well with a larger SLG affinity for hydrated K^+ when compared to Na^+ .

Importantly, in our experimental settings, the specific cation- π interaction at surface level could result in partial K^+ depletion from the extracellular solution facing the cell membrane (Fig. 6b) due to the tiny thickness of the cleft (see Supplementary Information and Fig. S6) between cells and substrates.

Substrate modulation of graphene cation- π interaction

Our experiments have shown that SLG behaves differently from MLG in respect to the K^+ homeostasis of neurons and subsequent improved excitability. The two culturing platforms, SLG and MLG, differ only in the conductive properties of the supporting structure immediately below the first mono-atomic carbon layer exposed to the biological milieu (i.e., on one hand glass and on the other multiple layers of graphene/graphite). From the point of view of a neuron growing on its top, MLG appears as a SLG film layered on the underlying, electrically conductive, MLG. In our hypothesis, graphene efficiency in trapping K^+ ions is tuned or influenced by the electronic properties of the supporting structure [42,43].

We compared PSCs in neurons directly grown on glass (Control), on free-standing SLG (suspended SLG, see Methods) and on SLG transferred on an insulating substrate (SLG on glass) and on a conductive substrate of Indium Tin-Oxide (SLG on ITO)[18]. The results of such experiments (Figure 6c) are in full agreement with our hypothesis: the PSCs frequency was boosted by SLG on glass, and even more by suspended SLG, with no detectable effects when SLG was layered on ITO. Notably, in suspended SLG, PSCs amplitude is also significantly higher than Control.

It is not trivial to understand the underlying exact mechanism, in the absence of any theoretical model describing the dependency of graphene π -cation interaction on supporting surface properties, we speculate that surface conductivity is playing a key role. In particular, in suspended SLG, environmental disturbances are minimized allowing access to the intrinsic properties of graphene close to the unperturbed Dirac point. Superficial charge inhomogeneity is reduced in this case compared to supported samples, giving rise to a “close-to-theory” system [1,44] (Fig. 6d, left) that will fit better to cation- π simulations’ results [33,45,46]. SLG laying on metal surfaces usually undergo electron-doping resulting in a down-shift of graphene Dirac point [47] (Fig. 6d, middle). This will induce a homogeneous charge-distribution [48] that could result in a reduction of graphene cation- π interaction force. On the other hand, in SLG transferred on insulating substrates there are significant local fluctuations in surface potential [49,50], thus inducing an inhomogeneous charge distribution on graphene surface where neutral areas, where SLG band structure is basically unperturbed as in the case of suspended graphene, and p-doped areas, coexist [51,52]. In the latter case, graphene- K^+ interactions will still be present even if with less pronounced effect than on suspended SLG. Our preliminary Raman results can

explain the different behaviour of SLG towards K^+ or Na^+ and the difference between SLG and MLG behaviour in ion solution. However, they do not directly demonstrate differences between SLG and MLG in ion absorbance.

Conclusions

SLG modifies neuronal excitability and we propose that this effect is mediated by graphene ability to restrict K^+ ions mobility in close proximity to the material surface, but only when SLG is deposited on electrically insulating substrates. We cannot exclude additional mechanisms related to non-uniform charge carrier densities, affecting surface concentrations of ions [53]. Alternatively, restricted ion mobility might affect the way astrocytes regulate the extracellular *milieu* between graphene and neurons.

Ultimately, we provided multiple lines of evidence to demonstrate that SLG, when engineered on an insulating glass-substrate, is able to tune neuronal excitability. Our physiological experiments demonstrate that the detected increase in neuronal synaptic activity is caused by increased cell firing, rather than by changes in network size, synaptic density [54,55,56], inhibition/excitation ratio or inhibition maturation. We also demonstrated that neurons, exposed to SLG, up-regulate K^+ currents and switch to functionally-*tonic* firing phenotypes. Our simulations support the notion that changes in the ratio of *adapting/tonic* firing neurons will impact the global network activity [57,58] and suggest the key contribution of up-regulated potassium currents in driving this change. All these effects are not mimicked by MLG or other conductive substrates, such as Au. We propose that, due to the cation- π interactions of graphene, cations, and K^+ in particular [37,38,39], will be trapped at the graphene surface, resulting in a graded ionic depletion at distances from the material

compatible with the nanometre-scale characterizing cell-adhesion mechanisms [59,60]. This hypothesis is grounded in earlier molecular dynamics simulations at the equilibrium [37], where ionic enrichment occurs at the interface.

Graphene properties might thus affect neuronal information processing *in virtue* of the physical interactions of such a nanomaterial with the biological environment.

Novel materials might then represent, in general, unconventional tools to gain insights into genuine biological processes.

REFERENCES

- [1] Novoselov, K. S. et al. Electric Field Effect in Atomically Thin Carbon Films. *Science* 306, 666-669 (2004).
- [2] Geim, A. K. & Novoselov, K. S. The rise of graphene. *Nat Mater.* 6, 183-191 (2007).
- [3] Yang, Y. et al. Graphene based materials for biomedical applications. *Mat Today* 16, 365-373 (2013).
- [4] Li, X., et al. Transfer of large-area graphene films for high-performance transparent conductive electrodes. *Nano Lett.* 9, 4359-4363 (2009)
- [5] Shin, S. R. et al. Graphene-based materials for tissue engineering. *Adv Drug Deliv Rev.* 105, 255-274 (2016).
- [6] Lu, Y. et al. Flexible Neural Electrode Array Based-on Porous Graphene for Cortical Microstimulation and Sensing. *Sci Rep.* 6, 33526 (2016).
- [7] Herbert, C. et al. Flexible graphene solution-gated field effect transistors, efficient transducers for micro-electrocorticography *Adv. Funct. Mater.* 28, 1703976 (2017).

- 377 [8] Kim, J. et al. Monolayer Graphene-Directed Growth and Neuronal
 378 Differentiation of Mesenchymal Stem Cells. J Biomed Nanotechnol. 11, 2024-
 379 2033 (2015).
- 380 [9] Fabbro, A. et al. Graphene-Based Interfaces Do Not Alter Target Nerve Cells.
 381 ACS Nano 10, 615-623 (2016).
- 382 [10] Rauti, R. et al. Graphene Oxide Nanosheets Reshape Synaptic Function in
 383 Cultured Brain Networks. ACS Nano 10, 4459-4471 (2016).
- 384 [11] Famm, K. et al. Drug discovery: a jump-start for electroceuticals. Nature
 385 496, 159-161 (2013).
- 386 [12] Rivnay, J. et al. Next-generation probes, particles, and proteins for neural
 387 interfacing. Sci Adv. 3, e1601649 (2017).
- 388 [13] Cançado, L.G. et al. Quantifying defects in graphene via Raman
 389 spectroscopy at different excitation energies. Nano Lett. 11, 3190-3196 (2011).
- 390 [14] Kim, J. et al, Monolayer Graphene-Directed Growth and Neuronal
 391 Differentiation of Mesenchymal Stem Cells. J Biomed. Nanotechnol. 11, 2024-
 392 2033 (2015).
- 393 [15] Baldrighi, M. et al, Carbon Nanomaterials Interfacing with Neurons: An In
 394 vivo Perspective. Front. Neurosci. 10, 250 (2016).
- 395 [16] Lovat, V. et al. Carbon nanotube substrates boost neuronal electrical
 396 signaling. Nano Lett. 5, 1107-1110 (2005).
- 397 [17] Cellot, G. et al. Carbon nanotubes might improve neuronal performance by
 398 favouring electrical shortcuts. Nat. Nanotechnol. 4, 126-133 (2009).
- 399 [18] Cellot, G. et al, Carbon nanotube scaffolds tune synaptic strength in cultured
 400 neural circuits: novel frontiers in nanomaterial-tissue interactions. J. Neurosci.
 401 31, 12945-12953 (2011).

- 402 [19] Raastad, M. et al. Putative Single Quantum and Single Fibre Excitatory
 403 Postsynaptic Currents Show Similar Amplitude Range and Variability in Rat
 404 Hippocampal Slices. *Eur J Neurosci.* 4, 113-117 (1992).
- 405 [20] Pampaloni, NP. et al. Sculpting neurotransmission during synaptic
 406 development by 2D nanostructured interfaces. *Nanomedicine.* S1549-9634,
 407 30082-5 (2017).
- 408 [21] Arosio, D. & Ratto, GM. Twenty years of fluorescence imaging of
 409 intracellular chloride. *Front Cell Neurosci.* 8, 258 (2014).
- 410 [22] Cherubini, E. GABA mediated excitation in immature rat CA3 hippocampal
 411 neurons. *Int J Dev Neurosci.* 8, 481-90 (1990).
- 412 [23] Marandi, N., Konnerth, A. & Garaschuk O. Two-photon chloride imaging in
 413 neurons of brain slices. *Pflugers Arch.* 445, 357-65 (2002).
- 414 [24] Ruscheweyh, R. & Sandkuhler, J. Lamina-specific membrane and discharge
 415 properties of rat spinal dorsal horn neurones in vitro. *J Physiol.* 541, 231-244
 416 (2002).
- 417 [25] Chang, YM. & Luebke, JI. Electrophysiological diversity of layer 5 pyramidal
 418 cells in the prefrontal cortex of the rhesus monkey: in vitro slice studies. *J*
 419 *Neurophysiol.* 98, 2622-32. (2007)
- 420 [26] Routh, B. N. et al. Anatomical and Electrophysiological Comparison of CA1
 421 Pyramidal Neurons of the Rat and Mouse. *J Neurophysiol.* 102, 2288–2302
 422 (2009).
- 423 [27] Renganathan, M., Cummins, TR. & Waxman, SG. Contribution of Na(v)1.8
 424 sodium channels to action potential electrogenesis in DRG neurons. *J*
 425 *Neurophysiol.* 86, 629-40 (2001).

- 426 [28] Platkiewicz, J. & Brette, R. Impact of Fast Sodium Channel Inactivation on
 427 Spike Threshold Dynamics and Synaptic Integration. PLoS Comput Biol. 7,
 428 e1001129 (2011).
- 429 [29] Kress, G. J. et al. Axonal sodium channel distribution shapes the
 430 depolarized action potential threshold of dentate granule neurons. Hippocampus
 431 20, 558-571 (2016).
- 432 [30] Storm, J. F. Action potential repolarization and a fast after-hyperpolarization
 433 in rat hippocampal pyramidal cells. J Physiol. 385, 733-759 (1987).
- 434 [31] Sah, P. & Faber, E. S. Channels underlying neuronal calcium-activated
 435 potassium currents. Prog Neurobiol. 66, 345-353 (2002).
- 436 [32] Dégenétais, E. et al. Electrophysiological properties of pyramidal neurons in
 437 the rat prefrontal cortex: an in vivo intracellular recording study. Cereb Cortex 12,
 438 1-16 (2002).
- 439 [33] Furlan, F. et al. ERG conductance expression modulates the excitability of
 440 ventral horn GABAergic interneurons that control rhythmic oscillations in the
 441 developing mouse spinal cord. J. Neurosci. 27, 919-928 (2007).
- 442 [34] Marom, S. & Shahaf, G. Development, learning and memory in large random
 443 networks of cortical neurons: lessons beyond anatomy. Q Rev Biophys. 35, 63-
 444 87 (2002).
- 445 [35] Sterratt, D. Principles of computational modelling in neuroscience.
 446 Cambridge University Press(2011).
- 447 [36] Kumpf R A, & Dougherty D A. A mechanism for ion selectivity in potassium
 448 channels: computational studies of cation- π interactions. Science. 261, 1708-10
 449 (1993).

- 450 [37] Shi, G, et al. Ion enrichment on the hydrophobic carbon-based surface in
451 aqueous salt solutions due to cation- π interactions. *Sci Rep.*, 3, 3436 (2013).
- 452 [38] Pham T A, et al. Salt Solutions in Carbon Nanotubes: The Role of Cation- π
453 Interactions *J. Phys. Chem. C*, 120, 7332-7338 (2016).
- 454 [39] Williams, C D, et al. Effective Polarization in Pairwise Potentials at the
455 Graphene-Electrolyte Interface *J. Phys. Chem. Lett.*, 8, 703-708 (2017)
- 456 [40] Dong, X et al. Doping single-layer graphene with aromatic molecules. *Small*.
457 5, 1422-6 (2009)
- 458 [41] Chacón-Torres, JC, Wirtz, L, Pichler, T. Manifestation of charged and
459 strained graphene layers in the Raman response of graphite intercalation
460 compounds. *ACS Nano*. 7, 9249-59 (2013).
- 461 [42] Novák M, et al. Solvent effects on ion-receptor interactions in the presence
462 of an external electric field. *Phys Chem Chem Phys*. 18, 30754-30760 (2016).
- 463 [43] Chen, K. et al, Electronic Properties of Graphene Altered by Substrate
464 Surface Chemistry and Externally Applied Electric Field. *J. Phys. Chem. C*, 116,
465 6259-6267 (2012)
- 466 [44] Novoselov, K. et al., Two-dimensional gas of massless Dirac fermions in
467 graphene. *Nature*. 438, 197-200 (2005).
- 468 [45] Gigante, G. et al. Network Events on Multiple Space and Time Scales in
469 Cultured Neural Networks and in a Stochastic Rate Model. *PLoS Comput Biol*.
470 11, e1004547 (2015).
- 471 [46] Gambazzi, L. et al. Diminished activity-dependent brain-derived neurotrophic
472 factor expression underlies cortical neuron microcircuit hypoconnectivity resulting
473 from exposure to mutant huntingtin fragments. *J. Pharmacol Exp Ther*. 335, 13-
474 22 (2010).

- 475 [47] González-Herrero, H et al. Graphene Tunable Transparency to Tunneling
476 Electrons: A Direct Tool to Measure the Local Coupling. ACS Nano. 10, 5131-44
477 (2016).
- 478 [48] Praveen, C. S. et al., Adsorption of alkali adatoms on graphene supported
479 by the Au/Ni(111) surface, Phys Rev. B 92, 075403 (2015).
- 480 [49] Kang, Y-J. et al. Electronic structure of graphene and doping effect on SiO₂
481 Phys Rev. B 78, 115404 (2008).
- 482 [50] Miwa, R. H. et al. Doping of graphene adsorbed on the a-SiO₂ surface. Appl
483 Phys Lett 99, 163108 (2011).
- 484 [51] Ao, Z. et al. Density functional theory calculations on graphene/ α -
485 SiO₂(0001) interface. Nanoscale Res Lett. 7, 158 (2012).
- 486 [52] Fan, X. F. et al. Interaction between graphene and the surface of SiO₂. J
487 Phys Condens Matter. 24, 305004 (2012).
- 488 [53] Hille, B. Ion channels of excitable membranes. Sunderland, Mass: Sinauer,
489 (2001).
- 490 [54] Slomowitz, E. et al. Interplay between population firing stability and single
491 neuron dynamics in hippocampal networks. eLife 4, e04378 (2015).
- 492 [55] Bogaard, A. et al. Interaction of cellular and network mechanisms in
493 spatiotemporal pattern formation in neuronal networks. J Neurosci. 29, 1677-
494 1687 (2009).
- 495 [56] Sahasranamam, A. et al. Dynamical state of the network determines the
496 efficacy of single neuron properties in shaping the network activity. Sci Rep. 6,
497 26029 (2016).

- 498 [57] Radulescu, R.A. Mechanisms Explaining Transitions between Tonic and
 499 Phasic Firing in Neuronal Populations as Predicted by a Low Dimensional Firing
 500 Rate Model. PLoS One 5, e12695 (2010).
- 501 [58] Naud, R. et al. Firing patterns in the adaptive exponential integrate-and-fire
 502 model. Biol Cybern. 99, 335-347 (2008).
- 503 [59] Wrobel G, Höller M, Ingebrandt S, Dieluweit S, Sommerhage F, Bochem HP,
 504 Offenhäusser A. Transmission electron microscopy study of the cell-sensor
 505 interface. J R Soc Interface. 5, 213-222. (2008).
- 506 [60] Braun, D. & Fromherz, P. Fluorescence interference-contrast microscopy of
 507 cell adhesion on oxidized silicon. Appl Phys A 65, 341-348 (1997).
- 508 [61] Ferrari A. C. et al. Raman spectrum of graphene and graphene layers. Phys.
 509 Rev. Lett. 97, 187401 (2006).
- 510 [62] Cançado, L. G. et al. Measuring the degree of stacking order in graphite by
 511 Raman spectroscopy. Carbon 46, 272-275 (2008).
- 512 [63] Alagem, N. et al, Mechanism of Ba(2+) block of a mouse inwardly rectifying
 513 K⁺ channel: differential contribution by two discrete residues. J Physiol. 534,
 514 381-393 (2001).
- 515 [64] Alger, B. E. & Nicoll, R. A. Epileptiform burst afterhyperpolarization: calcium-
 516 dependent potassium potential in hippocampal CA1 pyramidal cells. Science
 517 210, 1122-1124 (1980).
- 518 [65] Jiang, Y. & MacKinnon, R. The barium site in a potassium channel by x-ray
 519 crystallography. J Gen Physiol. 115, 269-272 (2000).
- 520 [66] Nisenbaum, E. S. & Wilson, C. J. Potassium currents responsible for
 521 inward and outward rectification in rat neostriatal spiny projection neurons. J
 522 Neurosci. 15, 4449-4463 (1995).

- [67] Pisirowski, R. & Aldrich, R. W. Calcium activation of BK(Ca) potassium channels lacking the calcium bowl and RCK domains. *Nature*. 420, 499-502 (2002).
- [68] Stocker, M. et al, An apamin-sensitive Ca^{2+} -activated K^+ current in hippocampal pyramidal neurons. *Prot Nat Acad Sci*. 96, 4662-4667(1999).
- [69] Hodgkin, A. L. & Huxley, A. F. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol*. 117, 500-544 (1952).
- [70] French, C. R. et al. Properties of an intermediate-duration inactivation process of the voltage-gated sodium conductance in rat hippocampal CA1 neurons. *J Neurophysiol*. 115, 790-802 (2016).

Acknowledgement

We are grateful to M. Lazzarino, S. Dal Zilio and the Facility of NanoFabrication of IOM of Trieste for experimental assistance in the fabrication of suspended SLG and FIB analysis; N. Secomandi and R. Rauti for assistance in imaging. We thank A. Laio, G. Scoles, A. Nistri and B. Cortés-Llanos for helpful discussion.

This paper is based on work supported by the European Union Seventh Framework Program under grant agreement no. 696656 Graphene Flagship and no. 720270 Human Brain Project Flagship, and by the Flanders Research Foundation (grant no. G0F1517N). MP, as the recipient of the AXA Chair, is grateful to the AXA Research Fund for financial support. MP was also supported by the Spanish Ministry of Economy and Competitiveness MINECO (project CTQ2016-76721-R), by the University of Trieste and by Diputación Foral de Gipuzkoa program Red (101).

548

549 **Author Contributions**

550 N.P.P. performed electrophysiological experiments, imaging, immunochemistry,
 551 confocal microscopy and all the related analysis; M.L. fabricated supported SLG
 552 and MLG and performed all material characterization; M.G. performed
 553 mathematical simulations and analysis and contributed to the writing of the
 554 manuscript; A.M. fabricate suspended SLG and gold plated samples; F.D.A. and
 555 A.M. performed Raman experiments and data analysis on SLG and MLG in wet
 556 and dried conditions; M. P., D.S. and L.B. conceived the study; D.S., L.B., and
 557 J.A.G. designed the experimental strategy, interpreted the results and wrote the
 558 manuscript.

559 All authors discussed the results and commented on the manuscript.

560

561 **Competing financial interests**

562 The authors declare no competing financial interests.

563

564 **FIGURE LEGENDS**

565 **Figure 1 | Characterization of the substrates.** **a**, AFM topography reconstructions
 566 of glass control, SLG, MLG and gold plated glass surfaces. AFM documented a
 567 surface roughness of the materials that varied from 0.23 ± 0.02 nm in Control ($n = 3$
 568 samples), 1.5 ± 0.5 nm in SLG ($n = 3$ samples), 20 ± 10 nm for MLG ($n = 3$ samples)
 569 and 0.47 ± 0.1 nm for Au ($n = 3$ samples). Scale bar: 5 μ m. **b**, Spatial maps of the
 570 2D/G and D/G peak amplitude ratio maps. Scale bar: 10 μ m. The single punctual
 571 Raman spectra of SLG (in red) and MLG (in blue) represent mapping data points
 572 with the corresponding average peak amplitude ratios. On the right, relative spatial

2D/G and D/G ratio maps. The FWHM of the 2D peak, as well as the I_{2D}/I_G ratio are indicative of a low bilayer content in the case of SLG [61] and turbostratic graphite in the case of MLG [62]. A graphitic Raman signature is to be expected as the MLG samples have a typical thickness of a few hundreds of layers. **c**, XPS spectrum (X-Ray source: Mg K α) of SLG (in red) and MLG (in blue) on Si₃N₄. Dotted lines highlight the relevant elements, while the unlabelled features around 750 eV and 980 eV correspond to the oxygen KL₁L₁ and the carbon KVV Auger lines respectively.

Figure 2 | Single-Layer graphene increase neuronal network activity. a,

Representative SEM micrographs depicting hippocampal neurons morphology after 10 DIV, supported by the different substrates (n = 5 fields each). Culture substrates were not pre-treated with any additional adhesion molecules, which might mask the effects of graphene. Scale bar: 10 μ m. **b**, Representative fluorescent microscopy images showing dissociated hippocampal networks labelled with class III β -tubulin (for neurons) in red and GFAP (for astrocytes) in green. Scale bar: 100 μ m.

The histograms show the density of cells (top) and the neuron/glia ratio (bottom) across all four conditions, which did not significantly differ (n = 30 fields, 3 culture series, each). In addition, SLG or MLG topography did not influence neuronal fibres' orientation in respect to Control or Au (see Methods). These observations, combined to the similarity of membrane passive electrical properties (see Methods), indicated the homogeneous growth of healthy neurons [17] across substrates, with comparable levels of cellular composition and organization. **c**, Representative traces of the spontaneous network activity of neurons grown on the different substrates are shown (left), the corresponding isolated PSCs are shown superimposed (middle; in black the average values). Box plots summarize the PSC amplitude values (right,

top) and the PSC frequency ones (right, bottom) in all experimental conditions (SLG, n = 45; MLG, n = 20; Au, n = 20; Control, n = 40). Note the significantly higher PSC frequency in SLG than all other substrates (i.e., Control 1.53 ± 0.22 Hz; SLG 3.21 ± 0.41 Hz; $P=0.0010$). Differences between the variables were assessed using one-way ANOVA and multiple comparisons were adjusted by Bonferroni correction.

Figure 3 | Single-layer graphene does not increase the number of synapses

and the network composition. a, Exemplificative traces of spontaneous synaptic activity, recorded in the presence of TTX, are shown together with their superimposed mPSCs (right, in black the average values). Control (n = 11) and SLG (n = 11) mPSC frequency and amplitude are summarized in the box plots, note that no differences were detected in these parameters. **b**, Confocal images of neuronal cultures (10 DIV) in Control and SLG identifying the presynaptic VGlut1 (in green) in III β -tubulin positive cells (orange). Scale bar: 20 μ m. Higher magnifications of the region highlighted by white boxes are displayed for clarity. Scale bar: 5 μ m. The histograms on the right summarize VGlut1 puncta densities in the two conditions (5.2 ± 1.14 a.u. in Control and 4.5 ± 0.64 a.u. in SLG; $P = 0.207$, n = 30 fields, 3 cultures each). **c**, Confocal images of neuronal cultures (10 DIV) in Control and SLG identifying positive cells for class III β -tubulin and GABA. Scale bar: 10 μ m. The histograms on the right summarize the percentage of double-positive cells in the two conditions (33 ± 2.7 % in Control and 30 ± 2.5 % in SLG; $P = 0.21$, n= 20 fields each). Statistically significant difference between two data sets was assessed by Student's t test for parametric data and by Mann-Whitney for non-parametric ones.

Figure 4 | SLG triggers changes in single cell intrinsic excitability. a, Representative current-clamp recordings of hippocampal neurons in culture (10 DIV) in Control and SLG. Control and SLG neurons displayed similar resting membrane potentials (-52 ± 10 mV in SLG; -50 ± 7 mV in Control). When hold at -60 mV, the cell's spontaneous AP firing was measured as summarized in the histograms (right). Note the significantly higher AP frequency in SLG (2.60 ± 0.36 Hz in SLG, $n = 21$; 1.37 ± 0.26 Hz in Control, $n = 19$; $P = 0.0054$). **b,** Evoked single AP in Control (top) and SLG (bottom). Note the pronounced AHP in SLG neurons, that was partially abolished by each of the treatments shown: BaCl, TEA or Apamin (right, superimposed tracings). The histogram quantifies the area below the Control and SLG post-AP voltage trajectories with respect to the resting membrane potential. The AHP in SLG neurons was significantly different than the ADP detected in control neurons (-86.96 ± 23.60 mV·ms in SLG, $n = 25$; $+107.12 \pm 21.85$ mV·ms in Control, $n = 20$; $P = 0.0010$). Interestingly, the AHP was reduced (by 88%) by bath applying Ba^{++} ($BaCl_2$, 2 mM; $n = 3$), which is known to block K_{ir} and K_{Ca} membrane channels [63,64,65,66]. The AHP was also reduced (by 58%) by bath applying tetraethylammonium (TEA, 1 mM; $n = 9$) a non-selective blocker of the large majority of voltage gated K^+ membrane channels (K_v) [30], including BK_{Ca} channels [67]. Finally, Apamin (200 μ M; $n = 5$), a specific inhibitor of SK_{Ca} membrane channels [68], also strongly affected (47% reduction) the AHP. **c,** Current-clamp recordings from neurons in control and SLG revealed different cell discharge patterns, classified as *adapting* or *tonic*. **d,** Bar charts illustrate probability distributions (expressed as percentage of sampled population) of each cell type in Control ($n = 13$) and SLG ($n = 15$) cultures. Note that in Control 81.8% are *adapting* and in SLG 83.3% are *tonic*. **e,** Scatter plot of after-potential area vs. the number of action potentials (APs) in SLG

neurons when a single AP is elicited (as in b; open circle) or when multiple APs are evoked (as in c, filled circle) by 1 s long depolarizing step (-352 ± 70 mV·ms upon long depolarizing steps). **f**, Representative records of voltage-activated outward currents evoked by depolarizing current steps in Control and SLG (capacitive transients were not removed). Plot summarizes the I/V relation in Control ($n = 13$) and SLG ($n = 15$) neurons obtained upon subtraction of leak currents. Note that SLG outward currents were significantly larger than control ones (10 mV step, $P=0.032$; 20 mV step, $P=0.031$; 30 mV step, $P=0.019$). Statistically significant difference between two parametric data sets was assessed by Student's *t* test.

Figure 5 | Spike-rate network model. The electrical activity of excitatory and inhibitory recurrently interacting neurons was described by a mathematical model (see Supplemental Methods). **a**, The increase in the fraction of non-adapting neurons, observed *in vitro* on graphene substrates, predicts a higher rate of occurrence *in silico* for spontaneous “bursts” of spikes, synchronized across the network. These bursts are presynaptic correlates of the synaptic potentials, observed experimentally by voltage-clamp. **b** and **c**, Samples of the simulated network firing rates, analysed in **a**, are shown for two fractions of non-adapting neurons, *i.e.* 20% and 80%, out of the total of excitatory neurons. **d**, Counter-intuitive effects of outward potassium currents on cell excitability are explored in a single-cell biophysical model (parameters as in Table 2). As a proof of concept, we considered the simplest possible model of AP generation, as proposed by Hodgkin and Huxley (HH) [69]. This model describes the generation of a (train of) AP(s) in terms of the known interplay between fast-inactivating (~ 1 ms) inward Na^+ currents and delayed rectifier outward K^+ potassium currents –of course, by no means these are the only

membrane currents underlying the electrophysiological behaviour of rat hippocampal neurons [70]. The membrane potential responses (black traces) to an external step current was simulated, as in the experiments (see Methods). **d** and **e**, Plots within each panel exemplify how an increase (from left to right subpanels) of the maximal K^+ conductance or, in **f**, its driving force, through a depletion of extracellular K^+ ions, may to some extent reverse the inactivation of inward Na^+ currents (green traces in **d**). Then, inactivating neuronal responses may turn into sustained firing thereby increasing cell excitability (as in **c**). All in all, this suggests a specific involvement of the extracellular concentration of K^+ in neuronal excitability: the less extracellular K^+ the higher the excitability, at least in those regimes where progressive sodium inactivation affects neuronal firing disfavoring sustained *tonic* response. Parameters: in **d**, G_K in $\{0.012; 0.0216\}$ mS/mm²; in **e**, G_K in $\{0.012, 0.04, 0.06\}$ mS/mm² from left to right, I_{stim} 5 nA/mm²; in **f**, I_{stim} as in **e**, while E_K in $\{-75, -77.6, -80.5\}$ mV from left to right, corresponding to a $\{0\%, 10\%, 20\%\}$ depletion of extracellular K^+ ions.

Figure 6 | Graphene deplete potassium at the cell/substrate cleft. **a**, Graphene-related G vibrational peak [61] was evaluated in wave number position for both SLG (left) and MLG substrates (right) with samples totally immersed in pure deuterium oxide (D_2O), in D_2O solution containing 4 mM KCl, and in D_2O solution containing 150 mM NaCl (see Supplementary Methods for technical details). The wavenumber maximum position relative to the G-peak Raman shift for control SLG sample was collocated at 1599 ± 0.5 cm⁻¹. It exhibited a change in shape associated to a G-peak position shift to 1600 ± 0.5 cm⁻¹ and to 1602 ± 0.5 cm⁻¹ in the presence of NaCl and KCl D_2O solutions, respectively (left, inset). Conversely, in MLG samples G peak maximum position did not change (right, inset). **b**, Sketch of the local amount of K^+

depletion in the membrane/surface cleft due to graphene trapping as function of cleft thickness. In light green the extrapolated values of such a distance (40 ± 100 nm) [34]. See Supplementary methods for technical details. **c**, Box plots summarize the average PSC frequency values (left) and the average PSC amplitudes ones (right) for neurons developed on glass control (in grey, $n = 21$), on glass supported SLG (in red, $n = 21$), on free-standing SLG (in green, $n = 26$) and on SLG deposited on ITO (in blue, $n = 8$). Note the significant increase in PSC frequency in SLG laying on insulating glass than Controls (3.11 ± 0.35 Hz vs. 1.72 ± 0.21 Hz, $P = 0.031$) or, even more, when grown on suspended SLG (4.22 ± 0.35 Hz vs. Controls, $P = 0.001$). SLG on conductive ITO does not change neuronal activity. Notably, in suspended SLG, PSCs amplitude is also significantly increased (59.2 ± 5.8 pA vs. 35.9 ± 4.9 pA, $P = 0.017$) when compared to control cultures. **d**, Hypothesis of Dirac point and Fermi level rearrangement as function of SLG supporting material (bottom), and an exemplification of the possible charge distribution in graphene layer as function of electrical characteristics of the underlying surface (top). Blue areas represent more positive regions (e.g. depletion of electrons), red areas represent more negative ones (e.g. persistency of electrons). Differences between the variables were assessed using one-way ANOVA and multiple comparisons were adjusted by Bonferroni correction.

METHODS

Substrate fabrication

SLG was CVD grown on ultra-flat Cu surfaces and transferred, as previously described [1], onto SiO₂ and Si₃N₄ substrates for subsequent Raman and XPS characterization. Briefly, after annealing the Cu foil in a 400 sccm :100 sccm argon and hydrogen atmosphere at 100 mbar and 1015 °C, a SLG layer is nucleated at 15 mbar with 0.2 sccm methane and closed by successively increasing the methane content to 0.5 sccm. The graphene is transferred using PMMA (PMMA 950K A2, MicroChem, USA) or PS (Polystyrene MW ~192k, Sigma-Aldrich, USA). For neuronal culturing, SLG was transferred to glass coverslips or indium tin oxide (ITO). Before the transfer procedure, hosting substrates were ultrasonicated in acetone and isopropanol to assure the required cleanness. Glass and ITO coverslips followed an additionally cleaning step in concentrated HCl overnight. MLG sheets were CVD grown on Ni ultra-flat surfaces as described previously [2] and transferred on hosting substrates following the same procedure adopted for SLG. Briefly, the Ni foil was annealed as described before, at a temperature of 900 °C. After annealing, a methane flow of 10 sccm at 50 mbar enabled to the diffusion of carbon into the foil, which then, during the following slow cooling step, precipitated to a layer of MLG on the surface.

Gold samples have been prepared starting from glass rectangular slides (24 mm x 12 mm, 0.2 mm thick), cleaned previously in hot Piranha solution (H₂SO₄:H₂O₂, 5:5 ratio in volume) in order to remove eventually present organic contaminants. Subsequently, 15 nm of Au were thermally evaporated at a rate of 0.5 Å/s. A thin adhesion layer of 5 nm of Cr was used in order to improve Au/glass adhesion. A quartz crystal microbalance was used as thickness control.

To obtain suspended graphene structures, graphene is transferred on patterned substrates obtained using OrmoComp[®] (micro resist technology, GmbH), a flexible

and biocompatible inorganic-organic material. The OrmoComp[®] substrates have been prepared on circular glasses (5 mm diameter, 0.12 mm thick), previously cleaned in hot Piranha solution (H₂SO₄:H₂O₂ 5:5 % v/v) in order to remove all organic contaminants. Subsequently, a poly-dimethylsiloxane (PDMS) master is prepared with replica molding process starting from a silicon stamp which is patterned with an array of parallel lines of width and periodicity of 10 µm and 20 µm, respectively. The PDMS master is used to press a drop of OrmoComp[®] on the circular glass in order to transfer the micropattern. Finally, the OrmoComp[®] is cured with UV light and the PDMS master is released. Commercially available single-layer CVD graphene on copper (GRAPHENEA – San Sebastián, Spain) is wet-transferred on the OrmoComp[®] substrates following the protocol described by Matruglio *et al.* [3]. Briefly, a layer of 250 nm of mr-I 7020 (a thermoplastic polymer of Micro Resist Technology GmbH) is used as sacrificial layer and spin coated on the graphene/Cu. The polymer/graphene/Cu membrane is placed in a copper etching solution (FeCl₃:H₂O 3:7 % v/v), etched overnight and finally washed in DI water in order to remove any residual due to the etching solution. The transfer of graphene is performed fishing the polymer/graphene/Cu membrane into the water directly on the OrmoComp[®] substrate. The water is left to evaporate at room temperature for 2 h, and mr-I 7020 is dissolved in cold acetone for 5 minutes. Critical point drying process is performed in order to avoid the collapse of the suspended structures.

Cell culture and electrophysiology

Isolation of primary brain tissue was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and the appropriate international and institutional

standards for the care and use of animals in research (Italian Ministry of Health, in agreement with the EU Recommendation 2007/526/CE). The protocols in this study and all performed experiments are approved by the local veterinary service and the institutional (SISSA) ethical committee, in accordance with the EU guidelines (2010/63/UE) and Italian law (decree 26/14).

Dissociated hippocampal cultures were obtained from neonatal rats (P0÷2) as previously described [4,5,6], and were plated on glass Control, SLG-, MLG- or Au-covered glass coverslips. As in our previous work with different carbon-based nanomaterial (e.g. carbon nanotubes, CNTs) [4,5,6], we did not pre-treat the culture substrates with any additional adhesion molecules, which might mask the effects of graphene. Cultured cells were incubated at 37 °C, 5% CO₂ in culture medium composed of Neurobasal-A (Thermo Fischer) containing B27 2% (Gibco) Glutamax 10 mM and Gentamycin 0.5 µM (Gibco), and used for experiments at 8÷10 days *in vitro* (DIV).

Somatic whole-cell patch clamp recordings were performed at room temperature (20÷22 °C) with pipettes (4÷7 MΩ) containing: 105 mM K gluconate, 20 mM KCl, 10 mM HEPES, 4 mM MgATP, 0.3 mM GTP, pH 7.35. The external saline solution contained: 150 mM NaCl, 4 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 10 mM HEPES, 10 mM Glucose, pH 7.4. Under voltage-clamp mode we measured the neuronal passive membrane properties: input resistance and cell capacitance did not significantly differ between the four groups (in Control 592 ± 51 MΩ, 74 ± 5 pF, n = 47; in SLG 664 ± 57 MΩ, 83 ± 4 pF, n = 54; in MLG 614 ± 74 MΩ, 85 ± 5 pF, n = 18; in Au 656 ± 65 MΩ, 80 ± 6 pF, n = 17). In voltage-clamp experiments, the holding potential (V_h) was -56 mV, not corrected for liquid junction potential, that was calculated to be -14

796 mV in our experimental conditions; the uncompensated value for series resistance
797 (R_s) was $< 8 \div 11 \text{ M}\Omega$.

798 Single spontaneous synaptic events (PSCs) and miniature PSCs (mPSCs) were
799 detected by the use of the AxoGraph X (Axograph Scientific) event detection
800 program and by the Clampfit 10 software (pClamp suite, Axon Instruments). On
801 average, ≥ 400 events were analysed from each cell in order to obtain mean
802 frequency and amplitude parameters. Glutamate AMPA-receptor and GABA_A-
803 receptor mediated PSCs were isolated offline by building two templates with different
804 kinetic parameters: respectively 0.1 ms rise-time; 3 and 30 ms decay time constant
805 (τ); 10 and 100 ms template length. Previous work [6,7] indicated that in our
806 experimental conditions, the vast majority of fast-decaying ($\tau < 5 \text{ ms}$) PSCs are
807 mediated by the glutamate AMPA-receptor type; while the slow-decaying ($\tau > 20 \text{ ms}$)
808 PSCs are mediated by the GABA_A-receptor type.

809 Current-voltage relations (I/V plots) were obtained by applying hyperpolarizing or
810 depolarizing voltage steps (15 steps of $\Delta V = 10 \text{ mV}$; 500 ms duration) from -110 mV
811 to $+30 \text{ mV}$ (values corrected for liquid junction potential) in the presence of $1 \text{ }\mu\text{M}$
812 Tetrodotoxin (TTX; Latoxan). A least square routine was fitted to the linear part of the
813 I/V curve, the slope of which was used to calculate leak conductance. Assuming that
814 a leak conductance is time and voltage independent, the I/V plot were corrected for
815 leak currents by subtracting the observed currents from the extrapolated leak
816 currents at the same level of test potential and the current values were then
817 normalized to the cell capacitance [8].

818 In current-clamp recordings, bridge balancing was continuously monitored and
819 adjusted. Action potentials (APs) were isolated off line by setting an appropriate
820 threshold voltage (10 mV). The fast voltage transients that crossed this value were

821 identified as APs and the spontaneous firing frequency for each neuron was
 822 calculated on a sample of at least 5 min of continuous recording keeping (by
 823 negative current injection) at -60 mV the resting membrane potential.
 824 At -60 mV resting membrane potential, the AP properties were experimentally
 825 determined by depolarizing (from 0 to 200 pA, in 20 pA increments) current steps
 826 (500 ms). The first AP produced by the current-clamp series was used for single AP
 827 measurements, including amplitude, threshold, duration at half-amplitude and
 828 maximal rising slope [9,10,11]. The threshold for firing was determined by measuring
 829 the voltage at the upward deflection of the trace, maximal AP amplitude was
 830 measured from threshold to the peak of the spike, the duration was measured at
 831 half-amplitude from threshold to peak and the maximal rising slope was measured as
 832 max dV/dt in the selected area of the voltage tracings (all measures performed by
 833 Clampfit; pClamp suite, 10.2 version; Axon Instruments). From this analysis, no
 834 significant difference was found between Control ($n = 21$) and SLG ($n = 19$) groups
 835 (amplitude: 56.4 ± 3.5 mV for Controls and 59 ± 3.1 mV in SLG, $P = 0.57$; width: 3.5
 836 ± 0.25 ms for Controls and 3.7 ± 0.38 ms for SLG, $P = 0.59$; threshold: -34.2 ± 1.5
 837 mV Control and -35.5 ± 1.2 mV in SLG, $P=0.34$; maximal rise slope: 61.6 ± 7.5
 838 mV/ms Control and 57.3 ± 5.8 mV/ms in SLG, $P= 0.32$).
 839 In evoked APs, the AHP was quantified over a window of 100 ms by calculating the
 840 area below or above the voltage curve, starting 20 ms after the beginning of the AP.
 841 AP discharge patterns were investigated by delivering depolarizing current steps (1
 842 s) of 200 pA while keeping the cells at -60 mV resting potential with steady
 843 intracellular current injection. “Adapting” and “tonic” responses were identified as
 844 previously described [12].

Beside the monitoring of the spontaneous firing frequency, all the current clamp experiments were carried out in presence of the synaptic blockers (all from Sigma) CNQX (10 μ M), Gabazine (5 μ M) and APV (50 μ M) added to the external solution. Current and voltage clamp responses were digitized at 20 kHz with the pCLAMP 10 software (Molecular Devices) and stored for further analysis.

Immunohistochemistry

Hippocampal neurons were fixed with 4% formaldehyde (prepared from fresh paraformaldehyde) in PBS for 20 min, permeabilized with 0.3% Triton-X-100 for 10 min and subsequently incubated with primary antibodies for 30 min at RT. After washing in PBS cultures were then incubated with secondary antibodies for 45 min and then mounted in *Vectashield* (Vector Laboratories) on 1 mm thick microscope glass slides. As primary antibodies were used rabbit polyclonal anti- β -tubulin III (Sigma T2200, 1:250 dilution), mouse monoclonal anti-GFAP (Sigma-Aldrich, 1:200 dilution), and guinea pig anti-vesicular glutamate transporter 1 (VGLUT1; Millipore, 1:2000). As secondary antibodies were used Alexa 594 goat anti rabbit (Invitrogen, dilution 1:500), Alexa 488 goat anti mouse (Invitrogen, dilution 1:500), and Alexa 488 goat anti guinea-pig (Invitrogen, 1:500). To stain cells nuclei, we used DAPI (Invitrogen, 1:200 dilution). To quantify cell density, images were acquired with an Epifluorescence Microscope (DM 6000, Leica; 10 $\times$ objective). We collected 10 fields (1000 μ m $\times$ 500 μ m) per coverslip and analysed fluorescence signals using ImageJ software (<http://rsb.info.nih.gov/ij/>).

To evaluate the orientation of the re-growing axons on the various substrates, we quantified their relative orientation based on the directionality analysis [13]. Briefly, the mean fibre's relative dispersion was computed from $n = 5$ randomly sampled

images per condition (Control, SLG, MLG and Au) where neuronal processes were visualized by class III β -tubulin immunofluorescence (Fig. 2a). The analysis was carried out using the Directionality plugin of Fiji software inferring the preferred orientation of “structures” present in the input image. Fibre orientation was calculated via a Fourier component analysis. We found no significant differences in the direction of the mean fibre’s dispersion among different conditions (Control = $30 \pm 9^\circ$; SLG = $29 \pm 7^\circ$; MLG = $38 \pm 8^\circ$; Au = $35 \pm 9^\circ$) indicating a negligible impact of the substrate on the orientation of the neuronal processes.

To quantify VGlut1 puncta, $n = 20 \pm 10$ z-stacks (acquired every $0.4 \mu\text{m}$) were taken from $n = 10$ randomly selected fields ($160 \mu\text{m} \times 80 \mu\text{m}$) per coverslip using an inverted confocal Microscope (Nikon Eclipse Ti-E; 40x oil immersion objective, 1.3 NA). We selected only VGlut1-positive puncta ($< 2 \mu\text{m}^3$) co-localized with β -tubulin III positive signal. For each image VGlut1 puncta were normalized to the β -tubulin III positive volume. Images were analysed using the Volocity software (Perkin Elmer). To highlight GABAergic neurons, cultures were stained with anti-GABA polyclonal primary antibody produced in rabbit (SIGMA, A2052; 1:500). To label the NKCC1 co-transporter, we used anti NKCC1 rabbit polyclonal primary antibody (Abcam; AB59791; $5 \mu\text{g/mL}$). Cultures were then stained with class III β -tubulin primary antibody produced in mouse (SIGMA; T5076; 1:500). As secondary antibody, we used AlexaFluor 488 goat anti rabbit (ThermoFisher A11034; 1:500), and AlexaFluor 594 goat anti mouse (ThermoFisher, A11032; 1:500).

To quantify the percentage of GABA-positive neurons, 10 ± 5 confocal z-stack (Nikon PlanFluor 40 $\times$ / 1.3 NA) were acquired from randomly selected fields and GABA-positive neurons were counted. This value was then normalized to the overall number of neurons (class III β -tubulin positive cells) for each field. To

quantify NKCC1 puncta, $n = 10$ z-stacks (acquired every $0.25\ \mu\text{m}$) were taken from selected fields ($106\ \mu\text{m} \times 106\ \mu\text{m}$) each group, using an inverted confocal Microscope (Nikon Eclipse Ti-E; Nikon Plan Apo Lambda 60x oil immersion objective, 1.4 NA). To quantify the amount of neuron-related NKCC1, only NKCC1-positive puncta in contact with the β -tubulin III signal were selected.

Imaging

For Cl^- imaging experiments, primary hippocampal cultures (DIV 8–10) were loaded with the fluorescent Chloride indicator MQAE (Abcam; ab145418) diluted in the standard extracellular solution at a final concentration of 1 mM for 10 min at $37\ ^\circ\text{C}$ in the cell culture incubator. Samples were then washed in the extracellular solution for 10 min at $37\ ^\circ\text{C}$. Samples were placed in a recording chamber mounted on an inverted microscope (Nikon Eclipse Ti-U) and observed with a $60\times$ objective (0.7 NA, PlanFluor, Nikon). Images (1024×1024 pixels) from fields containing 7 ± 4 neurons were acquired for 2 minutes at 5 Hz by a Hamamatsu Orca-Flash 4.0 digital camera, exciting the MQAE dye at the 365 nm Hg peak using a UV-2A Nikon filter set. Excitation light was attenuated by a neutral density filter (ND 16). Images of emitted fluorescence ($>420\ \text{nm}$) were displayed on a colour monitor controlled by an integrating imaging software package (HCImage, Hamamatsu) using a personal computer. Recorded images were analysed offline with the Clampfit software (pClamp suite, 10.2 version; Axon Instruments). Image time stacks were analysed in selected region of interest (ROI) to measure the variations in MQAE fluorescence intensity. Intracellular Cl^- transients were expressed as fractional amplitude variations ($\Delta F/F_0$, where F_0 is the baseline fluorescence level and ΔF is the change over the baseline); the onset time of

neuronal activation determined by detecting those events in the fluorescence signal that exceed at least five times the standard deviation of the noise. To elicit chloride influx/efflux through the membrane, an injection pipette (patch pipette with resistance of $1\div4$ M Ω , filled with 10 mM GABA diluted in the extracellular solution) was positioned at $20\div50$ μ m from the cell soma and connected to a pico-spritzer (PDES-02DX, npi Electronics) with 1 psi in-line pressure. 500 ms GABA puffs were delivered at fixed times. At the beginning of each experiment, a pipette containing pure saline was used to exclude artefacts due to the pressure injection (Supplementary Fig. 2). We found an overall percentage of 37 ± 6.5 % Control neurons and 35.6 ± 9.3 % SLG neurons that did not respond to the stimulation, and were excluded from further analysis.

Electron microscopy (EM)

Scanning EM imaging was conducted using collecting secondary electrons on a Gemini SUPRA 40 SEM (Carl Zeiss NTS GmbH, Oberkochen). Before SEM imaging, neuronal cells grown on the different substrates were fixed in 3% Glutaraldehyde in 0.1 M Sodium Cacodylate Buffer (pH 7.4), then dehydrated sequentially in ethanol solutions of 50, 75, 95, 99 and 100% (vol/vol in H₂O, 3 minutes each, 4 °C). After overnight drying in the fridge, and before imaging, samples were metalized with a 5 nm thick layer of platinum-iridium alloy using a metal sputter coater (Polaron SC7620). In order to prevent electron induced surface charging, low accelerating voltages ($1\div3$ keV) were used for cells visualization. SEM images of cells cross sections at membrane-substrate interface were obtained by focused ion beam (FIB) using a LEO-ZEISS Cross-Beam 1540 XB system. Gallium ion beam milling was performed with a current beam of 30 mA while SEM

images were collected at 3 kV. Samples were prepared following the same procedure described in the previous paragraph.

Substrate characterization

AFM topography data (MFP-3D, Asylum Research, Santa Barbara, California, USA) was acquired in tapping mode, using silicon cantilevers in ambient conditions. The roughness estimates were calculated using the standard deviation of elevation in mapped surface areas with sizes of 10 μm x 10 μm . XPS spectra were recorded in ultra-high vacuum conditions using a monochromatic SPECS XR-50 Mg K α X-Ray source ($E_{K\alpha}$ = 1253.6 eV) and a hemispherical energy analyser (Phoibos 100/150, Specs, Berlin, Germany). μ -Raman spectra were recorded with an in-house built system using an Ar-ion laser at 514.5 nm and operating with a spectral resolution of 0.75 cm^{-1} .

Raman measurements in aqueous conditions have been carried out at on the IUVS beamline at Elettra synchrotron radiation facility (Trieste, Italy). A complete description of the experimental apparatus can be found elsewhere [10.1016/j.nima.2012.11.037]. A 532 nm laser source, with a beam power near 5 mW, has been employed as excitation source. The scattered radiation was collected in a backscattering geometrical configuration. Slight modifications on the standard backscattering set-up have been introduced to allow measurements in liquid conditions (see Supplementary Methods). A 750 mm focal length Czerny-Turner spectrometer, equipped with an holographic reflection grating of 1800 g/mm and coupled with a Peltier-cooled back-thinned CCD, has been used to get the final Raman spectra.

Data Analysis, Statistics and Reproducibility

All values from samples subjected to the same experimental protocols were pooled together and expressed as histograms (mean $\pm$ SEM with n = biologically independent experiments, usually number of cells, unless otherwise indicated) or through box plot representation when one or more data set were found to follow a non normal distribution. In box plots, the thick horizontal bar indicates the median value, the boxed area extends from the 25th to 75th percentiles while whiskers from the 5th to the 95th percentiles. The homogeneity of variances was assessed through the Levene's test.

Statistically significant difference between two data sets was assessed by Student's t-test for parametric data and by Mann-Whitney for non-parametric ones. Differences between the logarithmic values of the analysed variables were assessed using one-way ANOVA [14] and multiple comparisons were adjusted by Bonferroni correction. Statistical significance was determined at $P < 0.05$, unless otherwise indicated. Significance was graphically indicated as follows: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Mathematical models of single neurons and neuronal networks

Full details on the mathematical models are reported in the Supplementary Information.

REFERENCES FOR METHODS

[1] Drieschner, S. et al. Frequency response of electrolyte-gated graphene electrodes and transistors. *Journal of Physics D: Applied Physics* 50, 095304 (2017).

- 995 [2] Drieschner, S. et al. High surface area graphene foams by chemical vapor
 996 deposition. 2D Materials 3, 045013 (2016).
- 997 [3] Matruglio, A. et al. Contamination-free suspended graphene structures by a
 998 Ti-based transfer method Carbon, 103, 305-310 (2016).
- 999 [4] Lovat, V. et al. Carbon nanotube substrates boost neuronal electrical
 1000 signaling. Nano Lett. 5, 1107-1110 (2005).
- 1001 [5] Cellot, G. et al. Carbon nanotubes might improve neuronal performance by
 1002 favouring electrical shortcuts. Nat. Nanotechnol. 4, 126-133 (2009).
- 1003 [6] Cellot, G. et al, Carbon nanotube scaffolds tune synaptic strength in cultured
 1004 neural circuits: novel frontiers in nanomaterial-tissue interactions. J. Neurosci.
 1005 31, 12945-12953 (2011).
- 1006 [7] Pampaloni, NP. et al. Sculpting neurotransmission during synaptic
 1007 development by 2D nanostructured interfaces. Nanomedicine. S1549-9634,
 1008 30082-5 (2017).
- 1009 [8] Sontheimer, H. & Ransom, C. Whole-Cell Patch-Clamp Recordings.
 1010 Neuromethods 35, 35-67 Humana Press Inc., (2007).
- 1011 [9] Ruscheweyh, R. & Sandkuhler, J. Lamina-specific membrane and discharge
 1012 properties of rat spinal dorsal horn neurones in vitro. J Physiol. 541, 231-244
 1013 (2002).
- 1014 [10] Chang, YM. & Luebke, JI. Electrophysiological diversity of layer 5 pyramidal
 1015 cells in the prefrontal cortex of the rhesus monkey: in vitro slice studies. J
 1016 Neurophysiol. 98, 2622-32. (2007)
- 1017 [11] Renganathan, M., Cummins, TR. & Waxman, SG. Contribution of Na(v)1.8
 1018 sodium channels to action potential electrogenesis in DRG neurons. J
 1019 Neurophysiol. 86, 629-40 (2001).

1020 [12] Furlan, F. et al. ERG conductance expression modulates the excitability of
 1021 ventral horn GABAergic interneurons that control rhythmic oscillations in the
 1022 developing mouse spinal cord. J. Neurosci. 27, 919-928 (2007).

1023 [13] Usmani S, et al. 3D meshes of carbon nanotubes guide functional
 1024 reconnection of segregated spinal explants. Sci Adv. e1600087 (2016).

1025 [14] Wilcox RR & Rousselet GA. A Guide to Robust Statistical Methods in
 1026 Neuroscience. Curr Protoc Neurosci. 82, 8.42.1-8.42.30 (2018).

1027

1028 **Additional information**

1029 Supplementary information are available in the online version of the paper.

1030 Correspondence and requests to Denis Scaini dscaini@sissa.it; Josè Antonio

1031 Garrido joseantonio.garrido@icn2.cat; Laura Ballerini laura.ballerini@sissa.it

1032

1033 **Data availability**

1034 The data that support the plots within this paper and other findings of this study

1035 are available from the corresponding authors upon reasonable request. The

1036 mathematical model is available online at ModelDB

1037 (<https://senselab.med.yale.edu/ModelDB/>).

a

b

c

d