

QUANTIFYING COMPLETION

ROBERT LOWEN and BART WINDELS

(Received 23 April 1999)

ABSTRACT. Approach uniformities were introduced in Lowen and Windels (1998) as the canonical generalization of both metric spaces and uniform spaces. This text presents in this new context of “quantitative” uniform spaces, a reflective completion theory which generalizes the well-known completions of metric and uniform spaces. This completion behaves nicely with respect to initial structures and hyperspaces. Also, continuous extensions of pseudo-metrics on uniform spaces and (real) compactification of approach spaces can be interpreted in terms of this completion.

Keywords and phrases. Completion, uniform space, approach space, approach uniformity, compactification, realcompactification.

2000 Mathematics Subject Classification. Primary 54E15, 18B30, 54B20.

1. Approach uniformities. The motivation for the introduction of approach uniformities (which is the subject of Lowen and Windels [6]) is twofold. On the one hand, the category **AP** of approach spaces (see Lowen [3]) is not fit to handle uniform concepts. Different results concerning completeness in the category **AP** imply that the theory is essentially *local*. For instance, the space of continuous functions between metric spaces, equipped with the pointwise distance, turns out to be complete. On the other hand, **AP** seems not to be the right context for the quantification of uniform properties, such as completeness and total boundedness, whereas for topological concepts everything works out quite well in **AP**. Therefore, it was natural to seek a new category that combines the quantitative aspects of **AP** (or **pMET**) and qualitative uniform concepts. To that end the concept of *approach uniformities* was introduced in [6].

For the reader's convenience we briefly recall the main definitions and results. An approach uniform space (X, Γ) is a set X together with an ideal Γ of functions from $X \times X$ into $[0, \infty]$, satisfying the following conditions:

$$(AU1) \quad \forall \gamma \in \Gamma, \forall x \in X : \gamma(x, x) = 0.$$

$$(AU2) \quad \forall \xi \in [0, \infty]^{X \times X} : (\forall \varepsilon > 0, \forall N < \infty : \exists \gamma \in \Gamma \text{ s.t. } \xi \wedge N \leq \gamma^N + \varepsilon) \Rightarrow \xi \in \Gamma.$$

$$(AU3) \quad \forall \gamma \in \Gamma, \forall N < \infty, \exists \gamma^N \in \Gamma \text{ s.t. } \forall x, y, z \in X : \gamma(x, z) \wedge N \leq \gamma^N(x, y) + \gamma^N(y, z).$$

$$(AU4) \quad \forall \gamma \in \Gamma : \gamma^s \in \Gamma.$$

Equivalently, an approach uniform space can be described by means of a *uniform tower*, i.e., a family of filters $(\mathfrak{U}_\varepsilon)_{\varepsilon \in \mathbb{R}^+}$ on $X \times X$, such that

$$(UT1) \quad \forall \varepsilon \in \mathbb{R}^+, \forall U \in \mathfrak{U}_\varepsilon : \Delta_X \subset U.$$

$$(UT2) \quad \forall \varepsilon \in \mathbb{R}^+, \forall U \in \mathfrak{U}_\varepsilon : U^{-1} \in \mathfrak{U}_\varepsilon.$$

$$(UT3) \quad \forall \varepsilon, \varepsilon' \in \mathbb{R}^+ : \mathfrak{U}_\varepsilon \circ \mathfrak{U}_{\varepsilon'} \supset \mathfrak{U}_{\varepsilon + \varepsilon'}.$$

$$(UT4) \quad \forall \varepsilon \in \mathbb{R}^+ : \mathfrak{U}_\varepsilon = \bigcup_{\alpha > \varepsilon} \mathfrak{U}_\alpha.$$

or equivalently, a family $(\mathfrak{U}_\varepsilon)_{\varepsilon \in \mathbb{R}^+}$ of semi-uniformities, satisfying (UT3) and (UT4).

For example, if d is a pseudo-metric, then the collection $\Gamma(d) := \{\gamma \mid \gamma \leq d\}$ is an approach uniformity. It is referred to as the *metric approach uniformity* induced by d .

If \mathfrak{U} is a uniformity, then the trivial tower $(\mathfrak{U})_\varepsilon$ (\mathfrak{U} on every level ε), is a uniform tower, defining an approach uniformity $\Gamma(\mathfrak{U})$, which is referred to as the *uniform approach uniformity* induced by \mathfrak{U} . If (X, Γ) and (Y, Ψ) are approach uniform spaces, then a function $f : (X, \Gamma) \rightarrow (Y, \Psi)$ is called a *uniform contraction* if and only if $\forall \psi \in \Psi : \psi \circ (f \times f) \in \Gamma$.

The category **AUnif** of approach uniform spaces and uniform contractions is a topological category. It contains **Unif** both reflectively and coreflectively and **pMET** coreflectively.

Approach uniformities establish a context for quantifying uniform concepts. In [5], a measure of total boundedness (and precompactness), of completeness and of uniform connectedness is presented.

For every approach uniformity (X, Γ) and for any $x \in X$ we can consider the set

$$\mathfrak{A}(x) := \{\gamma(x, \cdot) \mid \gamma \in \Gamma\} \subset [0, \infty]^X. \tag{1.1}$$

The family $(\mathfrak{A}(x))_{x \in X}$ defines an ordinary approach structure on X , which we shall call the *underlying approach structure* of Γ . This procedure yields a forgetful functor $A : \mathbf{AUnif} \rightarrow \mathbf{AP}$.

Also recall that in any approach space $(X, (\mathfrak{A}(x))_{x \in X})$ and for any filter \mathfrak{f} on X and any $x \in X$, we denote

$$\lambda_{\mathfrak{f}}(x) := \sup_{\varphi \in \mathfrak{A}(x)} \inf_{F \in \mathfrak{f}} \sup_{y \in F} \varphi(y). \tag{1.2}$$

2. Cauchy filters. Let X be a set, and let $U \subset X \times X$. Then we denote

$$\theta_U : X \times X \rightarrow [0, \infty] : \begin{cases} (x, y) \mapsto 0 & \text{if } (x, y) \in U, \\ (x, y) \mapsto 1 & \text{if } (x, y) \notin U. \end{cases} \tag{2.1}$$

PROPOSITION 2.1. *Let (X, Γ) be an approach uniform space, and let \mathfrak{f} be a filter on X . Then the following are equivalent.*

- (1) \mathfrak{f} is Cauchy with respect to the uniform coreflection.
- (2) $\forall \gamma \in \Gamma, \forall \varepsilon > 0, \exists F \in \mathfrak{f} : \theta_{F \times F} + \varepsilon > \gamma$.

PROOF. \mathfrak{f} is Cauchy with respect to the uniform coreflection \mathfrak{U}_0 if and only if

$$\begin{aligned} \forall U \in \mathfrak{U}_0, \exists F \in \mathfrak{f} : F \times F \subset U \\ \iff \forall \gamma \in \Gamma, \forall \varepsilon > 0, \exists F \in \mathfrak{f} : F \times F \subset \{\gamma < \varepsilon\}. \\ \iff \forall \gamma \in \Gamma, \forall \varepsilon > 0, \exists F \in \mathfrak{f} : \theta_{F \times F} + \varepsilon > \gamma. \quad \square \end{aligned} \tag{2.2}$$

A filter on (X, Γ) is called (Γ) -Cauchy if and only if it satisfies the conditions of the previous proposition.

DEFINITION 2.2. Two filters \mathfrak{f} and \mathfrak{g} on (X, Γ) are called *equivalent* if and only if $\mathfrak{f} \cap \mathfrak{g}$ is Cauchy.

In Lowen [3] Cauchy filters in approach spaces are introduced: a filter \mathfrak{f} on X is called Cauchy with respect to the approach space $(X, (\mathfrak{A}(x))_x)$ if and only if $\inf_{x \in X} \lambda \mathfrak{f}(x) = 0$. This definition yields a categorically nice completion theory in **AP** (see [3]).

We have the following relationship between **AP**-Cauchy filters and **AUnif**-Cauchy filters.

PROPOSITION 2.3. *Let (X, Γ) be an approach uniform space, and let $(X, A(\Gamma))$ denote its underlying approach space. For any filter \mathfrak{f} on X , we have*

$$\mathfrak{f} \text{ is } A(\Gamma)\text{-Cauchy} \implies \mathfrak{f} \text{ is } \Gamma\text{-Cauchy.} \tag{2.3}$$

PROOF. \mathfrak{f} is $A(\Gamma)$ -Cauchy if and only if $\inf_{x \in X} \lambda \mathfrak{f}(x) = 0$, i.e., if and only if

$$\forall \varepsilon > 0, \exists x \in X, \forall \gamma \in \Gamma, \exists F \in \mathfrak{f}, \forall \gamma \in F : \gamma(x, \gamma) < \varepsilon. \tag{2.4}$$

Let $\gamma \in \Gamma$ and $\varepsilon > 0$. Take $\gamma^N \in \Gamma$, γ^N symmetric, such that $\forall x, \gamma, z \in X : \gamma(x, z) \wedge N \leq \gamma^N(x, \gamma) + \gamma^N(\gamma, z)$ for some $N > \varepsilon$, and take $x_0 \in X$ and $F \in \mathfrak{f}$ such that $\forall \gamma \in F : \gamma^N(x_0, \gamma) < \varepsilon/2$. Clearly, we have $\forall \gamma, z \in F$:

$$\gamma(\gamma, z) \wedge N \leq \gamma^N(\gamma, x_0) + \gamma^N(x_0, z) \leq \varepsilon. \tag{2.5}$$

Consequently,

$$\forall \gamma \in \Gamma, \forall \varepsilon > 0, \exists F \in \mathfrak{f}, \forall x, \gamma \in F : \gamma(x, \gamma) \leq \varepsilon, \tag{2.6}$$

which means that \mathfrak{f} is Γ -Cauchy. □

PROPOSITION 2.4. *Let (X, Γ) be an approach uniformity, and let \mathfrak{f} be a Cauchy filter on X . Then $\mathfrak{B} := \{\{\gamma \mid \inf_{x \in F} \gamma(x, \gamma) < \alpha\} \mid \gamma \in \Gamma, \alpha > 0, F \in \mathfrak{f}\}$ is a filterbasis for a Cauchy filter \mathfrak{M} , coarser than \mathfrak{f} . Moreover, \mathfrak{M} is the coarsest Cauchy filter with that property. Consequently, \mathfrak{M} and \mathfrak{f} are equivalent.*

PROOF. Let \mathfrak{U}_0 denote the uniform coreflection of Γ . The proof becomes folklore, if one observes that

$$\mathfrak{B} = \{U(F) \mid U \in \mathfrak{U}_0, F \in \mathfrak{f}\}. \tag{2.7}$$

□

DEFINITION 2.5. An approach uniformity (X, Γ) is called complete if and only if every Γ -Cauchy filter converges (with respect to the underlying topology of the uniform coreflection of Γ).

Thus an approach uniformity is called complete if and only if its uniform coreflection is complete.

By Proposition 2.3, we know that if (X, Γ) is **AUnif**-complete, then $(X, A(\Gamma))$ is **AP**-complete. The converse need not be true.

EXAMPLE 2.6. Let $(E, \|\cdot\|)$ be a non-reflexive Banach space. If E is equipped with the initial **AP**-structure δ for the source

$$(f : E \longrightarrow \mathbb{R})_{f \in E^*, \|f\|_{E^*} \leq 1}, \tag{2.8}$$

(where \mathbb{R} is equipped with the usual distance), then E is complete in **AP**.

However, if E is equipped with the initial **AUnif**-structure Γ for the source

$$(f : E \rightarrow \mathbb{R})_{f \in E^*, \|f\|_{E^*} \leq 1}, \tag{2.9}$$

(where \mathbb{R} is equipped with the usual approach uniformity), then E is *not* complete in **AUnif**.

Indeed, we know that $\forall x \in E, \forall A \subset E$

$$\delta(x, A) = \sup_{F \in 2(E^*)} \inf_{\mathcal{Y} \in A} \sup_{f \in F} |f(x) - f(y)|, \tag{2.10}$$

and therefore the metric coreflection d_δ is given by $\forall x, y \in E$

$$d_\delta(x, y) = \sup_{F \in 2(E^*)} \sup_{f \in F} |f(x) - f(y)| = \|x - y\|. \tag{2.11}$$

Since E is Banach, (E, δ) is complete.

By [6, Proposition 2.16] we know that the uniform coreflection \mathfrak{U}_0 of Γ is initial for the **Unif**-source

$$(f : E \rightarrow \mathbb{R})_{f \in E^*, \|f\|_{E^*} \leq 1}, \tag{2.12}$$

(where \mathbb{R} is equipped with the usual uniformity). Let $B := \{x \in E \mid \|x\| \leq 1\}$. Then $f(B) \subset [-1, 1]$, which is totally bounded. Thus B is totally bounded. However, B is not compact, since E is non-reflexive, hence B is not complete. Therefore (E, Γ) is not complete.

3. Completion. By Proposition 2.4, it makes sense to define \hat{X} to be the set of all minimal Cauchy filters on X .

PROPOSITION 3.1. *Let (X, Γ) be an approach uniformity. If $\gamma \in \Gamma$, define $\forall \mathfrak{W}, \mathfrak{N} \in \hat{X}$*

$$\hat{\gamma}(\mathfrak{W}, \mathfrak{N}) := \inf_{F \in \mathfrak{W} \cap \mathfrak{N}} \sup_{x, y \in F} \gamma(x, y). \tag{3.1}$$

Then $\hat{\Gamma} := \{\{\hat{\gamma} \mid \gamma \in \Gamma, \gamma \text{ symmetric}\}\}$ is an approach uniformity on \hat{X} .

PROOF. The fact that $\hat{\Gamma}$ is an ideal follows from the observation that $\forall \gamma_1, \gamma_2 \in \Gamma : \hat{\gamma}_1 \vee \hat{\gamma}_2 \leq \widehat{\gamma_1 \vee \gamma_2}$.

(AU1) Let $\gamma \in \Gamma$ and $\mathfrak{W} \in \hat{X}$. Since \mathfrak{W} is Cauchy, we have that $\forall \varepsilon > 0, \exists F \in \mathfrak{W} : \theta_{F \times F} + \varepsilon \geq \gamma$. Therefore $\hat{\gamma}(\mathfrak{W}, \mathfrak{W}) = \inf_{F \in \mathfrak{W}} \sup_{x, y \in F} \gamma(x, y) = 0$.

(AU2) This is obvious.

(AU3) Let $\gamma \in \Gamma$ and $N < \infty$. Then there is some $\gamma^N \in \Gamma$ such that $\forall x, y, z \in X : \gamma(x, z) \wedge N \leq \gamma^N(x, y) + \gamma^N(y, z)$. We shall prove that $\forall \mathfrak{W}, \mathfrak{N}, \mathfrak{F} \in \hat{X} : \hat{\gamma}(\mathfrak{W}, \mathfrak{N}) \wedge N \leq \hat{\gamma}^N(\mathfrak{W}, \mathfrak{F}) + \hat{\gamma}^N(\mathfrak{F}, \mathfrak{N})$. Let $F_1 \in \mathfrak{F} \cap \mathfrak{W}$ and $F_2 \in \mathfrak{F} \cap \mathfrak{N}$. Then $F := F_1 \cup F_2 \in \mathfrak{W} \cap \mathfrak{N}$. Since $F_1 \cap F_2 \neq \emptyset$, we have that

$$\sup_{x, y \in F} \gamma(x, y) \wedge N \leq \sup_{x, z \in F_1} \gamma^N(x, z) + \sup_{z, y \in F_2} \gamma^N(z, y), \tag{3.2}$$

and therefore

$$\inf_{F \in \mathfrak{W} \cap \mathfrak{N}} \sup_{x, y \in F} \gamma(x, y) \wedge N \leq \inf_{F_1 \in \mathfrak{F} \cap \mathfrak{W}} \sup_{x, z \in F_1} \gamma^N(x, z) + \inf_{F_2 \in \mathfrak{F} \cap \mathfrak{N}} \sup_{z, y \in F_2} \gamma^N(z, y). \tag{3.3}$$

(AU4) This follows from the observation that $\forall \gamma \in \Gamma : \hat{\gamma}^s = \hat{\gamma}^s$. □

The approach uniformity $(\hat{X}, \hat{\Gamma})$ in the previous proposition is called the *completion* of (X, Γ) . We have the following obvious relationship between the completion of the uniform coreflection, and the uniform coreflection of the completion.

PROPOSITION 3.2. *If $(\mathfrak{U}_\varepsilon)_\varepsilon$ and $(\hat{\mathfrak{U}}_\varepsilon)_\varepsilon$ are the towers of (X, Γ) and $(\hat{X}, \hat{\Gamma})$ respectively, then $(\hat{\mathfrak{U}})_0$ coincides with the uniform completion $\hat{\mathfrak{U}}_0$ of \mathfrak{U}_0 .*

PROOF. We see that

$$\begin{aligned} (\hat{\mathfrak{U}})_0 &= \langle \{ \hat{\gamma} < \alpha \mid \hat{\gamma} \in \hat{\Gamma}, \alpha > 0 \} \rangle \\ &= \langle \{ (\mathfrak{M}, \mathfrak{N}) \mid \exists F \in \mathfrak{M} \cap \mathfrak{N}, \forall a, b \in F : \gamma(a, b) < \alpha \} \rangle \\ &= \langle \{ (\mathfrak{M}, \mathfrak{N}) \mid \exists F \in \mathfrak{M} \cap \mathfrak{N} : F \times F \subset \{ \gamma < \alpha \} \} \rangle \\ &= \hat{\mathfrak{U}}_0. \end{aligned} \tag{3.4} \quad \square$$

From now on we can write $\hat{\mathfrak{U}}_0$ without ambiguity.

Before we move on, we check that completion behaves nicely with respect to bases. An ideal basis \mathfrak{D} in $[0, \infty]^{X \times X}$ is called a basis for Γ , if \mathfrak{D} satisfies (AU1), (AU2), and (AU4) and

$$\Gamma = \langle \mathfrak{D} \rangle := \{ \xi \in [0, \infty]^{X \times X} \mid \forall \varepsilon > 0, \forall N < \infty : \exists \gamma_\varepsilon^N \in \mathfrak{D} : \xi \wedge N \leq \gamma_\varepsilon^N + \varepsilon \}. \tag{3.5}$$

PROPOSITION 3.3. *If $\langle \mathfrak{D} \rangle = \Gamma$, then $\langle \{ \hat{d} \mid d \in \mathfrak{D} \} \rangle = \hat{\Gamma}$.*

PROOF. This is a direct consequence of the fact that $\forall \gamma, \xi \in [0, \infty]^{X \times X}, \forall \varepsilon > 0, \forall N < \infty$:

$$\xi \wedge N \leq \gamma + \varepsilon \implies \hat{\xi} \wedge N \leq \hat{\gamma} + \varepsilon. \tag{3.6} \quad \square$$

If $x \in X$, then let \mathfrak{V}_x denote the neighbourhoodfilter of x with respect to the underlying topology of the uniform coreflection.

PROPOSITION 3.4. *Let (X, Γ) be an approach uniformity. Then*

$$i : (X, \Gamma) \longrightarrow (\hat{X}, \hat{\Gamma}) \quad x \longmapsto \mathfrak{V}_x \tag{3.7}$$

is initial, and $i(X)$ is dense in \hat{X} . Moreover, if (X, Γ) is Hausdorff, then i is an embedding.

PROOF. It is well-known (cf. [10]) that $\forall x \in X : \mathfrak{V}_x$ is a minimal Cauchy filter, and therefore i is well-defined; and also, that $i(X)$ is dense in \hat{X} .

In order to show that Γ is the initial structure, first notice that if $\gamma \in \Gamma$, then $\forall x, y \in X$:

$$\hat{\gamma} \circ (i \times i)(x, y) = \hat{\gamma}(\mathfrak{V}_x, \mathfrak{V}_y) = \inf_{F \in \mathfrak{V}_x \cap \mathfrak{V}_y} \sup_{a, b \in F} \gamma(a, b). \tag{3.8}$$

Let $\gamma \in \Gamma$ be bounded. Let $\tilde{\gamma} \in \Gamma$ be such that $\forall x, y, z, u \in X : \gamma(x, u) \leq \tilde{\gamma}(x, y) + \tilde{\gamma}(y, z) + \tilde{\gamma}(z, u)$. We shall prove that

- (1) $\gamma \leq \hat{\gamma} \circ (i \times i)$.
- (2) $\hat{\gamma} \circ (i \times i) \leq \tilde{\gamma}$.

PROOF OF (1). Let $x, y \in X$. Suppose $\hat{\gamma} \circ (i \times i)(x, y) < M \in \mathbb{R}^+$, then there is some $F \in \mathfrak{V}_x \cap \mathfrak{V}_y$ such that $\forall a, b \in F : \gamma(a, b) < M$. Since $\{x, y\} \subset F$, this means that $\gamma(x, y) < M$. □

PROOF OF (2). Let $x, y \in X$ and $\varepsilon > 0$. Put $F := \{z \in X \mid \tilde{y}(x, z) < \varepsilon/2 \text{ or } \tilde{y}(y, z) < \varepsilon/2\} \in \mathfrak{V}_x \cap \mathfrak{V}_y$. Let $a, b \in F$.

- If $\tilde{y}(x, a) < \varepsilon/2$ and $\tilde{y}(x, b) < \varepsilon/2$, then $\gamma(a, b) \leq \tilde{y}(a, x) + \tilde{y}(x, b) < \varepsilon$.
- If $\tilde{y}(y, a) < \varepsilon/2$ and $\tilde{y}(y, b) < \varepsilon/2$, then also $\gamma(a, b) < \varepsilon$.
- If $\tilde{y}(x, a) < \varepsilon/2$ and $\tilde{y}(y, b) < \varepsilon/2$, then $\gamma(a, b) \leq \tilde{y}(a, x) + \tilde{y}(x, y) + \tilde{y}(y, b) \leq \tilde{y}(x, y) + \varepsilon$.
- If $\tilde{y}(x, b) < \varepsilon/2$ and $\tilde{y}(y, a) < \varepsilon/2$, then $\gamma(a, b) \leq \tilde{y}(x, y) + \varepsilon$ too. □

In any case $\sup_{a, b \in F} \gamma(a, b) \leq \tilde{y}(x, y) + \varepsilon$, and thus $\hat{y} \circ (i \times i) \leq \tilde{y} + \varepsilon$, which by arbitrariness of ε proves the claim.

If (X, Γ) is Hausdorff, then for all $x \neq y : \mathfrak{V}_x \neq \mathfrak{V}_y$, and therefore i is an embedding. □

Now we have to prove that a completion really is complete.

PROPOSITION 3.5. *Let (X, Γ) be an approach uniformity. Then $(\hat{X}, \hat{\Gamma})$ is a complete approach uniform space.*

PROOF. This is a consequence of Proposition 3.2. □

PROPOSITION 3.6. *Let (Y, Ψ) be a Hausdorff complete approach uniformity. If $f : (X, \Gamma) \rightarrow (Y, \Psi)$ is a uniform contraction, then there is a unique uniform contraction $\hat{f} : (\hat{X}, \hat{\Gamma}) \rightarrow (Y, \Psi)$ such that $\hat{f} \circ i = f$.*

PROOF. If $(\mathfrak{U}_\varepsilon)_\varepsilon$ and $(\mathfrak{V}_\varepsilon)_\varepsilon$ are the towers of Γ and Ψ respectively, then $f : (X, \mathfrak{U}_0) \rightarrow (Y, \mathfrak{V}_0)$ is uniformly continuous and (Y, \mathfrak{V}_0) is a Hausdorff complete uniform space. It is well-known that there is a unique $\hat{f} : (\hat{X}, \hat{\mathfrak{U}}_0) \rightarrow (Y, \mathfrak{V}_0)$ uniformly continuous such that $\hat{f} \circ i = f$. In fact we know that for all $\mathfrak{M} \in \hat{X} : \hat{f}(\mathfrak{M}) = \lim f(\mathfrak{V}_{\mathfrak{M}} \mid X)$. We shall prove that \hat{f} is a uniform contraction. To that end, let $\psi \in \Psi$; we shall show that $\psi \circ (\hat{f} \times \hat{f}) \in \hat{\Gamma}$.

Recall that if \mathfrak{M} is a Cauchy-filter, then

$$\forall \gamma \in \Gamma, \forall \alpha > 0, \forall F \in M, \exists x \in F, \exists M \in \mathfrak{M} \cap \mathfrak{V}_x, \forall a, b \in M : \gamma(a, b) < \alpha. \quad (3.9)$$

Let $\mathfrak{M}, \mathfrak{N} \in \hat{X}$ and let $\varepsilon > 0$. We have that $f(\mathfrak{V}_{\mathfrak{M}} \mid X) \rightarrow \hat{f}(\mathfrak{M})$, i.e., $f(\mathfrak{V}_{\mathfrak{M}} \mid X) \supset \mathfrak{N}(\hat{f}(\mathfrak{M}))$. Since

$$\begin{aligned} f(\mathfrak{V}_{\mathfrak{M}} \mid X) &= \langle \{ \{ f(x) \mid \tilde{y}(\mathfrak{M}, \mathfrak{V}_x) < \alpha \} \mid \gamma \in \Gamma, \alpha > 0 \} \rangle \\ \mathfrak{N}(\hat{f}(\mathfrak{M})) &= \langle \{ \{ \gamma \mid \psi(\hat{f}(\mathfrak{M}), \gamma) < \beta \} \mid \psi \in \Psi, \beta > 0 \} \rangle, \end{aligned} \quad (3.10)$$

we find that

$$\forall \beta > 0, \forall \psi \in \Psi, \exists \alpha_1 > 0, \exists \gamma_1 \in \Gamma, \forall x \in X : \gamma_1(\mathfrak{M}, \mathfrak{V}_x) < \alpha_1 \implies \psi(\hat{f}(\mathfrak{M}), f(x)) < \beta. \quad (3.11)$$

Analogously we have that

$$\forall \beta > 0, \forall \psi \in \Psi, \exists \alpha_2 > 0, \exists \gamma_2 \in \Gamma, \forall x \in X : \gamma_2(\mathfrak{N}, \mathfrak{V}_x) < \alpha_2 \implies \psi(\hat{f}(\mathfrak{N}), f(x)) < \beta. \quad (3.12)$$

Let $\tilde{\psi} \in \Psi$ such that $\forall x, y, z, u \in Y : \psi(x, u) \leq \tilde{\psi}(x, y) + \tilde{\psi}(y, z) + \tilde{\psi}(z, u)$. Then we obtain that there exist $\alpha > 0$ and $\gamma \in \Gamma$ such that for all $x \in X$:

$$\hat{\gamma}(\mathfrak{M}, \mathfrak{V}_x) < \alpha \implies \tilde{\psi}(\hat{f}(\mathfrak{M}), f(x)) < \frac{\varepsilon}{2}, \quad \hat{\gamma}(\mathfrak{N}, \mathfrak{V}_x) < \alpha \implies \tilde{\psi}(\hat{f}(\mathfrak{N}), f(x)) < \frac{\varepsilon}{2}. \quad (3.13)$$

From (3.9) we deduce that $\forall F \in \mathfrak{M} \cap \mathfrak{N}$:

$$\exists x_1 \in F : \hat{\gamma}(\mathfrak{M}, \mathfrak{V}_{x_1}) < \alpha, \quad \exists x_2 \in F : \hat{\gamma}(\mathfrak{N}, \mathfrak{V}_{x_2}) < \alpha, \quad (3.14)$$

and thus

$$\exists x_1 \in F : \tilde{\psi}(\hat{f}(\mathfrak{M}), f(x_1)) < \frac{\varepsilon}{2}, \quad \exists x_2 \in F : \tilde{\psi}(\hat{f}(\mathfrak{N}), f(x_2)) < \frac{\varepsilon}{2}. \quad (3.15)$$

Then

$$\begin{aligned} \psi(\hat{f}(\mathfrak{M}), \hat{f}(\mathfrak{N})) &\leq \tilde{\psi}(\hat{f}(\mathfrak{M}), f(x_1)) + \tilde{\psi}(f(x_1), f(x_2)) + \tilde{\psi}(f(x_2), \hat{f}(\mathfrak{N})) \\ &\leq \tilde{\psi}(f(x_1), f(x_2)) + \varepsilon \leq \sup_{a, b \in F} \tilde{\psi}(f(a), f(b)) + \varepsilon. \end{aligned} \quad (3.16)$$

Thus

$$\psi \circ (\hat{f} \times \hat{f})(\mathfrak{M}, \mathfrak{N}) \leq \inf_{F \in \mathfrak{M} \cap \mathfrak{N}} \sup_{a, b \in F} \tilde{\psi} \circ (f \times f)(a, b) + \varepsilon. \quad (3.17)$$

Since $\tilde{\psi} \circ (f \times f) \in \Gamma$ and by arbitrariness of ε , we conclude that $\psi \circ (\hat{f} \times \hat{f}) \in \hat{\Gamma}$. \square

Let \mathbf{cAUnif} denote the full subcategory of \mathbf{AUnif} consisting of all complete Hausdorff approach uniformities.

COROLLARY 3.7. *\mathbf{cAUnif} is a reflective subcategory of \mathbf{AUnif} .*

4. Examples

4.1. Extensions of pseudo-metrics. Any collection \mathfrak{D} of pseudo-metrics that is closed under the formation of finite suprema, is a basis for some approach uniformity Γ . In that case, we call \mathfrak{D} a *uniform gauge* for Γ . We already noticed [Proposition 3.3] that if \mathfrak{D} is a uniform gauge for Γ , then $\tilde{\mathfrak{D}} := \{\hat{d} \mid d \in \mathfrak{D}\}$ is a basis for $\hat{\Gamma}$. As a matter of fact, $\tilde{\mathfrak{D}}$ is a uniform gauge too. Moreover, $\tilde{\mathfrak{D}}$ consists of the unique uniformly continuous extensions of the pseudo-metrics in \mathfrak{D} from $X \times X$ to $\hat{X} \times \hat{X}$.

PROPOSITION 4.1. *Let (X, Γ) be an approach uniformity and let $d \in \Gamma$ be a pseudo-metric. Then $\forall \mathfrak{M}, \mathfrak{N} \in \hat{X}$:*

$$\hat{d}(\mathfrak{M}, \mathfrak{N}) = \sup_{M \in \mathfrak{M}} \sup_{N \in \mathfrak{N}} d(M, N). \quad (4.1)$$

PROOF. Let $M \in \mathfrak{M}$, $N \in \mathfrak{N}$, $F \in \mathfrak{M} \cap \mathfrak{N}$. Since $N \cap F \neq \emptyset$ and $M \cap F \neq \emptyset$ it is apparent that

$$\forall \varepsilon > 0, \exists p \in N, \exists q \in M : d(p, q) \leq \sup_{x, y \in F} d(x, y) + \varepsilon. \quad (4.2)$$

This is equivalent to the fact that

$$\inf_{p \in N} \inf_{q \in M} d(p, q) \leq \sup_{x, y \in F} d(x, y) \quad (4.3)$$

and therefore we have that

$$\sup_{M \in \mathfrak{M}} \sup_{N \in \mathfrak{N}} \inf_{p \in N} \inf_{q \in M} d(p, q) \leq \inf_{F \in \mathfrak{M} \cap \mathfrak{N}} \sup d(x, y). \quad (4.4)$$

Conversely, let $\varepsilon > 0$ and choose $M \in \mathfrak{M}$ and $N \in \mathfrak{N}$ such that $\sup_{x, y \in M} d(x, y) < \varepsilon/2$ and $\sup_{x, y \in N} d(x, y) < \varepsilon/2$. If $F := M \cup N$, then $\forall x, y \in F$, $\forall p \in M$, $\forall q \in N$: $d(x, y) \leq d(p, q) + \varepsilon$ since

- If $(x, y) \in N \times N$ or $(x, y) \in M \times M$ then this is trivial.
- If $x \in M$ and $y \in N$, then

$$d(x, y) \leq d(x, p) + d(p, q) + d(q, y) \leq \frac{\varepsilon}{2} + d(p, q) + \frac{\varepsilon}{2}. \quad (4.5)$$

Thus

$$\inf_{F \in \mathfrak{M} \cap \mathfrak{N}} \sup_{x, y \in F} d(x, y) \leq \sup_{M \in \mathfrak{M}} \sup_{N \in \mathfrak{N}} \inf_{p \in N} \inf_{q \in M} d(p, q). \quad (4.6)$$

□

4.2. Metric and uniform approach uniformities. The categories **pMET** and **Unif** also allow a nice completion theory (cf. [8] for metric spaces, [10] for uniform spaces). As a matter of fact, the completion theory in **AUnif** generalizes these constructions.

PROPOSITION 4.2. (a) *Let (X, Γ) be a metric approach uniform space, $\Gamma = \Gamma(d)$. Then $\hat{\Gamma} = \Gamma(\hat{d})$.*

(b) *Let (X, Γ) be a uniform approach uniform space, $\Gamma = \Gamma(\mathfrak{U})$. Then $\hat{\Gamma} = \Gamma(\hat{\mathfrak{U}})$.*

PROOF. Part (a) is a special case of Proposition 4.1. Part (b) is a direct consequence of Proposition 3.2. □

4.3. Initial structures. In this section, we show that, as in **Unif**, completion commutes with the formation of subspaces and products.

PROPOSITION 4.3. *Let $(f_i : X \rightarrow Y_i)_{i \in I}$ be an initial **AUnif**-source.*

(1) *The source of canonical extensions $(\hat{f}_i : \hat{X} \rightarrow \hat{Y}_i)_{i \in I}$ is initial too.*

(2) *If $j_i : Y_i \rightarrow \hat{Y}_i$ are the canonical injections and if*

$$e : X \rightarrow \prod_{i \in I} \hat{Y}_i : x \mapsto (j_i \circ f_i(x))_{i \in I}, \quad (4.7)$$

then $\hat{X} \cong \overline{e(X)}$.

PROOF. The (categorical) proof (e.g., [9]) of the equivalent statements in **Unif**, can be imitated. □

PROPOSITION 4.4. *If Y is a subspace of X in **AUnif**, then $\hat{Y} = \text{cl}_{\hat{X}}(Y)$.*

PROOF. Apply Proposition 4.3(2) to the embedding $i : Y \rightarrow X$. □

PROPOSITION 4.5. $\widehat{\prod_{i \in I} Y_i} = \prod_{i \in I} \widehat{Y_i}$.

PROOF. Since the source of projections $(\text{pr}_i : \prod Y_i \rightarrow Y_i)_{i \in I}$ is initial, we have by Proposition 4.3(2), that (closures with respect to $\prod \widehat{Y_i}$)

$$\widehat{\prod Y_i} = e(\overline{\prod Y_i}) = \overline{\prod Y_i} = \prod \widehat{Y_i}. \tag{4.8}$$

4.4. Hyperspaces. Let (X, d) be a metric space. In [6] it was shown that the hyperspace $\text{CL}(X) := \{A \subset X \mid A \text{ closed}\}$ allows a natural approach uniformity generated by the ∞p -metrics $\{d_H \mid H \in 2^{(X)}\}$, where $\forall A, B \in \text{CL}(X)$:

$$d_H(A, B) := \sup_{x \in H} |d(x, A) - d(x, B)|. \tag{4.9}$$

Its uniform coreflection is the Wijsman uniformity (see [1]). We shall show that $\text{CL}(\widehat{X}) = \widehat{\text{CL}(X)}$.

LEMMA 4.6. *Let X be a metric space. Then $\text{CL}(X)$ is a dense subspace of $\text{CL}(\widehat{X})$ in **AUnif**.*

PROOF. In [7, Theorem 1.5], it is shown that $i : \text{CL}(X) \rightarrow \text{CL}(\widehat{X}) : A \mapsto \text{cl}_{\widehat{X}} A$ is a dense embedding in **Unif**. We only need to show that $\text{CL}(X)$ carries the **AUnif**-subspace structure with respect to this embedding. Let d denote the metric on X , and let \hat{d} be the metric completion.

The initial approach uniformity for the singleton source i is generated by ∞p -metrics of the form

$$d_K \circ (i \times i)(A, B) = \sup_{x \in K} |\hat{d}(x, \text{cl}_{\widehat{X}}(A)) - \hat{d}(x, \text{cl}_{\widehat{X}}(B))|, \tag{4.10}$$

where K is a finite subset of \widehat{X} . If K is a finite subset of \widehat{X} and $\varepsilon > 0$, then there exists a finite subset $H \subset X$ such that $\forall x \in K, \exists y \in H : \hat{d}(x, y) < \varepsilon/2$. Then $d_K \circ (i \times i) \leq d_H + \varepsilon$. On the other hand, $\forall K \in 2^{(X)} : d_K \leq d_{i(K)} \circ (i \times i)$. \square

PROPOSITION 4.7. *Let X be a metric space. Then $\text{CL}(\widehat{X}) = \widehat{\text{CL}(X)}$.*

PROOF. First notice that if Y is complete, then its Wijsman uniformity (which is the uniform coreflection of $\text{CL}(Y)$) is complete by [7, Theorem 1.5]. Hence $\text{CL}(Y)$ is complete in **AUnif**. Since $\text{CL}(X)$ is a dense subspace of $\text{CL}(\widehat{X})$ (Lemma 4.6), we obtain by Proposition 4.4 that $\widehat{\text{CL}(X)}$ is the closure of $\text{CL}(X)$ in $\widehat{\text{CL}(\widehat{X})} = \text{CL}(\widehat{X})$, i.e., $\widehat{\text{CL}(X)} = \text{CL}(\widehat{X})$. \square

4.5. Compactification. The Čech-Stone compactification of a topological space can be described in two essentially different ways: as a closed subspace of a product of the unit interval $[0, 1]$ and as the underlying topology of the completion of the initial uniformity for all continuous functions into $[0, 1]$ (cf. [2]).

On the analogy of the former description a compactification in **AP** was defined in [4] (or [6]) as follows. (Suppose $[0, 1]$ is equipped with the usual, i.e., euclidean, approach structure.) If (X, δ) is an approach space, then denote

$$\mathfrak{K}^*(X) := \{f : X \rightarrow [0, 1] \mid f \text{ is a contraction}\}. \tag{4.11}$$

Suppose $X' := \prod_{f \in \mathfrak{K}^*(X)} [0, 1]$ is equipped with the product distance δ' . Consider $e : (X, \delta) \rightarrow (X', \delta') : x \mapsto (f(x))_{f \in \mathfrak{K}^*(X)}$ and let $\beta^* X := \overline{e(X)}$ be equipped with the subspace distance $\beta^* \delta$. The space $(\beta^* X, \beta^* \delta)$ is called the *compactification* of (X, δ) . This construction generalizes the Čech-Stone compactification in **Top**.

The alternative description mentioned above can be generalized in the context **AUnif**. (Henceforth, suppose $[0, 1]$ is equipped with the usual, i.e., euclidean, approach uniformity.)

PROPOSITION 4.8. *Let (X, δ) be a Hausdorff uniform approach space. If $K^*(X)$ is the initial approach uniformity for the source*

$$(f : X \rightarrow [0, 1])_{f \in \mathfrak{K}^*(X)}, \quad (4.12)$$

and $\hat{\delta}$ is the underlying approach space of $\widehat{K^*(X)}$, then $(\hat{X}, \hat{\delta}) = (\beta^* X, \beta^* \delta)$.

PROOF. By definition, the source

$$(f : (X, K^*(X)) \rightarrow [0, 1])_{f \in \mathfrak{K}^*(X)}. \quad (4.13)$$

is initial. By Proposition 4.3(2) we obtain that $(\hat{X}, \widehat{K^*(X)}) \cong (\overline{e(X)}, \Gamma)$ where Γ is the relativization of product approach uniformity on $[0, 1]^{\mathfrak{K}^*(X)}$. Consequently (and using [6, Proposition 4.1]),

$$(\hat{X}, \hat{\delta}) = (\hat{X}, A(\widehat{K^*(X)})) \cong (\overline{e(X)}, A(\Gamma)) = (\beta^* X, \beta^* \delta). \quad (4.14)$$

□

Still on the analogy of the situation in **Top** and **Unif**, we have the following corollary.

PROPOSITION 4.9. *Let (X, δ) be a uniform approach space. Then (X, δ) is compact if and only if $(X, K^*(X))$ is complete.*

4.6. Realcompactification. If in the above definition of compactification in **AP** $[0, 1]$ is replaced by \mathbb{R} (equipped with the usual approach structure) and the stars are dropped (in particular $\mathfrak{K}(X) := \{f : X \rightarrow \mathbb{R} \mid f \text{ is a contraction}\}$), then the space $(\beta X, \beta \delta)$ is a generalization of the well-known Hewitt-Nachbin realcompactification in **Top** (cf. [2]).

Here too, we have an alternative description in terms of completion.

PROPOSITION 4.10. *let (X, δ) be a Hausdorff uniform approach space. If $K(X)$ is the initial approach uniformity for the source*

$$(f : X \rightarrow \mathbb{R})_{f \in \mathfrak{K}(X)}, \quad (4.15)$$

and $\hat{\delta}$ is the underlying approach space of $\widehat{K(X)}$, then $(\hat{X}, \hat{\delta}) = (\beta X, \beta \delta)$.

COROLLARY 4.11. *Let (X, δ) be a uniform approach space. Then (X, δ) is realcompact if and only if $(X, K(X))$ is complete.*

COROLLARY 4.12. *Every realcompact uniform approach space is completely uniformizable. Every realcompact uniform approach space is AP-complete.*

PROOF. If (X, δ) is realcompact, then $K(X)$ is complete and induces δ . By Proposition 2.3 we see that (X, δ) is AP-complete. \square

ACKNOWLEDGEMENT. The second author is Aspirant of the Foundation for Scientific Research Flanders.

REFERENCES

- [1] G. Beer and R. Lucchetti, *Weak topologies for the closed subsets of a metrizable space*, Trans. Amer. Math. Soc. **335** (1993), no. 2, 805–822. MR 93d:54023. Zbl 810.54011.
- [2] L. Gillman and M. Jerison, *Rings of Continuous Functions*, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London-New York, 1960. MR 22#6994. Zbl 093.30001.
- [3] R. Lowen, *Approach Spaces*, The Clarendon Press Oxford University Press, New York, 1997. MR 98f:54002. Zbl 891.54001.
- [4] R. Lowen and K. Robeys, *Compactifications of products of metric spaces and their relations to Čech-Stone and Smirnov compactifications*, Topology Appl. **55** (1994), no. 2, 163–183. MR 94m:54063. Zbl 802.54018.
- [5] R. Lowen and B. Windels, *On the quantification of uniform properties*, Comment. Math. Univ. Carolin. **38** (1997), no. 4, 749–759. MR 98m:54029.
- [6] ———, *AUnif: a common supercategory of pMET and Unif*, Internat. J. Math. Math. Sci. **21** (1998), no. 1, 1–18. MR 98j:54014. Zbl 890.54024.
- [7] K. Morita, *Completion of hyperspaces of compact subsets and topological completion of open-closed maps*, General Topology and Appl. **4** (1974), 217–233. MR 50#3175. Zbl 288.54009.
- [8] C. G. C. Pitts, *Introduction to Metric Spaces*, Oliver & Boyd [Longman Group Ltd.], Edinburgh, 1972. MR 57#4112. Zbl 269.54019.
- [9] G. Preuss, *Theory of Topological Structures*, D. Reidel Publishing Co., Dordrecht, 1988. MR 89m:54014. Zbl 649.54001.
- [10] S. Willard, *General Topology*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1970. MR 41#9173. Zbl 205.26601.

LOWEN: UNIVERSITEIT ANTWERPEN, DEPARTEMENT WISKUNDE-INFORMATICA, GROENENBORGERLAAN 171 B-2020, ANTWERPEN, BELGIUM
E-mail address: lowen@ruca.ua.ac.be

WINDELS: DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF ANTWERP, RUCA, GROENENBORGERLAAN 171, 2020, ANTWERPEN, BELGIUM
E-mail address: windels@ruca.ua.ac.be

Special Issue on Impact of Kirk's Results in the Development of Fixed Point Theory

Call for Papers

Kirk's fixed point theorem published in 1965 had a profound impact in the development of the fixed point theory over the last 40 years. Through the concept of generalized distance, Tarski's classical fixed point theorem may be seen as a variant of Kirk's fixed point theorem in discrete sets. This shows among other things the power of this theorem.

This special issue will focus on any type of applications of Kirk's fixed point theorem. It includes:

- Nonexpansive mappings in Banach and metric spaces
- Multivalued mappings in Banach and metric spaces
- Monotone mappings in ordered sets
- Multivalued mappings in ordered sets
- Applications to nonmetric spaces like modular function spaces
- Applications to logic programming and directed graphs

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/fpta/guidelines.html>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	October 1, 2009
First Round of Reviews	January 1, 2010
Publication Date	April 1, 2010

Lead Guest Editor

M. A. Khamsi, Department of Mathematical Science,
University of Texas at El Paso, El Paso, TX 79968, USA;
mohamed@utep.edu

Guest Editor

Tomas Domínguez-Benavides, Departamento de Análisis
Matemático, Universidad de Sevilla, 41080 Sevilla, Spain;
tomasd@us.es

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br