This item is the archived peer-reviewed author-version of:

Do glucocorticoids mediate the link between environmental conditions and telomere dynamics in wild vertebrates? A review

Reference:

Angelier Frederic, Costantini David, Blevin Pierre, Chastel Olivier.- Do glucocorticoids mediate the link between environmental conditions and telomere dynamics in wild vertebrates? A review
General and comparative endocrinology - ISSN 0016-6480 - San diego, Academic press inc elsevier science, 256(2018), p. 99-111
Full text (Publisher's DOI): <https://doi.org/10.1016/J.YGCEN.2017.07.007>
To cite this reference: <http://hdl.handle.net/10067/1484340151162165141>

Accepted Manuscript

Review

Do glucocorticoids mediate the link between environmental conditions and telomere dynamics in wild vertebrates? A review

Frédéric Angelier, David Costantini, Pierre Blévin, Olivier Chastel

PII: S0016-6480(17)30185-5
DOI: <http://dx.doi.org/10.1016/j.ygcen.2017.07.007>
Reference: YGCEN 12691

To appear in: *General and Comparative Endocrinology*

Received Date: 9 March 2017
Revised Date: 6 June 2017
Accepted Date: 6 July 2017

Please cite this article as: Angelier, F., Costantini, D., Blévin, P., Chastel, O., Do glucocorticoids mediate the link between environmental conditions and telomere dynamics in wild vertebrates? A review, *General and Comparative Endocrinology* (2017), doi: <http://dx.doi.org/10.1016/j.ygcen.2017.07.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Do glucocorticoids mediate the link between environmental conditions and telomere dynamics in wild vertebrates? A review

Frédéric Angelier^{1,*}, David Costantini^{2,3}, Pierre Blévin¹, Olivier Chastel¹

¹ Centre d'Etudes Biologiques de Chizé, CNRS-ULR, UMR 7372, Villiers en Bois, France

² Muséum National d'Histoire Naturelle, UMR 7221, Paris, France

³ Behavioural Ecology & Ecophysiology group, Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

Revised manuscript submitted to *General and Comparative Endocrinology*

ISAE 2016 special issue

* Author for correspondence

F. Angelier

angelier@cebc.cnrs.fr

+33 (0)5 49 09 96 19

Abstract

Following the discoveries of telomeres and of their implications in terms of health and ageing, there has been a growing interest into the study of telomere dynamics in wild vertebrates. Telomeres are repeated sequences of non-coding DNA located at the terminal ends of chromosomes and they play a major role in maintaining chromosome stability. Importantly, telomeres shorten over time and shorter telomeres seem to be related with lower survival in vertebrates. Because of this potential link with longevity, it is crucial to understand not only the ecological determinants of telomere dynamics but also the regulatory endocrine mechanisms that may mediate the effect of the environment on telomeres. In this paper, we review the relationships that link environmental conditions, glucocorticoids (GC, the main hormonal mediator of allostasis) and telomere length in vertebrates. First, we review current knowledge about the determinants of inter-individual variations in telomere length. We emphasize the potential strong impact of environmental stressors and predictable life-history events on telomere dynamics. Despite recent progress, we still lack crucial basic data to fully understand the costs of several life-history stages and biotic and abiotic factors on telomere length. Second, we review the link that exists between GCs, oxidative stress and telomere dynamics in vertebrates. Although circulating GC levels may be closely and functionally linked with telomere dynamics, data are still scarce and somewhat contradictory. Further laboratory and field studies are therefore needed not only to better assess the proximate link between GC levels and telomere dynamics, but also to ultimately understand to what extent GCs and telomere length could be informative to measure the fitness costs of specific life-history stages and environmental conditions. Finally, we highlight the importance of exploring the functional links that may exist between coping styles, the GC stress response, and telomere dynamics in a life-history framework. To conclude, we raise new hypotheses regarding the potential of the GC stress response to drive the trade-off between immediate survival and telomere protection.

Key-words: *corticosterone; cortisol; stress; HPA axis; oxidative stress; telomere; telomerase*

1. Introduction: The relevance of studying telomeres in an ecological and evolutionary context

In 2009, Elizabeth H. Blackburn, Carol W. Greider and Jack W. Szostack received the Nobel Prize in Physiology and Medicine for the discovery of how chromosomes are protected by telomeres. These discoveries have had an important impact on biomedicine because telomeres and telomere dysfunctions have been closely associated with cancer, ageing and hereditary disease syndromes (Blackburn et al. 2006). Telomeres are well-conserved repeated sequences of non-coding DNA (TTAGGG) located at the terminal ends of chromosomes (Blackburn & Gall 1978). They play a major role in maintaining chromosome stability during replication processes (Blackburn 2005). Although telomeres can be restored by the enzyme telomerase, a ribonucleoprotein that adds new sequences onto the ends of chromosomes at each DNA replication (Greider & Blackburn 1989), telomeres shorten over time and are therefore considered as a molecular marker of cellular ageing (Harley et al. 1990). Importantly, this shortening has been associated with the occurrence of diseases and with increased mortality in humans (Cawthon et al. 2003; Lansdorp 2009).

Following these discoveries, there has been a growing interest into the study of telomeres in wild animals (Fig. 1). In the early 2000's, Haussmann and his collaborators have initiated the study of telomere biology in an ecological and evolutionary context (Haussmann & Vleck 2002; Haussmann et al. 2003a; Vleck et al. 2003). By studying telomeres in wild vertebrates, they were the first to emphasize the relevance of this molecular proxy of ageing to explore life-history trade-offs and selection processes (Haussmann et al. 2003b; Haussmann et al. 2005; Monaghan & Haussmann 2006; Vleck et al. 2007; Haussmann & Mauck 2008). Telomere length was initially thought to be a reliable proxy of age (i.e., "mitotic clock", Haussmann & Vleck 2002), leading to the idea that it could be a useful tool to obtain detailed information on wild populations. Although the relationship between chronological age and telomere length appears more complex than initially thought (Monaghan 2010; Dunshea et al. 2012; Boonekamp et al. 2013), there is now strong evidence that telomere length and telomere dynamics are tightly linked to fitness (Haussmann & Marchetto 2010; Bauch et al. 2013; Monaghan 2010, 2014; Ouyang et al. 2016), and especially to longevity and survival in

captive and wild vertebrates (e.g. Haussmann et al. 2005; Pauliny et al. 2006; Bize et al. 2009; Salomons et al. 2009; Heidinger et al. 2012; Bauch et al. 2013; Barrett et al. 2013; Ashgar et al. 2015; Fairlie et al. 2016).

These findings have raised a huge interest in measuring telomere length and telomere dynamics in wild animals (Nussey et al. 2014). Specifically, several studies have aimed to understand the ecological and life-history determinants of telomere dynamics. Such studies have led to the idea that telomere dynamics are tightly linked to the occurrence of stressors (reviewed in Haussmann & Marchetto 2010; Monaghan 2014; Haussmann & Heidinger 2015; Bateson 2016; Fig. 1), and more generally to allostasis (*sensu* McEwen & Wingfield 2003, i.e. maintaining stability through change). As a consequence, there has recently been a growing interest in understanding the links that may exist between the endocrine regulation of allostasis and telomere dynamics in wild vertebrates (reviewed in Haussmann & Marchetto 2010; Monaghan 2014; Fig. 1). In that context, glucocorticoids (GCs) obviously deserve a specific attention because of their involvement in allostasis and the stress response (McEwen & Wingfield 2003; Wingfield 2003, 2013; Romero et al. 2009; Angelier & Wingfield 2013; Wingfield et al. 2015; Romero & Wingfield 2016). Because of the link between telomeres and survival, integrating telomere dynamics into the framework linking environmental conditions, GCs and life-history decisions may contribute to a better understanding of the life-history/physiology nexus (Zera & Harshman 2001; Ricklefs & Wikelski 2002).

In this review, our aim is to emphasize the links that exist among telomere dynamics, environmental conditions and endocrine stress mechanisms in wild vertebrates. We review current knowledge about (1) the determinants of inter-individual variations in telomere length by emphasizing the role of environmental conditions and life-history stages on telomere dynamics; (2) the link that exists between GCs (i.e. the main hormonal mediator of allostasis), oxidative stress and telomere dynamics in wild vertebrates. Finally, we highlight the importance of exploring the

functional link that may exist between the GC stress response and telomere dynamics in a life-history framework.

2. Determinants of inter-individual variation in telomere length

2.1. A large inter-individual variability in telomere dynamics through life

In vertebrates, individual telomere length seems to be determined mainly through three steps: pre-natal telomere dynamics, developmental telomere dynamics, and adult telomere dynamics (Fig. 2). Firstly, telomere length is heritable and also affected by environmental, maternal and epigenetic effects (h^2 varying from 0.18 to 1.23 depending on the species, reviewed in Reichert et al. 2015), meaning that all individuals do not start their life with similar telomere length (Shalev et al. 2013). Thus, biomedical studies have first explored the potential genetic and environmental mechanisms affecting telomere length determination (Njajou et al. 2007; Broer et al. 2013) and, for example, have reported a strong positive influence of paternal age on offspring telomere length in humans (Unryn et al. 2005; Broer et al. 2013). More recently, ecological studies have confirmed that offspring telomere length is related to parents' telomere length through heritability, parental effect (Olsson et al. 2011; Reichert et al. 2015; Ashgar et al. 2015; Atema et al. 2015; Becker et al. 2015), and even trans-generational effects in wild vertebrates (Hausmann & Heidinger 2015; Bebbington et al. 2016). Secondly, apart from the first steps of the development when the telomeres of the oocytes and early embryo elongate (Liu et al. 2007), growth and development are classically associated with strong telomere attrition in humans (Frenck et al. 1998; Zeichner et al. 1999). Accordingly, most longitudinal studies have found that telomere shortens through growth and development in wild vertebrates (e.g. Pauliny et al. 2006, 2012; Salomons et al. 2009; Boonekamp et al. 2014; but see Ujvari & Madsen 2009; Ujvari et al. 2016), confirming the idea that development and growth are associated with accelerated cellular ageing in birds (Ricklefs 2010). Thirdly, after the developmental phase, this shortening seems to slow down as an individual ages and it is even difficult to detect in some species during adulthood (e.g. Foote et al. 2010; Pauliny et al. 2012; Rattiste et al. 2015).

Interestingly, longitudinal studies have clearly shown that telomere dynamics vary between individuals during development and adulthood (e.g. Bize et al. 2009; Foote et al. 2011; Salomons et al. 2009; Barrett et al. 2013; Parolini et al. 2015; Bateson et al. 2015; Stier et al. 2016; Fig. 2). The telomeres of some individuals shorten quickly while the telomeres of other individuals remain steady or even elongate with time (e.g. Bize et al. 2009; Parolini et al. 2015). This large inter-individual variation in telomere dynamics is the main reason why telomere length is not really a reliable predictor of chronological age in wild vertebrates (Dunshea et al. 2012; Pauliny et al. 2012; Boonekamp et al. 2013). However, the strong link between telomere length and remaining lifespan is promising because this suggests not only that telomere length could be a molecular marker of biological age or/and “wear and tear”, but also that telomere dynamics and telomere length could help us assess the influence of life-history events and/or environmental conditions on wild vertebrates.

2.2. Oxidative stress as a mediator of telomere dynamics

At the proximate level, oxidative stress is thought to be the primary cause of telomere shortening in wild vertebrates (reviewed in Haussmann & Marchetto 2010). Oxidative stress is a complex biochemical condition of the organism that is dependent on the rate of oxidative damage generation and oxidation of non-protein and protein thiols that regulate the cell oxidative balance (Jones 2006; Halliwell & Gutteridge 2007; Costantini & Verhulst 2009; Sohal & Orr 2012). Oxidative damage is caused by reactive oxygen species (ROS), such as free radicals, which are mainly produced by metabolic processes and immune cells (Halliwell & Gutteridge 2015). ROS can oxidize biomolecules, such as lipids, proteins, and DNA, and importantly, telomeres are especially sensitive to ROS because guanine is a dominant site for oxidatively-generated damage (Kawanishi & Oikawa 2004; Monaghan 2014). Supporting the importance of oxidative stress in determining telomere dynamics, a few studies have found an association between telomere dynamics and oxidative stress/antioxidant defenses in wild and domestic vertebrates (e.g. Geiger et al. 2012; Haussmann et al. 2012; Stier et al.

2014; Badás et al. 2015; Kim & Velando 2015; but see Ouyang et al. 2016 and Boonekamp et al. 2017). For example, Geiger et al. (2012) found that small king penguin chicks (*Aptenodytes patagonicus*) that died early during the growth period had the highest level of oxidative damage and the shortest telomere lengths prior to death. Badás et al. (2015) found that supplementation with antioxidants (which protect against oxidative stress) reduced telomere loss a year following treatment in wild blue tits (*Cyanistes caeruleus*). Mechanistic studies have provided stronger evidence for a link between oxidative stress and telomere shortening (reviewed in von Zglinicki 2002; but see Boonekamp et al. 2017), providing support for a possible causal link of results obtained in wild animals. For example, in cultures of human umbilical vein endothelial cells exposed to oxidative stress, terminal restriction fragment analysis demonstrated faster telomere shortening than in cells not exposed to oxidative stress (Kurz et al. 2004). In addition, telomere dynamics is also certainly dependent on DNA repair mechanisms that include the enzyme telomerase. Here again, most evidence comes from mechanistic studies, which have shown that this enzyme is the main actor of telomere elongation and restoration (Blackburn et al. 2006). Interestingly, oxidative stress induced a rapid and sustained decrease in the activity of this enzyme (Kurz et al. 2004). Moreover, Kurz et al. (2004) also demonstrated a key role for glutathione-dependent redox homeostasis (e.g., glutathione peroxidase) in the preservation of telomere function. Further support for a role of peroxidase activity in the protection of telomeres from oxidation was found in human embryonic kidney 293 (HEK293) cell lines (Aeby et al. 2016). All these mechanisms have rarely been studied in wild vertebrates (but see Haussmann et al. 2007). The connection between oxidative stress and telomere dynamics may also be mediated by glucocorticoids, suggesting one probable route through which environmental stress impacts on cellular ageing. For example, glucocorticoids may affect the oxidative balance through either genomic (Atasanova et al. 2009) or non-genomic (reviewed in Costantini et al. 2011) mechanisms. For example, exposure to dexamethasone enhances the expression of antioxidant genes under some circumstances (Atasanova et al. 2009). Similarly, glucocorticoids can suppress

cellular antioxidant defences though genomic effects in the rat (Kratschmar et al. 2012). Moreover, they may also affect telomere length through a modulation of telomerase activity (Figure 3).

3. Environmental conditions and telomere shortening

3.1. *Telomere dynamics vary between habitats, sites and years*

Biomedical studies have first suggested that lifestyle could affect telomere dynamics in humans. For example, obesity and cigarette smoking have been associated with shorter telomeres in humans (Valdes et al. 2005). More recently, ecological studies have explored the links that exist between environmental conditions and telomere dynamics in wild vertebrates (Table 1). A few studies have specifically compared telomere length and telomere dynamics of wild birds living in contrasting habitats. Overall, they found that an *a priori* less suitable habitat is associated with faster telomere attrition, and consequently, shorter telomeres (Angelier et al. 2013; Young et al. 2013; Salmon et al. 2016). Similarly, other studies reported that telomere dynamics or telomere length vary between years or sites that were characterized by contrasting environmental conditions (Watson et al. 2015; Becker et al. 2015; Quirici et al. 2016; Gangoso et al. 2016; Kirby et al. 2017). Moreover, Young et al. (2015) also nicely demonstrated in thick-billed murres that telomere length is linked with spatial habitat use, foraging efficiency and prey selection. Altogether, these studies emphasize that the environment has certainly a strong influence on telomere dynamics. However, all these studies compared some habitats or periods that drastically differed in terms of environmental conditions, and therefore, several biotic and abiotic variables could be responsible for these differences between habitats and periods. We need experimental studies to tease apart the effects of biotic and abiotic variables on telomeres.

3.2. *Ecological determinants of telomere dynamics*

In that context, other correlative and experimental studies have shed some light on the impact of specific environmental variables on telomere length. Infection and diseases may be associated with

reduced telomere length in wild animals (Table 2). For example, Ilmonen et al. (2008) found that repeated exposure to bacteria was associated with rapid telomere attrition in wild-derived mice (*Mus musculus*). More recently, Ashgar et al. (2015) found that chronic malaria infection was associated with significantly faster telomere attrition in a wild bird species, the great reed warbler (*Acrocephalus arundinaceus*). However, Sebastiano et al. (2017) did not find any association between telomere length and herpes viral disease in nestling Magnificent frigatebirds (*Fregata magnificens*).

Although psychological stress and associated disorders have been correlated with reduced telomere length in humans (Epel et al. 2004; Shalev et al. 2013), the influence of social variables on telomere dynamics has been overlooked in wild vertebrates and we currently lack data on the influence of social ranks and social bonds on telomere dynamics (Lewin et al. 2015). To our knowledge, a single study has tested the influence of psychological stress on telomere dynamics in birds and it found that social isolation was associated with fast telomere attrition in African grey parrots (*Psittacus erithacus erithacus*), a highly sociable species (Aydinonat et al. 2014). In addition to pathogens and social factors, nutritional deficit could also represent a major constraint for wild animals. Several studies have shown that nutritional constraints during the developmental phase are associated with short telomeres in birds (see Table 2). However, the influence of nutritional constraints (food abundance and food quality) on adult telomere dynamics has rarely been tested in adults, especially in wild vertebrates (Table 2). Recently, Hoelzl et al. (2016) showed that food supplementation reduces telomere attrition and is even associated with telomere elongation in a wild mammal species, the dormouse (*Glis glis*).

Human-induced disturbance may also cause a reduction of telomere length in wild vertebrates. For instance, urban noise has been experimentally shown to reduce telomere length in house sparrow chicks (*Passer domesticus*, Meillère et al. 2015). Similarly, exposure to handling stress was associated with telomere shortening in European shag nestlings (*Phalacrocorax aristotelis*, Herborn et al. 2014). However, there is currently no direct or experimental data on the influence of multiple

stressors on telomere dynamics that are known to occur frequently in wild vertebrate populations (e.g. predation risk). Work done on captive animals suggests that exposure to multiple stressors may influence telomere dynamics. Hau et al. (2015) exposed hand-raised adult Eurasian blackbirds (*Turdus merula*) to a combination of repeated immune and disturbance stressors for over one year to determine the effects of chronic stress on telomeres. By the end of the experiment, stress-exposed birds showed greater decreases in telomere lengths as compared to controls, suggesting that repeated exposure to experimental stressors might affect the rate of biological ageing (Hau et al. 2015).

Although a few studies have examined the impact of rising ambient temperature on telomere dynamics in fish (Rollings et al. 2014; Debes et al. 2016; Simide et al. 2016), there is currently little data on the effect of climatic variables on telomere dynamics or telomere length in wild birds or mammals. In black-tailed gulls (*Larus crassirostris*), mild weather and low sea-surface temperature were associated with a slow telomere shortening (Mizutani et al. 2013). This suggests that climate could affect telomere dynamics but additional studies are clearly needed. Finally, a few studies have reported that environmental contaminants can potentially have detrimental effects on telomere dynamics in free-living birds (Sletten et al. 2016; Stauffer et al. 2017). For instance, Blévin et al. (2016) recently found that oxychlorane contamination, an organochlorine pesticide known to be very toxic for wildlife, is associated with shorter telomeres in female black-legged kittiwakes (*Rissa tridactyla*). Our current world is facing multiple sources of contamination and their effects on physiology and fitness are often dose-dependent. Therefore, further experimental and correlative studies linking telomere dynamics and contamination are necessary to better understand the impact of anthropogenic pollution on wild vertebrates.

As detailed in the previous paragraphs, many studies have found that telomere dynamics and telomere length are tightly connected to a broad panel of environmental conditions that are experienced by individuals. Importantly, this connection is apparent both during development and

adulthood, demonstrating that environmental conditions are probably one of the main drivers of telomere dynamics in wild vertebrates. Further studies are however necessary to better understand not only the link between specific overlooked environmental conditions and telomere dynamics but also the potential additive, interactive, and antagonistic effects of the occurrence of multiple environmental constraints on telomere dynamics. Overall, all these results demonstrate that studying telomere dynamics or telomere length can be appropriate to assess the survival costs of specific environmental conditions in wild animals.

4. Telomere shortening through the life cycle

In addition to environmental conditions, predictable life-history events are also likely to have a strong impact on telomere dynamics, especially when they are associated with physiological and metabolic modifications (Monaghan 2014). During its life, an organism will face obligatory stages such as development and growth in early life and breeding, wintering, migration, or molt during adulthood (Wingfield 2008). Although necessary for reproduction and survival, all these stages can be associated with specific physiological constraints that may affect telomere dynamics.

There is now strong evidence that telomeres shorten very quickly during early life in most wild vertebrates (e.g. Salomons et al. 2009; Foote et al. 2011; Geiger et al. 2012; Heidinger et al. 2012; Reichert et al. 2015; Hammers et al. 2015; Ashgar et al. 2015). The pace of early life shortening may differ between males and females (Barrett & Richardson 2011; Parolini et al. 2015) and seem to be dependent on developmental conditions (Table 2). For instance, the rate of early-life telomere shortening is overall faster when sibling competition is harsher (e.g. Boonekamp et al. 2014; Nettle et al. 2015; Stier et al. 2015; Costanzo et al. 2017, table 2). Because replication is known to accelerate telomere attrition, such shortening certainly results from the replication processes that are inherent to growth and development of the organism. Moreover, growth is also associated with strong metabolic modification and with the production of oxidative stress (Smith et al. 2016) that is known to accelerate telomere attrition (von Zgliniki et al. 2002; Kawanishi & Oikawa 2004). In comparison,

telomere attrition is much slower during adulthood (e.g. Rattiste et al. 2015), but can still reflect exposure to chronic stress (Hau et al. 2015). Because offspring telomere length could be related to survival probability and lifespan (Pauliny et al. 2006; Heidinger et al. 2012; Barrett et al. 2013; Fairlie et al. 2016), this developmental telomere attrition supports the existence of a trade-off between growth and longevity (Ricklefs 2006; Lee et al. 2013). Therefore, comparing developmental telomere dynamics among species could help to understand the importance of this trade-off in mediating inter-specific life-history strategies (e.g. fast vs. slow development). Moreover, at the individual level, developmental telomere dynamics could help better assessing the influence of early developmental events on the future performances and the fitness of individuals (Monaghan 2014; Boonekamp et al. 2014).

Several studies have focused on the influence of reproductive effort on telomere dynamics and they have reported contrasting results (Table 2). Although some of them found that the reproductive phase, and especially the parental phase, is correlated with quick telomere shortening (Bauch et al. 2013; Sudyka et al. 2014; Reichert et al. 2014), others did not find such an effect (Beaulieu et al. 2010; Sudyka et al. 2016). Interestingly, some of these studies have experimentally shown that parental effort is associated with quick telomere attrition (Sudyka et al. 2014; Reichert et al. 2014). This supports the idea that reproduction is a demanding life-history stage for wild vertebrates and it illustrates perfectly the trade-off that exists between reproduction and survival (or at least organism's "wear and tear"). Surprisingly, the influence of other life-history stages on telomere dynamics has been much less studied in wild vertebrates. For example, the influence of molt on telomere dynamics has to our knowledge never been studied. The influence of migration on telomere dynamics has been studied in a few wild bird species so far. By comparing residents and migratory dark-eyed juncos (*Junco hyemalis*), Bauer et al. (2016) found that migration was associated with shorter telomeres, emphasizing a potential important cost of migration at least in this passerine species. Regarding wintering, studies are also scarce. Quick telomere attrition has been associated with a short time spent on the wintering ground in black-legged kittiwakes (Schultner et al. 2014) and

with an increased wintering foraging activity in thick-billed murres (Young et al. 2017). This suggests that this non-breeding stage may actually be associated with telomere preservation, at least when foraging conditions are appropriate. Such telomere preservation probably occurs because wintering individuals may face low energetic demands and may allocate a large part of their resources to protecting processes that limit aging and telomere shortening. Interestingly and supporting this interpretation, hibernation has also been studied in wild mammals and torpor seems to slow down telomere attrition in dormice (Hoelzl et al. 2016) and even to be associated with telomere elongation in sub-adult djungarian hamsters (*Phodopus sungorus*, Turbill et al. 2012, 2013). Overall, we currently lack crucial basic data to fully understand the costs of several life-history stages on telomere length, and thus organism's longevity. Moreover, to our knowledge, no study has examined the dynamics of telomeres through a whole annual life cycle in wild animals. Such an approach could be especially relevant to assess the relative costs of each life stage (reproduction, migration, wintering, molt, etc.). Finally, studying the impact of specific life-history stages on the telomere dynamics of individuals living in contrasting environments could shed some light on the costs and benefits of specific life-history decisions under contrasting environmental conditions.

5. Glucocorticoids and telomere dynamics

5.1. Are glucocorticoids the link between environmental constraints and telomere dynamics?

In the previous paragraphs, we reviewed the important links that exist between life-history demands, environmental conditions and telomere dynamics in wild vertebrates. Overall, demanding life-history stages and constraining environmental conditions seem to be associated with a rapid rate of telomere attrition and there is also a clear connection between stress and telomere attrition in humans, rodents and wild vertebrates (Epel et al. 2004; Kotrschal et al. 2007; Haussmann & Marchetto 2010; Shalev et al. 2013; Monaghan 2014; Hau et al. 2015; Bateson 2016). This suggests that telomere dynamics could be tightly related to allostatic load (*sensu* McEwen & Wingfield 2003; Romero et al. 2009) in wild vertebrates. When facing predictable and unpredictable events,

organisms develop a suite of behavioral and physiological adjustments to maintain their homeostasis (Wingfield 2003; Angelier & Wingfield 2013; Romero & Wingfield 2016). These adjustments are under control of a few central neurological and endocrine mechanisms (Romero et al. 2009). Among them, the Hypothalamus-Pituitary-Adrenal (HPA) axis appears especially relevant not only because it regulates several physiological functions but also because it is considered as a proxy of allostatic load in wild vertebrates (Romero et al. 2009). Supporting the relevance of this HPA axis when focusing on telomere dynamics, the activation of this axis and the resulting secretion of GCs are functionally linked with the proximate mechanisms regulating telomere dynamics (Fig. 3). Short-term exposure to increased GCs may increase oxidation but also antioxidant protection to limit spread of oxidative damage (reviewed in Costantini et al. 2011). On the other hand, prolonged exposure to high GCs results in increased oxidative stress and reduction of antioxidant defenses in vertebrates (reviewed in Costantini et al. 2011; see also work on wild marmots *Marmota marmota* by Costantini et al. 2012; and work on wild brown trouts *Salmo trutta* by Birnie-Gauvin et al. 2017). Importantly, biomedical studies have also shown that glucocorticoids may affect telomerase activity. Specifically, exposure to elevated cortisol levels is associated with a down-regulation of telomerase activity (Choi et al. 2008) and glutathione peroxidase (Patel et al. 2002), but mild increases in cortisol levels rather seems to up-regulate telomerase activity (Epel et al. 2010).

In addition, GCs are involved in several physiological systems that are also linked to oxidative stress or/and telomere shortening. For instance, GCs are involved in metabolic processes (reviewed in Landys et al. 2006) and an activation of these processes results in an increased production of ROS (Costantini 2014). Specifically, increased circulating GCs levels are associated with protein catabolism and are thought to promote glucose and lipid mobilization (reviewed in Landys et al. 2006). Moreover, GCs are closely connected to immune activation (Martin 2009) that is known to be associated with the production of ROS and with telomere attrition (Ilmonen et al. 2008; Ashgar et al. 2015). Increased circulating GC levels are known to enhance parts of the innate and adaptive immune responses in the short-term while the pathological maintenance of elevated GC levels during

a prolonged period is rather associated with immune suppression (reviewed in Martin 2009). Similarly, circulating GC levels are associated to disturbance and the occurrence of acute and chronic stressors (Dickens & Romero 2013; Madliger & Love 2016) that also seem to accelerate telomere attrition (Epel et al. 2004; Kotrschal et al. 2007; Herborn et al. 2014; Hau et al. 2015). Furthermore, a few studies have also shown that environmental contaminants can both disrupt GC regulation (Nordstad et al. 2012; Tartu et al. 2015; Meillère et al. 2016), and affect telomere dynamics in wild vertebrates (Blévin et al. 2016; Stauffer et al. 2017). Finally, circulating GC levels are known to fluctuate during the life-history cycle (reviewed in Romero 2002), and interestingly, telomere shortening also seems greater during stages when GC levels are elevated. For example, GC levels are elevated during reproduction (Romero 2002), which is associated with fast telomere attrition (Bauch et al. 2013; Sudyka et al. 2014; Reichert et al. 2014). Moreover, reproductive effort is positively associated with both elevated GCs (e.g. large brood size, Bonier et al. 2011; Love et al. 2014) and rapid telomere attrition (Sudyka et al. 2014; Reichert et al. 2014), emphasizing this potential link between telomere dynamics and GCs.

5.2. Telomere length and circulating glucocorticoids levels: a context-dependent link?

All these results strongly suggest that HPA regulation probably plays a major role in determining telomere dynamics in wild vertebrates. However, only a few studies have examined the link between circulating GCs levels and telomere dynamics in wild vertebrates (Table 3). In thorn-tailed rayaditos (*Aphrastura spinicauda*), telomere length was negatively correlated with circulating corticosterone levels. Similarly, birds with higher corticosterone levels had shorter telomeres in black-browed albatrosses (*Thalassarche melanophrys*, Angelier et al. unpublished data) and in Andean condors (Gangoso et al. 2016). Interestingly, Young et al. (2016) found that the link between telomere length and corticosterone levels was inconsistent in thick-billed murrelets (*Uria lomvia*). Indeed, corticosterone levels were positively correlated with telomere length in one colony but negatively correlated in another one. Other studies found no correlation between circulating GC levels and

telomere length (Young et al. 2016; Ouyang et al. 2016). Similarly, Bauch et al. (2016) found that corticosterone levels were negatively correlated with telomere length in male but not in female Common terns. All these studies suggest that the relationship between GCs and telomere length is complex and context dependent (Young et al. 2016; Bauch et al. 2016). This is not so surprising because GCs can be positively correlated to individual quality in some circumstances (the CORT-adaptation hypothesis, Bonier et al. 2009) and negatively in others (the CORT-fitness hypothesis, Bonier et al. 2009). Thus, elevated GC levels can be a proxy of the inability of an individual to cope with its environment, leading to the idea that elevated GC levels should be found in individuals of low quality (Angelier et al. 2010), and thus in individuals with short telomeres (Quirici et al. 2016; Angelier et al. unpublished data; Young et al. 2016; Gangoso et al. 2016). On the other hand, telomere length and GC levels could be positively correlated when GC levels are positively correlated with individual quality. For instance, elevated GC levels are necessary to sustain the energetic demands of reproduction, and therefore, elevated GC levels should be only found in individuals that are able to allocate resources to reproduction, i.e. individuals of high quality with long telomeres (Bauch et al. 2016; Young et al. 2016).

Importantly, a few experimental studies have also examined the link between GCs and telomere dynamics (Monaghan 2014). In developing offspring, telomere attrition was accelerated by handling stress but an additional experimental increase of circulating GCs levels did not amplify this pattern (Herborn et al. 2014). In domestic chickens (*Gallus gallus*), an embryonic exposure to GCs resulted in shorter telomeres 25 days after hatching (Hausmann et al. 2012). In adults, two experimental studies have to our knowledge examined the impact of corticosterone on telomere dynamics. In black-legged kittiwakes, a temporary increase in corticosterone levels was clearly associated with a faster rate of telomere attrition over a year (Schultner et al. 2014). In captive zebra finches (*Taeniopygia guttata*), injection of corticosterone induced a faster rate of attrition in reproductive females (Tissier et al. 2014). All these studies clearly demonstrate that there is a strong functional link between the HPA axis and telomere dynamics. However, most studies have focused on

circulating GC levels without exploring the whole and complex functioning of the HPA axis. The relevance of circulating GC levels as a proxy of physiological stress or fitness is currently debated (Bonier et al. 2009; Dickens & Romero 2013; Love et al. 2014) because the actions of GCs on physiology and behavior are regulated at multiple levels (e.g. receptors, corticosterone-binding globulins, negative feedback, etc. Romero 2004). Therefore, further laboratory and field studies are mandatory not only to better assess the proximate link between the HPA axis and telomere dynamics, but also to ultimately understand to what extent GCs and telomere length could be informative to measure the fitness costs of specific life-history stages and environmental conditions.

5.3. Coping styles, the glucocorticoid stress response and telomere dynamics

So far, most studies have focused on baseline circulating GC levels that provide information on the energetic status of individuals (i.e. allostatic load, McEwen & Wingfield 2003; Romero et al. 2009). However, the link between the physiological sensibility to stressors and telomere dynamics has been overlooked. In wild vertebrates, individuals adopt different strategies to cope with unpredictable events and these coping styles have been linked to the HPA axis (Wingfield 2003, 2013, 2015). In response to an unpredictable event, the HPA axis is activated and this results in the rapid and transitory acute secretion of GCs in the bloodstream (Romero 2004). These increased circulating GC levels act on specific behavioral and physiological systems to activate an emergency life-history stage (ELHS) that aims to promote immediate survival at the expense of other life-history components, such as reproduction (Wingfield et al. 1998). Importantly, there is a large inter-individual variation in this so called GC stress response (Cockrem 2013) and this variability is associated to different stress sensitivity and coping styles. Although this GC stress response is known to mediate life-history decisions (Wingfield & Sapolsky 2003; Lendvai et al. 2007; Krause et al. 2016) and to have important fitness consequences (Breuner et al. 2008), its link with telomere dynamics has to our knowledge never been examined. Here, we develop a theoretical framework and several hypotheses that may stimulate future studies.

Because elevated GC levels are linked with oxidative stress (Costantini et al. 2011) and reduced telomerase activity (Choi et al. 2008), mounting a GC stress response could be associated with increased DNA damage and telomere shortening. One hidden cost of an increased sensitivity to stress could therefore be an acceleration of telomere shortening, and more generally, ageing. Because telomere length is positively associated with longevity (Pauliny et al. 2006; Bize et al. 2009; Salomons et al. 2009; Heidinger et al. 2012; Barrett et al. 2013; Fairlie et al. 2016), the GC stress response may therefore mediate the trade-off between immediate survival and long-term survival (Hypothesis 1). Thus, a high sensitivity to stress (and a strong GC stress response) could promote immediate survival at the expense of telomere protection, and thus, longevity (Fig. 4A).

Importantly, this trade-off between immediate survival and telomere preservation may be exacerbated or alleviated depending on the environmental context and/or individual quality. Individual quality involves multiple phenotypic traits that are positively correlated with fitness (Wilson & Nussey 2010). These phenotypic traits certainly involve behavior, physiology and morphological components that can benefit to fitness under specific circumstances. Overall, we may expect individuals of higher quality to have a better access to resources and to be able to allocate more resources to multiple competing traits. If individuals of high quality or individuals living in a highly suitable environment have for instance better antioxidant defenses or better DNA repair mechanisms, they may be able to limit the oxidative damages that are associated with elevated GC levels. Therefore, they may bear less telomere shortening when mounting an intense stress response (Hypothesis 2). Under that scenario, the link between telomere dynamics and the intensity of the GC stress response would depend not only on the environmental context, but also on the state of the individual (Fig. 4B).

Finally, the situation could even be more complex when individuals are engaged in a specific life-history stage. The GC stress response mediates behavioral and physiological adjustments that shift the organism from specific stages to an ELHS, which prioritizes immediate survival at the expense of other demanding activities (Wingfield et al. 1998). When these demanding activities are associated

with oxidative stress and telomere shortening, the disruption of these activities may actually be associated with a slowing down of the rate of telomere attrition. Therefore, a high sensitivity to stress (and a strong GC stress response) could be counter-intuitively associated with reduced telomere attrition under some circumstances (Hypothesis 3, Fig. 4C). Given our current knowledge of the links between GCs, telomere shortening and reproductive effort, the best support for this hypothesis comes from studies that have focused on the reproductive stage. During reproduction, an intense GC stress response is classically associated with a substantial reduction of parental effort (Wingfield & Sapolsky 2003; Lendvai et al. 2007; Krause et al. 2016), which is known to translate into a reduction of the rate of telomere attrition (Bauch et al. 2013; Sudyka et al. 2014; Reichert et al. 2014). In contrast, a weak stress response is usually associated with the maintenance of parental effort (Wingfield & Sapolsky 2003; Lendvai et al. 2007; Krause et al. 2016), and thus with potential significant survival costs ("the cost of reproduction") that would be associated to an important telomere attrition (Bauch et al. 2013; Sudyka et al. 2014; Reichert et al. 2014).

The link between the GC stress response and telomere attrition is however probably not so straightforward and it certainly depends on the costs and benefits of activating an ELHS that are certainly species and context dependent. Specifically, these costs and benefits certainly depend on what aspects of behavior or physiology are up or down-regulated when an ELHS is activated. For instance, high stress sensitivity could be associated with a reduced rate of telomere shortening in breeding individuals if it limits the negative impact of breeding activities on telomere dynamics. However, high stress sensitivity could also be associated with a risk of mounting numerous and repeated GC stress responses that are known to accelerate telomere attrition. Under some circumstances, the negative impact of high stress sensitivity on telomere dynamics may therefore outweigh the benefits of a reduced breeding effort on telomere dynamics, especially when the costs of reproduction are limited. Therefore, it appears crucial to better assess the costs and benefits of elevated GC levels and specific life-history stages on telomere dynamics in multiple species with

contrasted life-history strategies and further studies are clearly needed to better understand the functional link between the GC stress response and telomere dynamics in wild vertebrates.

Although the GC stress response is known to be heritable and repeatable (Evans et al. 2006; Cockrem et al. 2009; Angelier et al. 2011; Jenkins et al. 2014), it is also flexible because it can be actively modulated by individuals (Wingfield & Sapolsky 2003; Lendvai et al. 2007; Krause et al. 2016). The ability of individuals to adaptively modulate this GC stress response is thought to be involved in their capacity to cope with a changing world (Angelier & Wingfield 2013; Wingfield et al. 2015). Exploring the link between the GC stress response, its flexibility, and telomere dynamics should further shed some light on the costs and benefits of the hormonal sensitivity to stress. Therefore, we believe that combining GCs and telomere dynamics measurements in several ecological contexts should open some exciting and promising new research areas in environmental endocrinology.

Acknowledgements

We are grateful to the ANR (URBASTRESS, ANR-16-CE02-0004-01), the Centre National de la Recherche Scientifique (CNRS), the University of La Rochelle (ULR), the Polar French Institute (IPEV, Prog. 109, 330), the Région Poitou-Charentes, and the Contrat Plan Etat region (CPER ECONAT) for funding this research.

References

- Aeby, E., Ahmed, W., Redon, S., Simanis, V., Lingner, J., 2016. Peroxiredoxin 1 protects telomeres from oxidative damage and preserves telomeric DNA for extension by telomerase. *Cell Reports* 17, 3107–3114.
- Angelier, F., Wingfield, J.C., Weimerskirch, H., Chastel, O., 2010. Hormonal correlates of individual quality in a long-lived bird: a test of the “corticosterone–fitness hypothesis.” *Biology Letters* 6, 846–849.

- Angelier, F., Ballentine, B., Holberton, R.L., Marra, P.P., Greenberg, R., 2011. What drives variations in the corticosterone stress response between subspecies? A common garden experiment of swamp sparrows (*Melospiza Georgiana*). *J. Evol. Biol.* 24, 1274–1283.
- Angelier, F., Wingfield, J.C., 2013. Importance of the glucocorticoid stress response in a changing world: theory, hypotheses and perspectives. *General and comparative endocrinology* 190, 118–128.
- Angelier, F., Vleck, C.M., Holberton, R.L., Marra, P.P., 2013. Telomere length, non-breeding habitat and return rate in male American redstarts. *Functional Ecology* 27, 342–350.
- Angelier, F., Vleck, C.M., Holberton, R.L., Marra, P.P., 2015. Bill size correlates with telomere length in male American redstarts. *Journal of Ornithology* 156, 525–531.
- Asghar, M., Bensch, S., Tarka, M., Hansson, B., Hasselquist, D., 2015. Maternal and genetic factors determine early life telomere length. *Proceedings of the Royal Society of London B: Biological Sciences* 282, 20142263.
- Asghar, M., Hasselquist, D., Hansson, B., Zehindjiev, P., Westerdahl, H., Bensch, S., 2015. Hidden costs of infection: chronic malaria accelerates telomere degradation and senescence in wild birds. *Science* 347, 436–438.
- Atanasova, S., Wieland, E., Schlumbohm, C., Korecka, M., Shaw, L., von Ahsen, N., Fuchs, E., Oellerich, M., Armstrong, V., 2009. Prenatal dexamethasone exposure in the common marmoset monkey enhances gene expression of antioxidant enzymes in the aorta of adult offspring. *Stress* 12, 215–224.
- Atema, E., Mulder, E., Dugdale, H.L., Briga, M., van Noordwijk, A.J., Verhulst, S., 2015. Heritability of telomere length in the Zebra Finch. *Journal of Ornithology* 156, 1113–1123.
- Aydinonat, D., Penn, D.J., Smith, S., Moodley, Y., Hoelzl, F., Knauer, F., Schwarzenberger, F., 2014. Social isolation shortens telomeres in African grey parrots (*Psittacus erithacus erithacus*). *PloS one* 9, e93839.
- Badás, E.P., Martínez, J., Rivero de Aguilar Cachafeiro, J., Miranda, F., Figuerola, J., Merino, S., 2015. Ageing and reproduction: antioxidant supplementation alleviates telomere loss in wild birds. *Journal of evolutionary biology* 28, 896–905.
- Barrett, E.L., Richardson, D.S., 2011. Sex differences in telomeres and lifespan. *Aging Cell* 10, 913–921.
- Barrett, E.L., Burke, T.A., Hammers, M., Komdeur, J., Richardson, D.S., 2013. Telomere length and dynamics predict mortality in a wild longitudinal study. *Molecular ecology* 22, 249–259.
- Bateson, M., Brilot, B.O., Gillespie, R., Monaghan, P., Nettle, D., 2015. Developmental telomere attrition predicts impulsive decision-making in adult starlings. *Proceedings of the Royal Society of London B: Biological Sciences* 282, 20142140.

- Bateson, M., 2016. Cumulative stress in research animals: Telomere attrition as a biomarker in a welfare context? *BioEssays* 38, 201–212.
- Bauch, C., Becker, P.H., Verhulst, S., 2013. Telomere length reflects phenotypic quality and costs of reproduction in a long-lived seabird. *Proceedings of the Royal Society of London B: Biological Sciences* 280, 20122540.
- Bauch, C., Riechert, J., Verhulst, S., Becker, P.H., 2016. Telomere length reflects reproductive effort indicated by corticosterone levels in a long-lived seabird. *Molecular Ecology* 25, 5785–5794.
- Bauer, C.M., Heidinger, B.J., Ketterson, E.D., Greives, T.J., 2016. A migratory lifestyle is associated with shorter telomeres in a songbird (*Junco hyemalis*). *The Auk* 133, 649–653.
- Bebbington, K., Spurgin, L.G., Fairfield, E.A., Dugdale, H.L., Komdeur, J., Burke, T., Richardson, D.S., 2016. Telomere length reveals cumulative individual and transgenerational inbreeding effects in a passerine bird. *Molecular Ecology* 25, 2949–2960.
- Becker, P.J., Reichert, S., Zahn, S., Hegelbach, J., Massemin, S., Keller, L.F., Postma, E., Criscuolo, F., 2015. Mother–offspring and nest-mate resemblance but no heritability in early-life telomere length in white-throated dippers. *Proceedings of the Royal Society of London B: Biological Sciences* 282, p. 20142924.
- Beirne, C., Delahay, R., Hares, M., Young, A., 2014. Age-related declines and disease-associated variation in immune cell telomere length in a wild mammal. *PloS one* 9, e108964.
- Birnie-Gauvin, K., Peiman, K.S., Larsen, M.H., Aarestrup, K., Willmore, W.G., Cooke, S.J., 2017. Short-term and long-term effects of transient exogenous cortisol manipulation on oxidative stress in juvenile brown trout. *Journal of Experimental Biology* *in press*.
- Bize, P., Criscuolo, F., Metcalfe, N.B., Nasir, L., Monaghan, P., 2009. Telomere dynamics rather than age predict life expectancy in the wild. *Proceedings of the Royal Society of London B: Biological Sciences* 276, 1679–1683.
- Blackburn, E.H., 2005. Telomeres and telomerase: their mechanisms of action and the effects of altering their functions. *FEBS letters* 579, 859–862.
- Blackburn, E.H., Gall, J.G., 1978. A tandemly repeated sequence at the termini of the extrachromosomal ribosomal RNA genes in *Tetrahymena*. *Journal of Molecular Biology* 120, 33–53.
- Blackburn, E.H., Greider, C.W., Szostak, J.W., 2006. Telomeres and telomerase: the path from maize, *Tetrahymena* and yeast to human cancer and aging. *Nature Medicine* 12, 1133–1138.
- Blévin, P., Angelier, F., Tartu, S., Ruault, S., Bustamante, P., Herzke, D., Moe, B., Bech, C., Gabrielsen, G.W., Bustnes, J.O., 2016. Exposure to oxychlordan is associated with shorter telomeres in arctic breeding kittiwakes. *Science of the Total Environment* 563, 125–130.

- Bonier, F., Martin, P.R., Moore, I.T., Wingfield, J.C., 2009. Do baseline glucocorticoids predict fitness? *Trends in Ecology & Evolution* 24, 634–642.
- Bonier, F., Moore, I.T., Robertson, R.J., 2011. The stress of parenthood? Increased glucocorticoids in birds with experimentally enlarged broods. *Biology Letters* 7, 944–946.
- Boonekamp, J.J., Bauch, C., Mulder, E., Verhulst, S., 2017. Does oxidative stress shorten telomeres? *Biology Letters* 20170164.
- Boonekamp, J.J., Mulder, G.A., Salomons, H.M., Dijkstra, C., Verhulst, S., 2014. Nestling telomere shortening, but not telomere length, reflects developmental stress and predicts survival in wild birds. *Proceedings of the Royal Society of London B: Biological Sciences* 281, 20133287.
- Boonekamp, J.J., Simons, M.J., Hemerik, L., Verhulst, S., 2013. Telomere length behaves as biomarker of somatic redundancy rather than biological age. *Aging Cell* 12, 330–332.
- Breuner, C.W., Patterson, S.H., Hahn, T.P., 2008. In search of relationships between the acute adrenocortical response and fitness. *General and comparative endocrinology* 157, 288–295.
- Broer, L., Codd, V., Nyholt, D.R., Deelen, J., Mangino, M., Willemsen, G., Albrecht, E., Amin, N., Beekman, M., De Geus, E.J., 2013. Meta-analysis of telomere length in 19 713 subjects reveals high heritability, stronger maternal inheritance and a paternal age effect. *European Journal of Human Genetics* 21, 1163–1168.
- Caprioli, M., Romano, M., Romano, A., Rubolini, D., Motta, R., Folini, M., Saino, N., 2013. Nestling telomere length does not predict longevity, but covaries with adult body size in wild barn swallows. *Biology Letters* 9, 20130340.
- Cawthon, R.M., Smith, K.R., O'Brien, E., Sivatchenko, A., Kerber, R.A., 2003. Association between telomere length in blood and mortality in people aged 60 years or older. *The Lancet* 361, 393–395.
- Choi, J., Fauce, S.R., Effros, R.B., 2008. Reduced telomerase activity in human T lymphocytes exposed to cortisol. *Brain, behavior, and immunity* 22, 600–605.
- Cockrem, J.F., 2013. Individual variation in glucocorticoid stress responses in animals. *General and Comparative Endocrinology* 181, 45–58.
- Cockrem, J.F., Barrett, D.P., Candy, E.J., Potter, M.A., 2009. Corticosterone responses in birds: individual variation and repeatability in Adelie penguins (*Pygoscelis adeliae*) and other species, and the use of power analysis to determine sample sizes. *General and comparative endocrinology* 163, 158–168.
- Colitz, C.M.H., Whittington, A., Carter, R., Warren, J., 2004. The effect of oxidative stress on telomerase activity and other stress-related proteins in lens epithelial cells. *Experimental Eye Research* 78, 235–242.

- Costantini, D., 2014. Oxidative stress and hormesis in evolutionary ecology and physiology. A marriage between mechanistic and evolutionary approaches. Springer.
- Costantini, D., Ferrari, C., Pasquaretta, C., Cavallone, E., Carere, C., von Hardenberg, A., Réale, D., 2012. Interplay between plasma oxidative status, cortisol and coping styles in wild alpine marmots, *Marmota marmota*. *Journal of Experimental Biology* 215, 374–383.
- Costantini, D., Marasco, V., Møller, A.P., 2011. A meta-analysis of glucocorticoids as modulators of oxidative stress in vertebrates. *Journal of Comparative Physiology B* 181, 447–456.
- Costantini, D., Verhulst, S., 2009. Does high antioxidant capacity indicate low oxidative stress? *Functional Ecology* 23, 506–509.
- Costanzo, A., Parolini, M., Bazzi, G., Khoraiuli, L., Santagostino, M., Possenti, C.D., Romano, A., Nergadze, S.G., Rubolini, D., Giolotto, E., 2016. Brood size, telomere length, and parent-offspring color signaling in barn swallows. *Behavioral Ecology in press*.
- Debes, P.V., Visse, M., Panda, B., Ilmonen, P., Vasemägi, A., 2016. Is telomere length a molecular marker of past thermal stress in wild fish? *Molecular Ecology* 25, 5412–5424.
- Dickens, M.J., Romero, L.M., 2013. A consensus endocrine profile for chronically stressed wild animals does not exist. *General and Comparative Endocrinology* 191, 177–189.
- Dunshea, G., Duffield, D., Gales, N., Hindell, M., Wells, R.S., Jarman, S.N., 2011. Telomeres as age markers in vertebrate molecular ecology. *Molecular Ecology Resources* 11, 225–235.
- Epel, E.S., Blackburn, E.H., Lin, J., Dhabhar, F.S., Adler, N.E., Morrow, J.D., Cawthon, R.M., 2004. Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of the United States of America* 101, 17312–17315.
- Epel, E.S., Lin, J., Dhabhar, F.S., Wolkowitz, O.M., Puterman, E., Karan, L., Blackburn, E.H., 2010. Dynamics of telomerase activity in response to acute psychological stress. *Brain, Behavior, and Immunity* 24, 531–539.
- Evans, M.R., Roberts, M.L., Buchanan, K.L., Goldsmith, A.R., 2006. Heritability of corticosterone response and changes in life history traits during selection in the zebra finch. *Journal of evolutionary biology* 19, 343–352.
- Fairlie, J., Holland, R., Pilkington, J.G., Pemberton, J.M., Harrington, L., Nussey, D.H., 2016. Lifelong leukocyte telomere dynamics and survival in a free-living mammal. *Aging Cell* 15, 140–148.
- Foote, C.G., Daunt, F., González-Solís, J., Nasir, L., Phillips, R.A., Monaghan, P., 2010. Individual state and survival prospects: age, sex, and telomere length in a long-lived seabird. *Behavioral Ecology* 22, 156–161.
- Foote, C.G., Gault, E.A., Nasir, L., Monaghan, P., 2011. Telomere dynamics in relation to early growth conditions in the wild in the lesser black-backed gull. *Journal of Zoology* 283, 203–209.

- Frenck, R.W., Blackburn, E.H., Shannon, K.M., 1998. The rate of telomere sequence loss in human leukocytes varies with age. *Proceedings of the National Academy of Sciences* 95, 5607–5610.
- Gangoso, L., Lambertucci, S.A., Cabezas, S., Alarcón, P.A., Wiemeyer, G.M., Sanchez-Zapata, J.A., Blanco, G., Hiraldo, F., Donázar, J.A., 2016. Sex-dependent spatial structure of telomere length in a wild long-lived scavenger. *Ecosphere* 7, e01544.
- Geiger, S., Le Vaillant, M., Lebard, T., Reichert, S., Stier, A., Le Maho, Y., Criscuolo, F., 2012. Catching-up but telomere loss: half-opening the black box of growth and ageing trade-off in wild king penguin chicks. *Molecular Ecology* 21, 1500–1510.
- Greider, C.W., Blackburn, E.H., 1989. A telomeric sequence in the RNA of *Tetrahymena* telomerase required for telomere repeat synthesis. *Nature* 337, 331–337.
- Halliwell, B., Gutteridge, J.M., 2015. *Free radicals in biology and medicine*. Oxford University Press, USA.
- Hammers, M., Kingma, S.A., Bebbington, K., van de Crommenacker, J., Spurgin, L.G., Richardson, D.S., Burke, T., Dugdale, H.L., Komdeur, J., 2015. Senescence in the wild: Insights from a long-term study on Seychelles warblers. *Experimental gerontology* 71, 69–79.
- Harley, C.B., Futcher, A.B., Greider, C.W., 1990. Telomeres shorten during ageing of human fibroblasts. *Nature* 345, 458.
- Hau, M., Haussmann, M.F., Greives, T.J., Matlack, C., Costantini, D., Quetting, M., Adelman, J.S., Miranda, A.C., Partecke, J., 2015. Repeated stressors in adulthood increase the rate of biological ageing. *Frontiers in zoology* 12, 4.
- Haussmann, M.F., Heidinger, B.J., 2015. Telomere dynamics may link stress exposure and ageing across generations. *Biology letters* 11, 20150396.
- Haussmann, M.F., Longenecker, A.S., Marchetto, N.M., Juliano, S.A., Bowden, R.M., 2012. Embryonic exposure to corticosterone modifies the juvenile stress response, oxidative stress and telomere length. *Proceedings of the Royal Society of London B: Biological Sciences*, 279, 1447–1456.
- Haussmann, M.F., Marchetto, N.M., 2010. Telomeres: linking stress and survival, ecology and evolution. *Current Zoology* 56, 714–27.
- Haussmann, M.F., Mauck, R.A., 2008. Telomeres and longevity: testing an evolutionary hypothesis. *Molecular Biology and Evolution* 25, 220–228.
- Haussmann, M.F., Vleck, C.M., 2002. Telomere length provides a new technique for aging animals. *Oecologia* 130, 325–328.
- Haussmann, M.F., Vleck, C.M., Nisbet, I.C., 2003a. Calibrating the telomere clock in common terns, *Sterna hirundo*. *Experimental gerontology* 38, 787–789.
- Haussmann, M.F., Winkler, D.W., Huntington, C.E., Nisbet, I.C., Vleck, C.M., 2007. Telomerase activity is maintained throughout the lifespan of long-lived birds. *Experimental gerontology* 42, 610–618.

- Hausmann, M.F., Winkler, D.W., O'Reilly, K.M., Huntington, C.E., Nisbet, I.C., Vleck, C.M., 2003b. Telomeres shorten more slowly in long-lived birds and mammals than in short-lived ones. *Proceedings of the Royal Society of London B: Biological Sciences* 270, 1387–1392.
- Hausmann, M.F., Winkler, D.W., Vleck, C.M., 2005. Longer telomeres associated with higher survival in birds. *Biology Letters* 1, 212–214.
- Heidinger, B.J., Blount, J.D., Boner, W., Griffiths, K., Metcalfe, N.B., Monaghan, P., 2012. Telomere length in early life predicts lifespan. *Proceedings of the National Academy of Sciences* 109, 1743–1748.
- Heidinger, B.J., Herborn, K.A., Granroth-Wilding, H., Boner, W., Burthe, S., Newell, M., Wanless, S., Daunt, F., Monaghan, P., 2016. Parental age influences offspring telomere loss. *Functional Ecology* 30, 1531–1538.
- Herborn, K.A., Heidinger, B.J., Boner, W., Noguera, J.C., Adam, A., Daunt, F., Monaghan, P., 2014. Stress exposure in early post-natal life reduces telomere length: an experimental demonstration in a long-lived seabird, *Proceedings of the Royal Society of London B: Biological Sciences* 281, 20133151.
- Hoelzl, F., Cornils, J.S., Smith, S., Moodley, Y., Ruf, T., 2016. Telomere dynamics in free-living edible dormice (*Glis glis*): the impact of hibernation and food supply. *Journal of Experimental Biology* 219, 2469–2474.
- Ilmonen, P., Kotrschal, A., Penn, D.J., 2008. Telomere attrition due to infection. *PloS one* 3, e2143.
- Jones, D.P., 2006. Redefining oxidative stress. *Antioxidants & Redox Signaling* 8, 1865–1879.
- Jenkins, B. R., Vitousek, M. N., Hubbard, J. K. & Safran, R. J. 2014 An experimental analysis of the heritability of variation in glucocorticoid concentrations in a wild avian population. *Proc. R. Soc. B* 281, 20141302.
- Kawanishi, S., Oikawa, S., 2004. Mechanism of telomere shortening by oxidative stress. *Annals of the New York Academy of Sciences* 1019, 278–284.
- Kim, S.-Y., Velando, A., 2015. Antioxidants safeguard telomeres in bold chicks. *Biology letters* 11, 20150211.
- Kirby, R., Alldredge, M.W., Pauli, J.N., 2017. Environmental, not individual, factors drive markers of biological aging in black bears. *Evolutionary Ecology in press*.
- Kotrschal, A., Ilmonen, P., Penn, D.J., 2007. Stress impacts telomere dynamics. *Biology letters* 3, 128–130.
- Kratschmar, D.V., Calabrese, D., Walsh, J., Lister, A., Birk, J., Appenzeller-Herzog, C., Moulin, P., Goldring, C.E., Odermatt, A., 2012. Suppression of the Nrf2-dependent antioxidant response by glucocorticoids and 11B-HSD1-mediated glucocorticoid activation in hepatic cells. *Plos One* 7, e36774.

- Krause, J.S., Pérez, J.H., Chmura, H.E., Meddle, S.L., Hunt, K.E., Gough, L., Boelman, N., Wingfield, J.C., 2016. The stress response is attenuated during inclement weather in parental, but not pre-parental, Lapland longspurs (*Calcarius lapponicus*). *Hormones and Behavior* 83, 68-74.
- Kurz, D.J., Decary, S., Hong, Y., Trivier, E., Akhmedov, A., Erusalimsky, J.D., 2004. Chronic oxidative stress compromises telomere integrity and accelerates the onset of senescence in human endothelial cells. *Journal of Cell Science* 117, 2417–2426.
- Landys, M.M., Ramenofsky, M., Wingfield, J.C., 2006. Actions of glucocorticoids at a seasonal baseline as compared to stress-related levels in the regulation of periodic life processes. *General and comparative endocrinology* 148, 132–149.
- Lansdorp, P.M., 2009. Telomeres and disease. *The EMBO Journal* 28, 2532–2540.
- Lee, W.-S., Monaghan, P., Metcalfe, N.B., 2013. Experimental demonstration of the growth rate–lifespan trade-off. *Proceedings of the Royal Society of London B: Biological Sciences* 280, 20122370.
- Lendvai, A.Z., Giraudeau, M., Chastel, O., 2007. Reproduction and modulation of the stress response: an experimental test in the house sparrow. *Proceedings of the Royal Society of London B: Biological Sciences* 274, 391–397.
- Lewin, N., Treidel, L.A., Holekamp, K.E., Place, N.J., Hausmann, M.F., 2015. Socioecological variables predict telomere length in wild spotted hyenas. *Biology letters* 11, 20140991.
- Liu, L., Bailey, S.M., Okuka, M., Muñoz, P., Li, C., Zhou, L., Wu, C., Czerwicz, E., Sandler, L., Seyfang, A., 2007. Telomere lengthening early in development. *Nature cell biology* 9, 1436–1441.
- Love, O.P., Madliger, C.L., Bourgeon, S., Semeniuk, C.A., Williams, T.D., 2014. Evidence for baseline glucocorticoids as mediators of reproductive investment in a wild bird. *General and Comparative Endocrinology* 199, 65–69.
- Madliger, C.L., Love, O.P., 2016. Employing individual measures of baseline glucocorticoids as population-level conservation biomarkers: considering within-individual variation in a breeding passerine. *Conservation physiology* 4, cow048.
- Martin, L.B., 2009. Stress and immunity in wild vertebrates: timing is everything. *General and comparative endocrinology* 163, 70–76.
- McEwen, B.S., Wingfield, J.C., 2003. The concept of allostasis in biology and biomedicine. *Hormones and Behavior* 43, 2–15.
- Meillère, A., Brischoux, F., Bustamante, P., Michaud, B., Parenteau, C., Marciau, C., Angelier, F., 2016. Corticosterone levels in relation to trace element contamination along an urbanization gradient in the common blackbird (*Turdus merula*). *Science of the Total Environment* 566, 93–101.
- Meillère, A., Brischoux, F., Ribout, C., Angelier, F., 2015. Traffic noise exposure affects telomere length in nestling house sparrows. *Biology Letters* 11, 20150559.

- Mizutani, Y., Niizuma, Y., Yoda, K., 2016. How Do Growth and Sibling Competition Affect Telomere Dynamics in the First Month of Life of Long-Lived Seabird? *PloS One* 11, e0167261.
- Mizutani, Y., Tomita, N., Niizuma, Y., Yoda, K., 2013. Environmental perturbations influence telomere dynamics in long-lived birds in their natural habitat. *Biology Letters* 9, 20130511.
- Monaghan, P., 2014. Organismal stress, telomeres and life histories. *Journal of Experimental Biology* 217, 57–66.
- Monaghan, P., 2010. Telomeres and life histories: the long and the short of it. *Annals of the New York Academy of Sciences* 1206, 130–142.
- Monaghan, P., Haussmann, M.F., 2006. Do telomere dynamics link lifestyle and lifespan? *Trends in Ecology & Evolution* 21, 47–53.
- Näslund, J., Pauliny, A., Blomqvist, D., Johnsson, J.I., 2015. Telomere dynamics in wild brown trout: effects of compensatory growth and early growth investment. *Oecologia* 177, 1221–1230.
- Nettle, D., Andrews, C., Reichert, S., Bedford, T., Gott, A., Parker, C., Kolenda, C., Martin-Ruiz, C., Monaghan, P., Bateson, M., 2016. Brood size moderates associations between relative size, telomere length, and immune development in European starling nestlings. *Ecology and Evolution* 6, 8138–8148.
- Nettle, D., Monaghan, P., Boner, W., Gillespie, R., Bateson, M., 2013. Bottom of the heap: having heavier competitors accelerates early-life telomere loss in the European starling, *Sturnus vulgaris*. *PLoS One* 8, e83617.
- Nettle, D., Monaghan, P., Gillespie, R., Brilot, B., Bedford, T., Bateson, M., 2015. An experimental demonstration that early-life competitive disadvantage accelerates telomere loss. *Proceedings of the Royal Society of London B: Biological Sciences* 282, 20141610.
- Njajou, O.T., Cawthon, R.M., Damcott, C.M., Wu, S.-H., Ott, S., Garant, M.J., Blackburn, E.H., Mitchell, B.D., Shuldiner, A.R., Hsueh, W.-C., 2007. Telomere length is paternally inherited and is associated with parental lifespan. *Proceedings of the National Academy of Sciences* 104, 12135–12139.
- Noguera, J.C., Metcalfe, N.B., Boner, W., Monaghan, P., 2015. Sex-dependent effects of nutrition on telomere dynamics in zebra finches (*Taeniopygia guttata*). *Biology letters* 11, 20140938.
- Nordstad, T., Moe, B., Bustnes, J.O., Bech, C., Chastel, O., Goutte, A., Sagerup, K., Trouvé, C., Herzke, D., Gabrielsen, G.W., 2012. Relationships between POPs and baseline corticosterone levels in black-legged kittiwakes (*Rissa tridactyla*) across their breeding cycle. *Environmental pollution* 164, 219–226.
- Nussey, D.H., Baird, D., Barrett, E., Boner, W., Fairlie, J., Gemmell, N., Hartmann, N., Horn, T., Haussmann, M., Olsson, M., 2014. Measuring telomere length and telomere dynamics in evolutionary biology and ecology. *Methods in Ecology and Evolution* 5, 299–310.

- Olsson, M., Pauliny, A., Wapstra, E., Blomqvist, D., 2010. Proximate determinants of telomere length in sand lizards (*Lacerta agilis*). *Biology Letters* 6, 651–653.
- Olsson, M., Pauliny, A., Wapstra, E., Uller, T., Schwartz, T., Blomqvist, D., 2011. Sex differences in sand lizard telomere inheritance: paternal epigenetic effects increases telomere heritability and offspring survival. *PLoS One* 6, e17473.
- Ouyang, J.Q., Lendvai, A.Z., Moore, I.T., Bonier, F., Haussmann, M.F., 2016. Do hormones, telomere lengths, and oxidative stress form an integrated phenotype? A case-study in free-living tree swallows. *Integrative and Comparative Biology* 56, 138–145.
- Parolini, M., Romano, A., Khorauli, L., Nergadze, S.G., Caprioli, M., Rubolini, D., Santagostino, M., Saino, N., Giulotto, E., 2015. Early life telomere dynamics differ between the sexes and predict growth in the Barn swallow (*Hirunda rustica*). *Plos One* 10:e0142530.
- Patel, R., McIntosh, L., McLaughlin, J., Brooke, S., Nimon, V., Sapolsky, R., 2002. Disruptive effects of glucocorticoids on glutathione peroxidase biochemistry in hippocampal cultures. *Journal of Neurochemistry* 82, 118–125.
- Pauliny, A., Wagner, R.H., Augustin, J., Szep, T., Blomqvist, D., 2006. Age-independent telomere length predicts fitness in two bird species. *Molecular Ecology* 15, 1681–1687.
- Pauliny, A., Larsson, K., Blomqvist, D., 2012. Telomere dynamics in a long-lived bird, the barnacle goose. *BMC Evolutionary Biology* 12, 257.
- Plot, V., Criscuolo, F., Zahn, S., Georges, J.-Y., 2012. Telomeres, age and reproduction in a long-lived reptile. *PloS One* 7, e40855.
- Quirici, V., Guerrero, C.J., Krause, J.S., Wingfield, J.C., Vásquez, R.A., 2016. The relationship of telomere length to baseline corticosterone levels in nestlings of an altricial passerine bird in natural populations. *Frontiers in Zoology* 13, 1.
- Rattiste, K., Klandorf, H., Urvik, J., Sepp, T., Asghar, M., Hasselquist, D., Coe, C., Hörak, P., 2015. Skin pentosidine and telomere length do not covary with age in a long-lived seabird. *Biogerontology* 16, 435–441.
- Reichert, S., Rojas, E.R., Zahn, S., Robin, J.P., Criscuolo, F., Maselein, S., 2015. Maternal telomere length inheritance in the king penguin. *Heredity* 114, 10–16.
- Reichert, S., Stier, A., Zahn, S., Arrivé, M., Bize, P., Maselein, S., Criscuolo, F., 2014. Increased brood size leads to persistent eroded telomeres. *Frontiers in Ecology and Evolution* 2, 9.
- Ricklefs, R.E., 2010. Embryo growth rates in birds and mammals. *Functional Ecology* 24, 588–596.
- Ricklefs, R.E., 2006. Embryo development and ageing in birds and mammals. *Proceedings of the Royal Society of London B: Biological Sciences* 273, 2077–2082.
- Ricklefs, R.E., Wikelski, M., 2002. The physiology/life-history nexus. *Trends in Ecology & Evolution* 17, 462–468.

- Rollings, N., Miller, E., Olsson, M., 2014. Telomeric attrition with age and temperature in Eastern mosquitofish (*Gambusia holbrooki*). *Naturwissenschaften* 101, 241–244.
- Romero, L.M., 2004. Physiological stress in ecology: lessons from biomedical research. *Trends in Ecology & Evolution* 19, 249–255.
- Romero, L.M., 2002. Seasonal changes in plasma glucocorticoid concentrations in free-living vertebrates. *General and Comparative Endocrinology* 128, 1–24.
- Romero, L.M., Dickens, M.J., Cyr, N.E., 2009. The reactive scope model—a new model integrating homeostasis, allostasis, and stress. *Hormones and Behavior* 55, 375–389.
- Romero, L.M., Wingfield, J.C., 2015. *Tempests, poxes, predators, and people: stress in wild animals and how they cope*. Oxford University Press.
- Salmón, P., Nilsson, J.F., Nord, A., Bensch, S., Isaksson, C., 2016. Urban environment shortens telomere length in nestling great tits, *Parus major*. *Biology Letters* 12, 20160155.
- Salomons, H.M., Mulder, G., van de Zande, L., Haussmann, M.F., Linskens, M.H., Verhulst, S., 2009. Telomere shortening and survival in free-living corvids. *Proceedings of the Royal Society of London B: Biological Sciences* 276, 3157–3165.
- Saretzki, G., 2009. Telomerase, mitochondria and oxidative stress. *Experimental Gerontology* 44, 485–492.
- Schultner, J., Moe, B., Chastel, O., Bech, C., Kitaysky, A.S., 2014. Migration and stress during reproduction govern telomere dynamics in a seabird. *Biology Letters* 10, 20130889.
- Sebastiano, M., Eens, M., Angelier, F., Pineau, K., Chastel, O., Costantini, D., 2017. Corticosterone, inflammation, immune status and telomere length in frigatebird nestlings facing a severe herpesvirus infection. *Conservation Physiology* 5, cow073.
- Shalev, I., Entringer, S., Wadhwa, P.D., Wolkowitz, O.M., Puterman, E., Lin, J., Epel, E.S., 2013. Stress and telomere biology: a lifespan perspective. *Psychoneuroendocrinology* 38, 1835–1842.
- Simide, R., Angelier, F., Gaillard, S., Stier, A., 2016. Age and heat stress as determinants of telomere length in a long-lived fish, the Siberian sturgeon. *Physiological and Biochemical Zoology* 89, 441–447.
- Sletten, S., Bourgeon, S., Bårdsen, B.-J., Herzke, D., Criscuolo, F., Massemin, S., Zahn, S., Johnsen, T.V., Bustnes, J.O., 2016. Organohalogenated contaminants in white-tailed eagle (*Haliaeetus albicilla*) nestlings: An assessment of relationships to immunoglobulin levels, telomeres and oxidative stress. *Science of the Total Environment* 539, 337–349.
- Smith, S.M., Nager, R.G., Costantini, D., 2016. Meta-analysis indicates that oxidative stress is both a constraint on and a cost of growth. *Ecology and evolution* 6, 2833–2842.
- Sohal, R.S., Orr, W.C., 2012. The redox stress hypothesis of aging. *Free Radical Biology and Medicine* 52, 539–555.

- Soler, J.J., Ruiz Castellano, C., Martínez-de la Puente, J., Tomás, G., Ruiz-Rodríguez, M., Figuerola, J., 2015. Telomere dynamics in parasitic great spotted cuckoos and their magpie hosts. *Journal of Evolutionary Biology* 28, 1610–1617.
- Stauffer, J., Panda, B., Eeva, T., Rainio, M., Ilmonen, P., 2017. Telomere damage and redox status alterations in free-living passerines exposed to metals. *Science of the Total Environment* 575, 841–848.
- Stier, A., Delestrade, A., Bize, P., Zahn, S., Criscuolo, F., Massemin, S., 2016. Investigating how telomere dynamics, growth and life history covary along an elevation gradient in two passerine species. *Journal of Avian Biology* 47, 134–140.
- Stier, A., Massemin, S., Zahn, S., Tissier, M.L., Criscuolo, F., 2015. Starting with a handicap: effects of asynchronous hatching on growth rate, oxidative stress and telomere dynamics in free-living great tits. *Oecologia* 179, 999–1010.
- Stier, A., Viblanc, V.A., Massemin-Challet, S., Handrich, Y., Zahn, S., Rojas, E.R., Saraux, C., Le Vaillant, M., Prud'Homme, O., Grosbellet, E., *et al.* 2014. Starting with a handicap: phenotypic differences between early-and late-born king penguin chicks and their survival correlates. *Functional ecology* 28, 601–611.
- Sudyka, J., Arct, A., Drobnik, S., Dubiec, A., Gustafsson, L., Cichoń, M., 2014. Experimentally increased reproductive effort alters telomere length in the blue tit (*Cyanistes caeruleus*). *Journal of Evolutionary Biology* 27, 2258–2264.
- Sudyka, J., Casasole, G., Rutkowska, J., Cichoń, M., 2016. Elevated reproduction does not affect telomere dynamics and oxidative stress. *Behavioral Ecology and Sociobiology* 70, 2223–2233.
- Tartu, S., Angelier, F., Bustnes, J.O., Moe, B., Hanssen, S.A., Herzke, D., Gabrielsen, G.W., Verboven, N., Verreault, J., Labadie, P., Budzinski, H., Wingfield, J.C., Chastel, O., 2015. Polychlorinated biphenyl exposure and corticosterone levels in seven polar seabird species. *Environmental Pollution* 197, 173–180.
- Tissier, M.L., Williams, T.D., Criscuolo, F., 2014. Maternal effects underlie ageing costs of growth in the zebra finch (*Taeniopygia guttata*). *PLoS One* 9, e97705.
- Turbill, C., Ruf, T., Smith, S., Bieber, C., 2013. Seasonal variation in telomere length of a hibernating rodent. *Biology Letters* 9, 20121095.
- Turbill, C., Smith, S., Deimel, C., Ruf, T., 2012. Daily torpor is associated with telomere length change over winter in Djungarian hamsters. *Biology Letters* 8, 304–307.
- Ujvari, B., Biro, P.A., Charters, J.E., Brown, G., Heasman, K., Beckmann, C., Madsen, T., 2016. Curvilinear telomere length dynamics in a squamate reptile. *Functional Ecology* 31, 753–759.
- Ujvari, B., Madsen, T., 2009. Short telomeres in hatchling snakes: erythrocyte telomere dynamics and longevity in tropical pythons. *PLoS one* 4, e7493.

- Valdes, A.M., Andrew, T., Gardner, J.P., Kimura, M., Oelsner, E., Cherkas, L.F., Aviv, A., Spector, T.D., 2005. Obesity, cigarette smoking, and telomere length in women. *The Lancet* 366, 662–664.
- Vleck, C.M., Haussmann, M.F., Vleck, D., 2007. Avian senescence: underlying mechanisms. *Journal of Ornithology* 148, 611–624.
- Vleck, C.M., Haussmann, M.F., Vleck, D., 2003. The natural history of telomeres: tools for aging animals and exploring the aging process. *Experimental Gerontology* 38, 791–795.
- Voillemot, M., Hine, K., Zahn, S., Criscuolo, F., Gustafsson, L., Doligez, B., Bize, P., 2012. Effects of brood size manipulation and common origin on phenotype and telomere length in nestling collared flycatchers. *BMC Ecology* 12, 17.
- von Zglinicki, T., 2002. Oxidative stress shortens telomeres. *Trends in Biochemical Sciences* 27, 339–344.
- Watson, H., Bolton, M., Monaghan, P., 2015. Variation in early-life telomere dynamics in a long-lived bird: links to environmental conditions and survival. *Journal of Experimental Biology* 218, 668–674.
- Wilson, A.J., Nussey, D.H., 2010. What is individual quality? An evolutionary perspective. *Trends in Ecology and Evolution* 25, 207–214.
- Wingfield, J.C., 2013. The comparative biology of environmental stress: behavioural endocrinology and variation in ability to cope with novel, changing environments. *Animal Behaviour* 85, 1127–1133.
- Wingfield, J.C., 2008. Organization of vertebrate annual cycles: implications for control mechanisms. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363, 425–441.
- Wingfield, J.C., 2003. Control of behavioural strategies for capricious environments. *Animal behaviour* 66, 807–816.
- Wingfield, J.C., Maney, D.L., Breuner, C.W., Jacobs, J.D., Lynn, S., Ramenofsky, M., Richardson, R.D., 1998. Ecological bases of hormone—behavior interactions: the “emergency life history stage.” *American Zoologist* 38, 191–206.
- Wingfield, J.C., Sapolsky, R.M., 2003. Reproduction and resistance to stress: when and how. *Journal of neuroendocrinology* 15, 711–724.
- Wingfield, J.C., Krause, J.S., Pérez, J.H., Chmura, H.E., Németh, Z., Word, K.R., Calisi, R.M., Meddle, S.L., 2015. A mechanistic approach to understanding the range shifts in a changing world: What makes a pionner? *General and Comparative Endocrinology* 222, 44–53.
- Young, R.C., Kitaysky, A.S., Haussmann, M.F., Descamps, S., Orben, R.A., Elliott, K.H., Gaston, A.J., 2013. Age, sex, and telomere dynamics in a long-lived seabird with male-biased parental care. *PLoS One* 8, e74931.

- Young, R.C., Kitaysky, A.S., Barger, C.P., Dorresteijn, I., Ito, M., Watanuki, Y., 2015. Telomere length is a strong predictor of foraging behavior in a long-lived seabird. *Ecosphere* 6, 39.
- Young, R.C., Barger, C.P., Dorresteijn, I., Haussmann, M.F., Kitaysky, A.S., 2016. Telomere length and environmental conditions predict stress levels but not parental investment in a long-lived seabird. *Marine Ecology Progress Series* 556, 251–259.
- Young, R.C., Orben, R.A., Will, A.P., Kitaysky, A.S., 2017. Relationship between telomere dynamics and movement and behavior during winter in the thick-billed murre. *Marine Ecology Progress Series* in press.
- Zeichner, S.L., Palumbo, P., Feng, Y., Xiao, X., Gee, D., Sleasman, J., Goodenow, M., Biggar, R., Dimitrov, D., 1999. Rapid telomere shortening in children. *Blood* 93, 2824–2830.
- Zera, A.J., Harshman, L.G., 2001. The physiology of life history trade-offs in animals. *Annual review of Ecology and Systematics* 32, 95–126.

Figure 1: Trends in the number of studies published each year on different topics from 1995 to 2016. These numbers were found by conducting a search in ISI web of knowledge (search terms, filled circles: telomere, empty circles: telomere and stress, triangles: telomere and glucocorticoids). For all searches, the results were limited to the following categories: ecology, zoology, behavioral sciences, environmental sciences and evolutionary biology.

Figure 2: Schematic representation of individual telomere dynamics through life in wild vertebrates (thick black line): (1) initial telomere length is partly heritable and determined by epigenetic/trans-generational effects; (2) telomeres shorten very quickly during the developmental phase; (3) telomere shortening is much slower during adulthood. For each of these steps, there is an important inter-individual variation in telomere dynamics: (1) Inter-individual variability in initial telomere length is represented by a grey vertical arrow; (2) inter-individual variability in developmental telomere dynamics is represented by a few examples of possible telomere trajectories (thin grey lines); (3) inter-individual variability in adult telomere dynamics is represented by a few examples of possible telomere trajectories (thin black lines). This variability in telomere dynamics translate into large inter-individual differences in telomere length (the theoretical possible range of telomere length is represented by thick grey dashed lines).

The diagram illustrates the role of telomerase in telomere maintenance and the impact of oxidative stress. On the left, the chemical structure of Corticosterone (or Cortisol) is shown. Two pathways lead from it: 'Antioxidant gene expression' and 'Metabolism'. 'Antioxidant gene expression' leads to a cycle involving GPX (Glutathione Peroxidase) and GR (Glutathione Reductase). GPX converts H_2O or ROH to H_2O_2 or ROOH , while GR converts GSSG to GSH. 'Metabolism' leads to a cycle involving NADPH and NADP^+ . The Fenton Reaction is shown where Fe^{2+} reacts with H_2O_2 or ROOH to produce Fe^{3+} and reactive oxygen species ($\text{OH}^\bullet$, $\text{RO}^\bullet$ or $\text{ROO}^\bullet$). These species lead to the formation of Telomeres (represented by a DNA double helix). Telomerase (represented by a yellow hexagon) is shown to maintain the length of Telomeres. A feedback loop is indicated by an arrow from Telomeres back to the Antioxidant gene expression pathway.

Figure 4: Theoretical links between the GC stress response, immediate survival and telomere preservation. A: An intense GC stress response is theoretically associated with an improved immediate survival when an acute stressor occurs (blue line) but also with a faster telomere attrition (orange line). B: The relationship between the intensity of the GC stress response and telomere dynamics may be different in high-quality individuals or in individuals living in a highly suitable environment. Because of better antioxidant defenses or DNA repair mechanisms, these individuals may be able to limit the negative effect of elevated GC levels on telomere length (the solid and the dashed line respectively represent individuals of low and high quality). C: Elevated GC levels can switch individual from a specific life-history stage (e.g. breeding stage) to an emergency life-history stage (e.g. non breeding stage). This switch is associated with reduced demanding activities, and therefore, with limited telomere attrition. This switch from a specific life-history stage to an emergency life history stage is represented by the orange arrow.

Variable	Relationship with telomere length/dynamics	Stage	Type of study
Cohorts	≠	Development	correlative
Cohorts	≠	Development	correlative
Cohorts	≠	Development	correlative
Cohorts	≠	Development	correlative
Years	≠	Adulthood	correlative
Sites (longitude)	≠	Adulthood	correlative
Sites (latitude)	≠	Adulthood	correlative
Sites (elevation)	≠	Development	correlative
Sites (elevation)	≠	Development	correlative
Sites (latitude)	≠	Adulthood	correlative
Sites (elevation)	≠	Adulthood	correlative
Habitat of low suitability	-	Adulthood	correlative
Habitat of low suitability	-	Adulthood	correlative
Habitat of low suitability	-	Development	experimental/correlative
Habitat of low suitability	=	Development	correlative

≠ denotes a significant difference in telomere length or telomere dynamics between cohorts, years or sites

= denotes similar telomere length or telomere dynamics between cohorts, years or sites

- denotes a negative effect of the variable on telomere length or telomere dynamics

Variable	Relationship with telomere length/dynamics	Stage	Type of study	
<i>Infection/pathogens/parasites</i>				
Multiple infections	-	Adulthood	experimental	M
Tuberculosis infection	-	Adulthood	correlative	M
Malaria infection	0	Adulthood	experimental	
Malaria infection	-	Adulthood	experimental	
Immune challenge	-	adulthood	experimental	
Herpes virus infection	0	Development	correlative	
<i>Social factors</i>				
Social isolation	-	Adulthood	correlative	
Dominance rank	+	Adulthood	correlative	M
<i>Nutritional factors</i>				
Nutritional antioxidant	-*	Adulthood	experimental	
Nutritional micronutrient supplementation	0	Development	experimental	
Nutritional micronutrient supplementation	+/0**	Adulthood	experimental	
Nuritional antioxidant	+/0***	Development	experimental	
Food availability	0	Adulthood	correlative	M
Food availability	+	Adulthood	correlative	
Food supplementation	+	Adulthood	experimental	M
<i>Developmental factors</i>				
Maternal age	+	Development	correlative	
Maternal age	+	Development	correlative	
Parental age	0	Development	correlative	
Parental age	+	Development	correlative	
Brood competition	0	Development	experimental	
Brood competition	-	Development	experimental	
Brood competition	-	Development	experimental	
Brood competition	-	Development	experimental	
Brood competition	-	Development	experimental	
Brood competition	-	Development	correlative	
Brood competition	-	Development	experimental	
Brood competition	-	Development	correlative	
Brood competition	-	Development	correlative	
Brood competition	-	Development	experimental	
Brood competition disadvantage	-	Development	correlative	
Growth	+	Development	correlative	
Growth	0	Development	correlative	

Growth	+	Development	correlative
Growth	+	Development	correlative
Accelerated growth	-	Development	correlative
Accelerated growth	0	Development	experimental
Slow growth	-	Development	correlative
Developmental constraints	-	Development	correlative

(Table 2 continued)

Reproductive factors

Reproductive effort	-	Adulthood	experimental
Reproductive effort	0	Adulthood	experimental
Reproductive effort	-	Adulthood	correlative
Reproductive effort	-	Adulthood	correlative
Reproductive effort	-	Adulthood	experimental
Reproductive effort	-	Adulthood	experimental
Reproductive effort	0	Adulthood	experimental

Climatic factors

Climate (El Nino)	+	Adulthood	correlative
Reduced temperature	-	Development	experimental
Increased temperature	-	Adulthood	experimental
Increased temperature	-	Adulthood	experimental

Stressors

Crowding	-	adulthood	experimental
Handling	-	Development	experimental
Disturbance	-	adulthood	experimental
Noise	-	Development	experimental

Contamination

Heavy metals	-	Development	correlative
Heavy metals	0	Development	correlative
Heavy metals	0	Adulthood	correlative
Heavy metals	0	Adulthood	correlative
Chlordanes	-/0****	Adulthood	correlative
Persistent organic pollutants (Σ POPs)	0	Adulthood	correlative
Polychlorinated biphenyls (Σ PCBs)	0	Adulthood	correlative
Hexachlorobenzen (HCBs)	0	Adulthood	correlative
Dichlorodiphenyldichloroethylen (pp'-DDE)	0	Adulthood	correlative
Perfluoroalkylated substances (PFASs)	+	Adulthood	correlative
Mercury	+/0**	Adulthood	correlative
Organochlorine pesticides (OCPs)	0	Development	correlative
Perfluoroalkylated substances (PFASs)	0	Development	correlative

Other factors

Tail regrowth

-

Adulthood

correlative

R

* This negative effect is indirectly linked to increased reproductive effort in experimental birds

** the effect was apparent in females only

*** The positive effect of antioxidant supplementation was only apparent for bold chicks

**** the effect was apparent for oxychlordanes in females only

ACCEPTED MANUSCRIPT

GC measurement	Relationship with telomere length	Stage	Taxon	Comment
<i>correlative studies</i>				
Feather corticosterone	+	Adulthood	Bird	Thi
Plasma corticosterone	-	Adulthood	Bird	Thon
Plasma corticosterone	.*	Adulthood	Bird	C
Plasma corticosterone	+	Adulthood	Bird	Thi
Plasma corticosterone	0	Adulthood	Bird	Thi
Plasma corticosterone	-	Adulthood	Bird	Thi
Plasma corticosterone	-	Adulthood	Bird	Black-
Feather corticosterone	-	Adulthood	Bird	A
Plasma corticosterone	0	Adulthood	Bird	T
+ and - respectively denote a positive and negative relationship between GC levels and telomere length				
0 denotes no significant relationship between GC levels and telomere length				
* the relationship was apparent in males only				
<i>Experimental studies</i>				
Embryonic injection of corticosterone	-	Development	Bird	Do
Ingestion of corticosterone	-	Development	Bird	E
Corticosterone implants	-	Adulthood	Bird	Black
Injection of corticosterone	-	Adulthood	Bird	
- denotes a negative effect of the experiment on telomere length				

Highlights

- Telomere length is often studied in wild animals because of its link with fitness
- We review the link among stress, glucocorticoids (GC), and telomere dynamics.
- Telomere dynamics is influenced by environmental stressors and life-history events.
- GCs could mediate the impact of the environment on telomere dynamics.
- Coping styles and the GC stress response may be involved in telomere dynamics.