

Ossifying fibroma of the mandible

Published on 04.12.2016

DOI: 10.1594/EURORAD/CASE.14188

ISSN: 1563-4086

Section: Head & neck imaging

Area of Interest: Head and neck Bones

Procedure: Diagnostic procedure

Imaging Technique: CT

Imaging Technique: Percutaneous

Special Focus: Neoplasia Case Type: Clinical Cases

Authors: Arne Impens¹, Adelard De Backer¹, Carine Petré², Herwig Van Dijck³, Filip Vanhoenacker^{2,4,5}

Patient: 54 years, male

Clinical History:

A 54-year-old man presented to the maxillofacial surgeon with a slowly increasing swelling of the lower left jaw for more than 6 months. The lesion was painless, hard and immobile on palpation.

Imaging Findings:

A cone beam CT (CBCT) of the mandible showed a well-circumscribed osseous lesion in the left mandible, measuring 46 x 22 x 33 mm. The lesion was radiopaque with a ground glass appearance. There was bone expansion with cortical thinning but absence of cortical breakthrough and no periosteal reaction (Fig.1-4). The mandibular canal was displaced inferiorly.

A biopsy was performed and subsequent histopathological examination showed cellular connective tissue with an osteoblastic rim on the surface of woven bone intermingled with a fibrous component without atypia (Fig. 5). Diagnosis of ossifying fibroma was made and the lesion was partially resected.

Discussion:

Ossifying fibroma is a benign neoplasm composed of fibrous tissue and mineralized material. The previously used term cemento-ossifying fibroma was renamed as ossifying fibroma according to the revised WHO classification (2005) [1, 2].

It is thought that ossifying fibromas of the jaw arise from the periodontal ligament but this remains controversial as similar lesions may appear in bones that lack this structure. Some authors suggest a possible relationship with previous trauma as a triggering factor [3]. The craniofacial bones are the most frequently involved with a predilection for the region inferior to the molars and premolars of the mandible. Ossifying fibromas are most frequently diagnosed during the third and fourth decades with a marked female predilection (female:male ratio: 4:1). A juvenile variant, ore aggressive, exists that most commonly affects boys younger than 15 years [1, 3, 4].

The clinical presentation usually consists of a painless slowly growing mass. Lesion expansion may result in tooth displacement, root resorption and facial asymmetry if untreated. The mass is covered with normal mucosa. Lymphadenopathy is not seen [3-6].

Different radiographic patterns have been described depending on the amount of mineralized tissue. Most frequently the lesion is mixed radiopaque-radiolucent and less commonly either radiolucent or radiopaque. The degree of opacity of the lesion depends on its age. With maturation the lesion becomes more radiopaque due to progressive matrix mineralization. The lesion is usually unilocular although infrequently a multilocular appearance is seen. The main differential diagnosis consists of fibrous dysplasia if the lesion has a ground-glass attenuation. The margins of the lesion are always well demarcated whereas fibrous dysplasia may have ill-defined borders. The mandibular

canal may be displaced inferiorly while in fibrous dysplasia the canal may be displaced in any direction. Usually the lesion displays concentric growth and buccolingual expansion. Tooth displacement and root resorption are seen in larger lesions [5, 7, 8].

The recommended treatment is complete surgical resection or enucleation since incomplete resection has a high recurrence rate. In our case complete resection was impossible due the proximity of the lesion to the mandibular canal. Surgical treatment may require bone grafting or reconstructive surgery. The estimated recurrence rate after appropriate surgical removal is 12% [5, 7].

Differential Diagnosis List: Ossifying fibroma, Fibrous dysplasia, Cementoblastoma, Paget's disease, Periapical cemental dysplasia


Final Diagnosis: Ossifying fibroma

References:

- Barnes L, Eveson JW, Reichart P, Sidransky D (2005) World Health Organization classification of tumors, pathology and genetics. Head and neck tumors p. 319
- de Andrade M, Silva-Sousa YT, Marques MF, Pontual ML, Ramos-Perez FM, Perez DE (2013) Ossifying Fibroma of the Jaws: A Clinicopathological Case Series Study. *Braz Dent J* 24(6), 662-6 (PMID: [24474366](#))
- Silvestre-Rangil J, Silvestre FJ, Requeni-Bernal J (2011) Cemento-ossifying fibroma of the mandible: Presentation of a case and review of the literature. *J. Clin. Exp. Dent* 3(1):e66-9
- Curé JK, Vattoth S, Shah R (2012) Radiopaque jaw lesions: an approach to the differential diagnosis. *Radiographics* 32:1909–1925 (PMID: [23150848](#))
- Liu Y, Wang H, You M, Yang Z, Miao J, Shimizutani K, Koseki T (2010) Ossifying fibromas of the jaw bone: 20 cases. *Dentomaxillofac Radiol* 39(1):57-63 (PMID: [20089746](#))
- Trijolet JP, Parmentier J, Sury F, Goga D, Mejean N, Laure B (2011) Cemento-ossifying fibroma of the mandible. *Eur Ann Otorhinolaryngol Head Neck Dis* 128(1):30-3 (PMID: [20846921](#))
- MacDonald-Jankowski DS (2009) Ossifying fibroma: a systematic review. *Dentomaxillofac Radiol* 38(8):495-513 (PMID: [20026707](#))
- Sontakke SA, Karjodkar FR, Umarji HR (2011) Computed tomographic features of fibrous dysplasia of maxillofacial region. *Imaging Sci Dent* 41(1): 23-8 (PMID: [21977470](#))

Figure 1


a



Description: Scout view demonstrates relative radiolucency at the left mandibular angle with an indistinct cortical margin (white arrows). **Origin:** F.M. Vanhoenacker, Department of radiology, AZ Sint-Maarten, Duffel-Mechelen, Belgium

Figure 2

a
P3



Description: Panoramic reformatted CBCT image of the mandible demonstrates a radiopaque lesion causing inferior displacement of the mandibular canal (white arrows). **Origin:** F.M. Vanhoenacker, Department of radiology, AZ Sint-Maarten, Duffel-Mechelen, Belgium

Figure 3

a



Description: CBCT of the mandible showing an expansile, well-circumscribed lesion with ground glass appearance and buccolingual expansion in the left mandible (white arrows). **Origin:** F.M. Vanhoenacker, Department of radiology, AZ Sint-Maarten, Duffel-Mechelen, Belgium

Figure 4


a



Description: Parasagittal reformatted image demonstrates inferior displacement of the mandibular canal (white arrow), buccolingual expansion and ground-glass attenuation. **Origin:** F.M. Vanhoenacker, Department of radiology, AZ Sint-Maarten, Duffel-Mechelen, Belgium

Figure 5

a



Description: Histological examination shows mineralized material composed by woven bone, predominantly bordered by osteoblasts and interspersed by cellularized connective tissue (H & E staining). **Origin:** F.M. Vanhoenacker, Department of radiology, AZ Sint-Maarten, Duffel-Mechelen, Belgium