

Avulsion fracture of the tibial tubercle in an adolescent

Published on 20.09.2016

DOI: 10.1594/EURORAD/CASE.14039

ISSN: 1563-4086

Section: Musculoskeletal system

Area of Interest: Musculoskeletal joint Musculoskeletal bone

Procedure: Diagnostic procedure

Procedure: Normal variants

Imaging Technique: MR

Imaging Technique: Conventional radiography

Special Focus: Trauma Case Type: Clinical Cases

Authors: Charlotte Vanhoenacker 1, Kris Van Crombrugge 2, Lieven Tack 2, Filip Vanhoenacker 2,3,4

Patient: 13 years, male

Clinical History:

A 13-year-old boy presented to the emergency department of our hospital with a painful, swollen left knee after high jumping in gym class. During take-off, he felt a sudden snap in his knee, after landing he was unable to bear weight. Clinical examination revealed the impossibility to extend the knee.

Imaging Findings:

Lateral radiograph revealed soft tissue swelling and an enlargement of the growth plate of the secondary ossification centre of the tibial tubercle (Fig. 1).

Subsequent MRI was performed for further evaluation of the extent of the lesion. MRI confirmed the presence of an avulsion fracture of the secondary ossification centre of the tibial tuberosity. The maximum gap of the growth plate was located anterolaterally and measured seven millimetres. In the ventral part of the proximal tibial epiphysis, bone marrow oedema was visualized in keeping with lesion extension in the femoro-tibial joint space (Fig. 2).

Haemorrhagic effusion of the deep tibial tubercle bursa and the infrapatellar subcutaneous bursa was seen as well.

Discussion:

An avulsion fracture is caused by a high energy tensile force on a ligament or tendon which exceeds the strength of the bony tissue. In adolescents the growth plate is the weakest link, in comparison to adults where a tensile force will cause injuries of the musculotendinous junctions [1]. Typically avulsion fractures occur in boys between the age of 13-17 years because of the cartilage features of the growth plate. Avulsion fractures of the knee are uncommon compared to avulsion fractures of the hip or the pelvis [2]. Fractures of the apophysis of the proximal tibia comprise 0.5% - 3% of all fractures [3].

The mechanism of injury is thought to be a high energy extension of the knee against a fixed leg, such as in jumping sports [4].

Avulsion fractures of the tibial tubercle (TT) are classified according to the modified Watson-Jones classification (Fig. 3). The original classification of Watson and Jones described three categories of avulsion fractures of the tibial tubercle [5]. Ryu and Debenham introduced an additional fourth category in which the fracture extends into the proximal tibial epiphysis. The extent of the injury is related to the stage of fusion of the epiphysis [6].

Plain radiographs are the first step to define the displacement of the fracture, the degree of comminution and the

extent of the injury [1].

Although MRI is not always mandatory, it may be of additional value to define the precise extent of the bone and soft tissue abnormalities, and allows a more accurate classification of the lesion. In our case, the plain radiograph showed an enlargement of the growth plate of the TT. In addition, MRI revealed bone marrow oedema in the ventral part of the proximal tibial epiphysis indicating lesion extension in the articular surface of the tibia. After MRI the lesion was classified as a type II fracture.

The treatment, either conservative versus surgical, depends on correct staging of the lesion. If displacement is marked or if the physis is comminuted, surgical approach is recommended [7]. In our case the maximum gap of the growth plate measured seven millimetres, therefore surgical repair, consisting of open reduction and internal fixation with cortical screws, was performed, followed by six weeks of immobilization in a plaster cast.

Further follow-up was uneventful five months after trauma.

TAKE HOME MESSAGE

Although rare, avulsion fracture of the TT should be considered in sportive adolescents particularly in jumping. MRI may add in correct staging of the lesion.

Differential Diagnosis List: Acute avulsion fracture of the apophysis of the tibial tubercle., Patellar tendon rupture, Osgood-Schlatter disease

Final Diagnosis: Acute avulsion fracture of the apophysis of the tibial tubercle.

References:

- Morrow AC, Reiter MP, Zbojniec AM, Laor T (2014) Avulsion fractures of the pediatric knee. *Pediatr Radiol* 44:1436-1445 (PMID: [25331022](#))
- Singer G, Eber R, Wegmann H, Marterer R, Kraus T, Sorantin E (2014) . Diagnosis and Treatment of apophyseal Injuries of the Pelvis in Adolescents. *Semin Musculoskelet Radiol* 18(05): 498-504 (PMID: [25350828](#))
- Crimaldi S, Calderazzi F, Becherucci L, Faidini A (2003) Upper tibial physeal fracture- a case report. Proposed mechanism of injury and classification. *Acta Orthop Scand* 74(6): 764-765 (PMID: [14763712](#))
- Clarke DO, Franklin SA, Wright DE (2016) Avulsion Fracture of the Tibial Tubercle Associated With Patellar Tendon Avulsion. *Orthopedics* 39(3): 561-4 (PMID: [26425381](#))
- Inoue G, Kuboyama K and Shido T (1991) Avulsion fractures of the proximal tibial epiphysis. *Br J Med* 25(1) (PMID: [1913033](#))
- Harb Z and Malhi A (2015) Bilateral Simultaneous Avulsion fractures of the Proximal tibia in a 14-Year-Old Athlete with Vitamin-D Deficiency. *Case Rep Orthop* (PMID: [26425381](#))
- Subbu R, Nadra R, Jordan R et al (2013) Lessons learnt from managing an avulsion fracture of the tibial tubercle extending into the tibial physis. *BMJ Case Rep* (PMID: [24057332](#))

Figure 1


a



Description: Lateral view: Soft tissue swelling (red arrow) and broadening of the growth plate of the apophysis of the tibial tubercle (yellow star). **Origin:** Vanhoenacker F, Department of Radiology, AZ Sint Maarten, Mechelen, Belgium.

Figure 2

a




Description: 1. Undulating course of patellar tendon (red arrow), adjacent soft tissue swelling. Bone marrow oedema in ventral proximal tibial epiphysis (yellow arrow) indicating lesion extension in the knee joint.

2. Broadening of the growth plate (star). **Origin:** Vanhoenacker F, Department of Radiology, AZ Sint Maarten, Mechelen, Belgium

Figure 3

a



Description: I: Upward displacement of tibial tubercle (TT).

II: Fracture extension through the articular surface.

III: Upward displacement of TT together with ossified part of tibial epiphysis.

IV: Fracture through growth plate of proximal tibial epiphysis. **Origin:** Vanhoenacker F, Department Radiology, AZ Sint Maarten, Mechelen, Belgium

Figure 4

a



Description: The avulsion fracture was treated surgically with open reduction and internal fixation with four cortical screws. Postoperative X-ray shows correct alignment of fracture and correct position of screws. **Origin:** Vanhoenacker F, Department of Radiology, AZ Sint Maarten, Mechelen, Belgium.