

Advancements in Non-Small Cell Lung Cancer

Developments in predictive biomarker testing and targeted therapy in advanced stage non-small cell lung cancer and their application across European countries



Vincent D. de Jager,^a Wim Timens,^a Arnaud Bayle,^b Johan Botling,^c Luka Brcic,^d Reinhard Büttner,^e Maria Gabriela O. Fernandes,^f Libor Havel,^g Maximilian J. Hochmair,^{h,j} Paul Hofman,^j Annelies Janssens,^k Mikael Johansson,^l Léon van Kempen,^m Izidor Kern,ⁿ Fernando Lopez-Rios,^o Margreet Lüchtenborg,^{p,q} José Carlos Machado,^{r,s} Katja Mohoric,^t Luis Paz-Ares,^u Sanjay Popat,^v Aleš Ryška,^w Phillipe Taniere,^x Jürgen Wolf,^y Ed Schuurin,^{a,aa,**} and Anthonie J. van der Wekken^{z,aa,*}



^aDepartment of Pathology and Medical Biology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

^bOncostat U1018, Inserm, Paris-Saclay University, Gustave Roussy, Villejuif, France

^cDepartment of Laboratory Medicine, Institute of Biomedicine, Sahlgrenska Academy of University of Gothenburg, Gothenburg, Sweden

^dDiagnostic and Research Institute of Pathology, Medical University of Graz, Graz, Austria

^eInstitute of Medical Genetics and Applied Genomics, University of Tuebingen, Tuebingen, Germany

^fPulmonology Department, Centro Hospitalar Universitário de São João, Porto, Portugal

^gCharles University and Thomayer Hospital, Prague, Czech Republic

^hKarl Landsteiner Institute of Lung Research and Pulmonary Oncology, Klinik Floridsdorf, Vienna, Austria

ⁱDepartment of Respiratory and Critical Care Medicine, Klinik Floridsdorf, Vienna Healthcare Group, Vienna, Austria

^jIHU RespirERA, FHU OncoAge, Nice University Hospital, Côte d'Azur University, Nice, France

^kDepartment of Oncology, University Hospital Antwerp, University of Antwerp, Edegem, Belgium

^lDepartment of Radiation Sciences, Oncology, Umeå University, Umeå, Sweden

^mDepartment of Pathology, University Hospital Antwerp, University of Antwerp, Edegem, Belgium

ⁿLaboratory for Cytology and Pathology, University Clinic Golnik, Golnik, Slovenia

^oHospital Universitario 12 de Octubre, Universidad Complutense de Madrid, Research Institute Hospital 12 de Octubre (i+12), Ciberonc, Madrid, Spain

^pNational Disease Registration Service, NHS England, London, United Kingdom

^qCentre for Cancer, Society & Public Health, King's College London, London, United Kingdom

^rInstitute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP), Porto, Portugal

^sFaculty of Medicine of the University of Porto, Institute for Research and Innovation in Health (i3S), Porto, Portugal

^tUniversity Clinic of Respiratory and Allergic Diseases, Golnik, Slovenia

^uHospital Universitario 12 de Octubre, Universidad Complutense de Madrid, H120-CNIO Lung Cancer Clinical Research Unit, Research Institute Hospital 12 de Octubre (i+12)/Spanish National Cancer Research Center (CNIO), Ciberonc, Madrid, Spain

^vLung Unit, Royal Marsden NHS Trust, London, United Kingdom

^wThe Fingerland Department of Pathology, Charles University Medical Faculty and University Hospital, Czech Republic

^xDepartment of Histopathology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom

^yLung Cancer Group Cologne, Department I for Internal Medicine and Center for Integrated Oncology Cologne/Bonn, University Hospital Cologne, Cologne, Germany

^zDepartment of Pulmonary Diseases and Tuberculosis, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

Summary

In the past two decades, the treatment of metastatic non-small cell lung cancer (NSCLC), has undergone significant changes due to the introduction of targeted therapies and immunotherapy. These advancements have led to the need for predictive molecular tests to identify patients eligible for targeted therapy. This review provides an overview of the development and current application of targeted therapies and predictive biomarker testing in European patients with advanced stage NSCLC. Using data from eleven European countries, we conclude that recommendations for predictive testing are incorporated in national guidelines across Europe, although there are differences in their comprehensiveness. Moreover, the availability of recently EMA-approved targeted therapies varies between European

The Lancet Regional Health - Europe
2024;38: 100838

Published Online 4 March 2024
<https://doi.org/10.1016/j.lanepe.2024.100838>

DOIs of original articles: <https://doi.org/10.1016/j.lanepe.2024.100841>, <https://doi.org/10.1016/j.lanepe.2024.100845>, <https://doi.org/10.1016/j.lanepe.2024.100840>, <https://doi.org/10.1016/j.lanepe.2024.100839>

*Corresponding author.

**Corresponding author.

E-mail addresses: a.j.van.der.wekken@umcg.nl (A.J. van der Wekken), e.schuuring@umcg.nl (E. Schuurin).

^{aa}These authors contributed equally to this manuscript.

countries. Unfortunately, routine assessment of national/regional molecular testing rates is limited. As a result, it remains uncertain which proportion of patients with metastatic NSCLC in Europe receive adequate predictive biomarker testing. Lastly, Molecular Tumor Boards (MTBs) for discussion of molecular test results are widely implemented, but national guidelines for their composition and functioning are lacking. The establishment of MTB guidelines can provide a framework for interpreting rare or complex mutations, facilitating appropriate treatment decision-making, and ensuring quality control.

Copyright © 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Predictive biomarker testing; Targeted therapy; Non-small cell lung cancer; Europe

Introduction

Lung cancer remains the leading cause of cancer-related mortality in Europe, resulting in a considerable number of deaths.¹ Projections for 2023 indicate that lung cancer will account for a combined total of 275,956 fatalities in the European Union and the United Kingdom.² Within the realm of lung cancer treatment, significant advancements have occurred over the past two decades, particularly in the management of non-small cell lung cancer (NSCLC), comprising approximately 85% of all lung cancer cases. These developments encompass various modalities such as minimally invasive surgery techniques and stereotactic body radiotherapy for localized disease, as well as targeted therapies and immunotherapy as systemic therapeutic options.³ As roughly 55–70% of NSCLC patients are diagnosed with metastatic disease at time of presentation, targeted therapies such as EGFR and ALK tyrosine kinase inhibitors are thought to have contributed significantly to the observed increase in overall survival rates between 2010 and 2016.⁴ Due to the expanding repertoire of European Medicines Agency (EMA) approved targeted therapies for advanced-stage NSCLC, it has become imperative to perform molecular testing to identify actionable molecular aberrations and determine the appropriate patient population for targeted treatment (Fig. 1).

The current Clinical Practice Guidelines of the European Society for Medical Oncology (ESMO) for oncogene-addicted advanced-stage NSCLC (2023) recommend molecular testing for nine predictive biomarkers.⁵ However, previous studies have described suboptimal NSCLC testing rates for targetable molecular aberrations in various European countries, suggesting that routine predictive biomarker testing is not keeping pace with the approval of targeted therapies.^{6–9} Notwithstanding the presence of European guidelines, discrepancies have also been observed between national guidelines for molecular testing in NSCLC and the European consensus, potentially contributing to variation in testing and treatment practices across individual European countries.^{6,8} Moreover, recent studies have described delayed uptake of next-generation sequencing (NGS) for oncology patients across Europe.^{9,10} Additionally, discrepancies were reported in the availability and reimbursement of precision medicines between

European countries.⁹ The latter two studies^{9,10} provided a broad overview of the European application of NGS and/or precision medicines for all (solid tumor) oncology patients, while the previously mentioned studies on NSCLC^{6–8} predominantly focused on biomarker testing rather than targeted therapy availability and by now, are from several years ago. The aim of this review is to provide an overview of the timeline, developments, and current application of targeted therapies and predictive biomarker testing in patients with advanced stage NSCLC in Europe in 2023.

The first part of this review highlights the major, recent developments in targeted therapy treatment in advanced-stage NSCLC in Europe (Section [EMA-approved targeted therapies](#)) and the changes in European guidelines for predictive biomarker testing (Section [European guidelines for predictive biomarker testing](#)). The second part provides an overview of national targeted therapy availability (Section [Availability of targeted therapies](#)), national guidelines for predictive biomarker testing (Section [National guidelines for predictive biomarker testing](#)), and molecular testing rates (Section [Predictive biomarker testing rates](#)), in a selection of European countries including Austria, Belgium, the Czech Republic, England, France, Germany, the Netherlands, Portugal, Slovenia, Spain, and Sweden. To accomplish this, one (molecular) pathologist or clinical scientist in molecular pathology and one (pulmonary) oncologist of these countries, with extensive expertise with regard to targeted treatment and molecular testing in patients with NSCLC, sought to obtain up-to-date information on their national guidelines, availabilities of targeted therapies, and molecular testing rates in patients with advanced stage non-squamous NSCLC, using a detailed questionnaire (see [Supplementary File 1](#)). Participants also provided information regarding molecular test types and molecular test result interpretation specific to their affiliated institutes (Sections [Interpretation of molecular test results](#) and [Barriers for off-label use, compassionate use programs and trial participation](#)). Collection of information from all countries was completed between March and September of 2023. The objective of this approach is to identify potential differences in predictive biomarker testing rates and targeted therapy availability for patients

Key messages

- The use of targeted therapies contributes to optimal treatment of patients with advanced stage non-small cell lung cancer (NSCLC);
- The number of European Medicine Agency (EMA)-approved targeted therapies for patients with advanced stage NSCLC is increasing rapidly;
- Predictive biomarker testing is required for the identification of the appropriate molecular aberrations that are actionable with specific targeted therapies;
- The availability of European Medicine Agency-approved targeted therapies for patients with advanced stage NSCLC varies across European countries;
- National guidelines of European countries contain recommendations for predictive biomarker testing in patients with NSCLC but only partially align with current ESMO guidelines;
- Evidence for high predictive biomarker testing rates in patients with NSCLC in Europe is limited due to the lack of (publicly) available data of national testing rates;
- National guidelines for the functioning of Molecular Tumor Boards are lacking in most European countries and should be established to provide a framework for multidisciplinary interpretation of rare or complex mutations in patients with NSCLC.

with advanced stage NSCLC among European countries that are linked by international collaboration and common legislation. In the corresponding viewpoint paper in this Clinical Series, a future perspective of the application of molecular testing in advanced stage NSCLC is described.¹¹


EMA-approved targeted therapies

Between 2005 and 2018, the EMA granted approval for eight targeted therapies in advanced stage NSCLC for specific molecular aberrations in four genes: *EGFR*,

ALK, *ROS1*, and *BRAF* (Fig. 1). These therapies included erlotinib, gefitinib, afatinib, and osimertinib for *EGFR* mutations, crizotinib, ceritinib, and alectinib for *ALK* fusions, crizotinib for *ROS1* fusions, and the combination treatment of dabrafenib and trametinib in *BRAF*-mutated advanced stage NSCLC.

Over the past five years (2018–2023), the EMA has authorized the use of targeted therapies for four other molecular biomarkers in advanced-stage NSCLC. These include *NTRK* fusions targeted by larotrectinib and entrectinib, *RET* fusions targeted by selpercatinib and pralsetinib, *KRAS* G12C mutations targeted by

Timeline of EMA-approved targeted therapies for patients with advanced stage NSCLC (July 2023)



a) Felip E, Gridelli C, Bass P, et al. Metastatic non-small-cell lung cancer: consensus on pathology and molecular tests, first-line, second-line, and third-line therapy. 1st ESMO Consensus Conference in Lung Cancer; Lugano 2010. *Ann Oncol*. 2011;22(7):1507-1519.
b) Kerr KM, Bubendorf L, Edelman MJ, et al. Second ESMO consensus conference on lung cancer: pathology and molecular biomarkers for non-small-cell lung cancer. *Ann Oncol*. 2014;25(9):1681-1690.
c) Planchard D, Popat S, Kerr K, et al. Metastatic non-small cell lung cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up [published correction appears in *Ann Oncol*. 2019 May;30(5):863-870]. *Ann Oncol*. 2018;29(Suppl 14):iv192-iv237.
d) Metastatic Non-Small-Cell Lung Cancer: ESMO Clinical Practice Guidelines for Diagnosis, Treatment and Follow-Up (2020). <https://www.esmo.org/content/download/347819/6934778/1/ESMO-CPG-mNSCLC-15SEP2020.pdf> (Accessed 20 March 2021).
e) Hendriks LE, Kerr KM, Menis J, et al. Oncogene-addicted metastatic non-small-cell lung cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up [published online ahead of print, 2023 Jan 23]. *Ann Oncol*. 2023;30(23):7534-7542[04781-0].

Fig. 1: Timeline of EMA-approved targeted therapies and the ESMO-recommendations for predictive biomarker testing associated with these targeted drugs for patients with advanced stage NSCLC (as of July 2023).

sotorasib, and *MET* exon 14 skipping mutations targeted by tepotinib and capmatinib. Furthermore, entrectinib was also approved for the treatment of *ROS1* fusion-positive NSCLC, brigatinib and lorlatinib were approved as ALK inhibitors, and the group of *EGFR*-targeting drugs was enriched with the approval of amivantamab for *EGFR* exon 20 insertion mutations. Lastly, the treatment indication of osimertinib was recently extended to include non-metastatic NSCLC patients (see also early-stage NSCLC review).¹²

The following paragraphs provide an overview of the key characteristics and recent developments associated with each targeted therapy, subdivided by their respective biomarker. For a comprehensive overview of all EMA-approved indications for targeted therapy use in patients with NSCLC, including references, see [Supplementary File 2](#).

EGFR-targeted treatment

Activating mutations in *EGFR* have a prevalence of approximately 12–15% among European patients with NSCLC, although they are more common in Asian populations.¹³ These mutations are generally divided into three groups: (1) common *EGFR* mutations (exon 19 deletions and L858R mutations), (2) *EGFR* exon 20 insertion mutations, and (3) uncommon/rare *EGFR* mutations.¹⁴ First- (erlotinib, gefitinib) and second-generation (afatinib) *EGFR* tyrosine kinase inhibitors (TKI) have been available for over a decade and were primarily developed for the treatment of common *EGFR* mutations. Osimertinib, a third-generation *EGFR*-TKI, was initially approved as second-line treatment for patients with the acquired T790M resistance mutation. However, in 2018, the EMA expanded the approval of osimertinib to include first-line treatment of *EGFR* mutation-positive NSCLC. This decision was based on the FLAURA study, which demonstrated improved progression-free survival, overall survival, and treatment of brain metastasis compared to first-generation *EGFR*-TKIs.¹⁵

Patients with *EGFR* exon 20 insertion mutations have generally shown poor response to treatment with first-, second-, and third-generation *EGFR*-TKIs.^{14,16} Newly developed *EGFR*-targeting therapies, including amivantamab and mobocertinib, have shown promising treatment results in patients with *EGFR* exon 20 insertion mutations.^{17–21} In 2021, amivantamab, a bispecific anti-*EGFR* and anti-*MET* antibody, was approved by the EMA as a second-line treatment option for advanced-stage NSCLC patients with *EGFR* exon 20 insertion mutations, based on the results of the phase I CHRYSLIS trial.¹⁹

The group of uncommon/rare *EGFR* mutations encompasses all remaining *EGFR* mutations and the actionability of these mutations may vary. In Dutch patients with advanced stage NSCLC, 18.7% of all *EGFR* mutations detected by multi-gene assays were uncommon/rare *EGFR* mutations, forming a significant

subgroup of *EGFR*-mutated NSCLC.¹⁴ Better clinical outcomes have typically been reported with afatinib compared to osimertinib, in particular in uncommon *EGFR* mutations with relatively high prevalence, such as G719X.^{16,22} While there is limited clinical data comparing the treatment outcome of TKI in rare *EGFR* mutations, pre-clinical studies have demonstrated significant differences in the efficacy of different *EGFR*-TKIs, using codon-based and/or structure-based classifications to explain and predict efficacy.²³

In 2021, the phase III ADAURA trial results led to the approval of osimertinib as an adjuvant therapy following complete tumor resection (with or without adjuvant chemotherapy) in patients with stage IB–IIIA NSCLC harboring either an *EGFR* exon 19 deletion or L858R mutation.²⁴ This marks the first-approved targeted therapy for use in adjuvant setting in NSCLC, opening up possibilities for potential benefits of other targeted therapies in an adjuvant treatment setting.^{12,25}

ALK-targeted treatment

Fusions involving anaplastic lymphoma kinase (*ALK*) are found in approximately 2–5% of patients with NSCLC.¹³ The introduction of crizotinib, a first-generation ALK inhibitor, revolutionized the treatment landscape for ALK fusion-positive NSCLC patients. However, in the past five years, there have been significant advancements with the approval of second-generation (alectinib, ceritinib and brigatinib) and third-generation (lorlatinib) ALK inhibitors by the EMA for their use in patients with ALK fusion-positive metastatic NSCLC. These newly approved ALK inhibitors not only provide additional options for first-line ALK-targeted treatment options, but they also demonstrate improved clinical efficacy in the central-nervous system (CNS) and in cases where certain ALK mutations may arise as resistance mechanisms to ALK inhibitors.²⁶ The development of resistance mutations is a common challenge in targeted therapies, and the emergence of these mutations can limit the effectiveness of initial treatment. However, brigatinib and lorlatinib have shown promising effects in overcoming resistance and maintaining therapeutic response in patients with specific ALK mutations.²⁶

The development of new ALK inhibitors and their sequential use in routine clinical practice has prolonged the lives of patients with ALK fusion-positive NSCLC up to five to seven years.²⁷ The continuous development and approval of next-generation ALK inhibitors highlight the importance of ongoing research and innovation in the field of targeted therapies for NSCLC.²⁸

ROS1-targeted treatment

A phase I trial of crizotinib demonstrated a significant overall response rate (ORR) of 72% (36 out of 50 patients) in individuals with *ROS1*-rearranged NSCLC.²⁹ Subsequently, in 2016, the EMA extended the treatment indication of crizotinib to include patients with

advanced stage *ROS1* fusion-positive NSCLC. An integrated analysis of three ongoing separate trials with entrectinib, a combined pan-TRK, *ROS1* and ALK-inhibitor, demonstrated an ORR of 77% (41 out of 53 patients) with entrectinib.³⁰ Notably, the response rates to entrectinib were similar between patients with and without CNS metastases at baseline, whereas the efficacy of crizotinib may be compromised in these patients due to its limited ability to cross the blood-brain barrier and reduced intracerebral drug activity.³¹ In 2020, entrectinib received simultaneous approval as a treatment option for patients with advanced stage *ROS1* fusion-positive NSCLC, and as pan-tumor therapy in *NTRK* fusion-positive tumors (see Section [NTRK-targeted treatment](#). *NTRK* fusions). However, it is important to note that entrectinib is indicated for patients not previously treated with *ROS1* inhibitors as its effectiveness may be limited to treatment-naïve patients.

BRAF-targeted treatment

Mutations in V-raf murine sarcoma viral oncogene homolog B1 (*BRAF*) occur in approximately 2–4% of patients with NSCLC.^{13,32,33} Among these mutations, missense mutations at codon 600 of *BRAF* (known as V600 mutations) account for around 40% of all *BRAF* mutations in NSCLC. These *BRAF* V600 mutations are classified as class I *BRAF* mutations, as they lead to increased *BRAF* kinase activity that is not dependent on *BRAF* dimerization or activation of *RAS*.³² In 2017, the EMA approved the use of dabrafenib in combination with trametinib for the treatment of advanced NSCLC patients with V600 mutations in *BRAF*.³³ However, it is important to note that there is currently no EMA approval for the use of combined *BRAF* and MEK inhibition in NSCLC with class II (*RAS*-independent dimerization) or class III (*RAS*-dependent dimerization with *CRAF*) *BRAF* driver mutations. These different classes of *BRAF* mutations have distinct molecular characteristics and response profiles to targeted therapies. A recent meta-analysis described that MAPK-targeted therapies, including *BRAF* and MEK inhibitors, have demonstrated clinical activity in some tumors with non-V600 *BRAF* mutations, in particular those with class II mutations.³⁴ However, clinical trials regarding the efficacy of MAPK-targeted therapies in patients with NSCLC with class II or class III *BRAF* mutations are currently lacking. This is primarily due to the heterogeneity of mutations within each class. Certain non-V600 *BRAF* mutations may still be targetable in NSCLC using existing *BRAF* and/or MEK inhibitors, although further research is needed to determine their effectiveness.

NTRK-targeted treatment

Fusions involving the neurotrophic tropomyosin receptor kinase genes (*NTRK1*, *NTRK2*, and *NTRK3*) are known oncogenic drivers in various types of

malignancies, including NSCLC, and have a combined prevalence of <1% in NSCLC.^{13,35} In 2019, the EMA approved larotrectinib, a selective pan-TRK (*TRKA*, *TKRB*, *TRKC*) inhibitor, followed in 2020 by the approval of entrectinib, a combined pan-TRK, *ROS1* and ALK inhibitor, based on two studies with combined results of multiple phase I/II trials of these therapies in solid tumors.^{36,37} A clinical study later showed the efficacy of larotrectinib in previously treated patients with lung cancer harboring an *NTRK1/2/3* fusion. This study demonstrated an ORR of 73% and a median progression-free survival (PFS) of 33.9 months among evaluable patients (11 out of 15 patients).³⁸ Entrectinib showed a comparable ORR of 70% in patients with *NTRK1/2/3* fusion-positive NSCLC (7 out of 10 patients).³⁹ These findings suggest that targeted therapy directed at *NTRK1/2/3* fusion-positive NSCLC can provide significant and durable clinical benefits in metastatic NSCLC, despite representing a small subset of patients.

RET-targeted treatment

Rearranged during transfection (*RET*) fusions have a prevalence of 1–3% in patients with NSCLC.¹³ In 2021, two selective *RET* kinase-inhibitors, selpercatinib and pralsetinib, were approved by the EMA for treatment of patients with advanced stage *RET* fusion-positive NSCLC, including application as first-line therapy. The phase I/II LIBRETTO-001 trial evaluating the efficacy of selpercatinib in previously treated patients with *RET* fusions showed promising results, demonstrating an ORR of 64% (67 out of 105 patients) and a median PFS of 17.5 months.⁴⁰ In previously untreated patients, selpercatinib as first-line systemic therapy resulted in an ORR of 85% (33 out of 39 patients).⁴⁰ Similarly, pralsetinib has demonstrated favorable response rates in patients with *RET* fusion-positive NSCLC. In patients previously treated with platinum-based chemotherapy, pralsetinib treatment resulted in an ORR of 61% (53 out of 87 patients), while treatment naïve patients showed an ORR of 70% (19 out of 27 patients).⁴¹ These results suggest that selpercatinib and pralsetinib are an effective therapeutic option for both previously treated and treatment-naïve NSCLC patients with *RET* fusions. Ongoing clinical trials will further refine the understanding of *RET* fusions and optimize the use of *RET* kinase inhibitors compared to currently standard first-line chemo-immunotherapy in the management of NSCLC.

KRAS^{G12C}-targeted treatment

Somatic mutations in rat sarcoma virus (*RAS*) genes, including *KRAS*, *NRAS*, and *HRAS*, are the most commonly observed oncogenic aberrations in human cancers, occurring in approximately 25% of all cancers.⁴² Among the *RAS* genes, driver mutations in the Kirsten rat sarcoma virus (*KRAS*) gene account for about 85% of

all RAS-driven malignancies. In NSCLC, the prevalence of *KRAS* mutations ranges from 26 to 41%.^{42–45} Directly targeting mutated *KRAS* has posed a significant challenge due to the high affinity of *KRAS* for GDP/GTP and the absence of identified allosteric regulatory binding sites.⁴⁶ However, in 2013, the discovery of the switch-II pocket in *KRAS*^{G12C} provided a breakthrough for the development of clinically effective *KRAS*^{G12C} inhibitors.^{47,48} Sotorasib has shown promising results in pretreated NSCLC patients with an ORR of 37% (46 out of 124 patients) and disease control rate (DCR) of 81% (100 out of 124 patients).⁴⁹ Moreover, in the phase III CodeBreaK 200 trial, sotorasib improved PFS compared to docetaxel treatment (5.6 months versus 4.5 months) with more favorable safety profile, in patients pretreated with chemo-immunotherapy.⁵⁰ As *KRAS* G12C mutations account for more than a third of all *KRAS* mutations in patients with NSCLC (overall prevalence of 12–17%), *KRAS*^{G12C} inhibitors such as sotorasib hold the potential to improve clinical outcome in a significant group of patients with advanced stage NSCLC.^{13,44,45} In 2021, the EMA approved the use of sotorasib for patients with advanced stage *KRAS*^{G12C}-mutated NSCLC, who have received at least one prior line of systemic therapy. The development of *KRAS*^{G12C} inhibitors represents a significant advancement in the field of targeted therapies for NSCLC and offers new hope for improving treatment outcomes in patients with *KRAS*^{G12C}-mutated tumors. Ongoing research and clinical trials will further explore the potential of *KRAS*-targeted therapies and their combination with other treatment modalities in the management of NSCLC.

MET-targeted treatment

Intron 13 and splice-site mutations of mesenchymal-epithelial transition (*MET*) gene result in loss of transcription of exon 14 lead to increased activation of *MET* signaling, due to the loss of the Y1003 binding site of CBL on the juxtamembrane domain of *MET*.⁵¹ These *MET* exon 14 skipping mutations occur in 1–3% of patients with NSCLC.⁵² Unlike crizotinib, a type 1a *MET* inhibitor, capmatinib and tepotinib are selective *MET* kinase domain inhibitors (type 1b *MET* inhibitor).⁵³ In the phase II GEOMETRY mono-1 study, capmatinib has demonstrated response rates of 41% in pretreated patients (28 out of 69 patients) and 68% in treatment-naïve patients (19 out of 28 patients) with NSCLC harboring a *MET* exon 14 skipping mutation.⁵⁴ With simultaneously published results, in the phase II VISION study, tepotinib demonstrated an ORR of 46% in patients with advanced stage NSCLC (46 out of 99 patients), with similar response rates observed in both pretreated and treatment-naïve patients.⁵⁵

Both capmatinib and tepotinib were approved by the EMA in 2021 for the treatment of advanced stage NSCLC patients harboring *MET* exon 14 skipping mutations who require systemic therapy after treatment

with immunotherapy and/or platinum-based chemotherapy. Though capmatinib and tepotinib have demonstrated comparable response rates, the clinical trial evaluating capmatinib reported a higher occurrence of grade III or IV adverse events compared to the tepotinib trial.^{54,55} The safety and tolerability profiles of these drugs should be considered when making treatment decisions for patients with *MET* exon 14 skipping-mutated NSCLC.

European guidelines for predictive biomarker testing

The ESMO Clinical Practice Guidelines for patients with metastatic NSCLC, as of 2018, recommended molecular testing for detection of *EGFR* mutations in exons 18–21, *BRAF* V600 mutations, *ALK* fusions, and *ROS1* fusions (Table 1).⁵⁶ In the 2020 guideline update, *NTRK* fusions were added to the previously recommended biomarkers. In the current ESMO guidelines (2023), this list of recommended predictive biomarkers has been expanded by the addition of *MET* exon 14 skipping mutations, *MET* amplifications, *KRAS*^{G12C} mutations, *RET* fusions, and *HER2/ERBB2* mutations.⁵ Generally it takes one to two years for the ESMO guidelines to include recommendations for molecular biomarker testing after EMA-approval of corresponding targeted therapies (Fig. 1). However, the inclusion of *ERBB2* exon 20 insertions as a recommended biomarker in the current ESMO guidelines is an exception. Although *ERBB2*-targeted therapies such as trastuzumab-deruxtecan and pyrotinib have been approved by the FDA, these targeted therapies have not yet received approval from the EMA.⁵ Nevertheless, in many European countries patients can still be treated with *ERBB2*-targeted treatment in clinical trials (see also Section Availability of targeted therapies).

Future guidelines are likely to incorporate additional biomarkers that have predictive and/or prognostic value for targeted therapies (e.g., *TP53* mutations) and/or for non-targeted therapies (e.g., tumor-mutation burden, *STK11* mutations, *KEAP1* mutations). The ongoing development of novel targeted therapies as well as availability of tumor-agnostic therapies already approved in other cancers, and the necessary expansion of biomarkers associated with these novel therapies will continue to shape the landscape of predictive biomarker testing in advanced-stage NSCLC. For more detailed information and a future perspective of predictive biomarker testing in advanced-stage NSCLC, the corresponding viewpoint in this Clinical Series should be read.¹¹

Availability of targeted therapies

Using a questionnaire sent to all participants (see Supplemental File 1), we collected present-day information on the availability of targeted therapies in eleven European countries (summarized in Fig. 2; Table 1).

Indication	Therapy	Austria	Belgium	Czech Republic	England	France	Germany	Netherlands	Portugal	Slovenia	Spain	Sweden
EGFR	gefitinib	●	●	●	●	●	●	●	●	●	●	●
	erlotinib	●	●	●	●	●	●	●	●	●	●	●
	afatinib	●	●	●	●	●	●	●	●	●	●	●
	dacomitinib	●	●	●	●	●	●	●	●	● [#]	●	●
	osimertinib	●	●	●	●	●	●	●	●	● [#]	●	●
	amivantamab	●	● CUP/ [†]	● [†]	● [†]	●	● nfc	● nfc	● CUP/ [†]	●	●	●
	mobocertinib*	● CUP/ [†]	●	●	●	● CUP	●	●	●	● CUP	●	●
KRAS	sotorasib	●	● CUP/ [†]	● [§]	●	●	●	●	● CUP/ [†]	●	●	●
	adagrasib*	● [†]	● CUP/ [†]	●	● [†]	●	●	● CUP	● [†]	●	●	●
BRAF	dabrafenib+trametinib	●	●	● [§]	●	●	●	●	● off-label	● nfc,&,#	●	●
	vemurafenib*	●	●	● [§]	●	●	●	● off-label	● off-label	●	●	● off-label
MET	crizotinib*	●	● CUP	● [§]	●	●	● off-label	● CUP	●	● off-label,&,#	●	●
	capmatinib	●	●	●	●	●	●	●	● CUP	● CUP	●	● nfc
	tepotinib	●	●	● [§]	●	●	●	● nfc	● CUP	●	●	●
HER2/ERBB2	trastuzumab deruxtecan*	● [†]	● [†]	● [§]	●	● [†]	● off-label	● [†]	● off-label	● off-label,&,#	●	● off-label
	pyrotinib*	●	●	●	●	●	●	●	●	●	●	●
ALK	crizotinib	●	●	●	●	●	●	●	●	●	●	●
	alectinib	●	●	●	●	●	●	●	●	●	●	●
	ceritinib	●	●	●	●	●	●	●	●	●	●	●
	brigatinib	●	●	●	●	●	●	●	●	●	●	●
	lorlatinib	●	●	●	●	●	●	●	●	●	●	●
ROS1	crizotinib	●	●	●	●	●	●	●	●	● nfc,&,#	●	●
	lorlatinib*	●	●	● [§]	● CUP	●	● off-label	● nfc	● off-label	● off-label,&,#	●	●
	entrectinib	●	●	● [§]	●	●	●	●	● CUP	●	●	●
NTRK	repotrectinib*	● CUP	● [†]	●	● [†]	● CUP	● [†]	● CUP	● CUP	●	●	●
	larotrectinib	●	●	● [§]	●	●	●	● nfc	● CUP	●	●	●
	entrectinib	●	●	● [§]	●	●	●	● nfc	●	●	●	●
RET	cabozantinib*	●	●	●	●	●	● off-label	● CUP	● off-label	● off-label,&,#	●	● off-label
	vandetanib*	●	●	●	●	●	● off-label	●	● off-label	● off-label,&,#	●	● off-label
	selpercatinib	●	●	●	●	● [†]	●	● nfc	● CUP	●	●	●
	pralsetinib	●	●	● [§]	● [†]	●	●	● nfc	● CUP	●	●	●

●, available as standard-of-care treatment and covered by health insurance or reimbursed fully by government; ● (nfc), available as standard-of-care treatment, but not (fully) covered by health insurance nor reimbursed fully by government; ●, not available as standard-of-care treatment, but available in other setting (e.g., compassionate use); ●[†], only available in trial setting; ●, not available as standard-of-care treatment or in other setting. CUP, compassionate use program (used in table as overarching term for compassionate use program, patient named program, patient access program, and managed access program); n/a, data not available. [†]Trial or CUP if ineligible; [§]Can be requested by treating oncologist to be reimbursed by the state, but not all patients receive reimbursement; [#]Individual patient request to national health insurance company to get reimbursement; [#]Molecular Tumor Board recommendation required; *Treatment indication in advanced stage NSCLC only approved by FDA.

Table 1: Availability of targeted therapies (EMA- and/or FDA-approved) for advanced stage NSCLC in a selection of European countries.

Any targeted therapy with a treatment indication approval for patients with advanced stage NSCLC by either the EMA and/or the FDA, were included in the questionnaire to form a comprehensive overview of available targeted treatments. Overall, not all EMA-approved therapies are available in every participating European country. First-, second-, and third-generation EGFR TKIs, first-, second- and third-generation ALK inhibitors, and ROS1 inhibitors are available as standard-of-care treatment in all European countries participating in this study (Fig. 2; Table 1). Amivantamab is only available as standard-of-care treatment in

four out of eleven countries. Targeted therapy as standard-of-care treatment is not available in all countries for patients with advanced stage NSCLC with an NTRK fusion (available as standard-of-care in seven out of eleven countries), RET fusion (8/11 countries), BRAF V600 mutation (9/11 countries), MET exon 14 skipping mutation (6/11 countries), or KRAS G12C mutation (5/11 countries). It should be noted that in some countries the only available MET-targeted treatment is crizotinib, which is FDA-approved but not EMA-approved for its application in patients with advanced stage NSCLC harboring an MET exon 14 skipping mutation.




Fig. 2: Availability of targeted therapies for patients with metastatic NSCLC across eleven European countries. For each country, the displayed information was provided by a detailed questionnaire (Supplementary File 1) that was completed by a (molecular) pathologist, clinical scientist in molecular pathology and/or (pulmonary) oncologist with expertise in the field of NSCLC. *Presented availability concerns crizotinib (FDA-approved), due to its better availability compared to tepotinib and capmatinib (EMA-approved) (see also Table 2).

National guidelines for predictive biomarker testing

The representatives of all eleven participating European countries reported current national guidelines with recommendations regarding predictive biomarkers testing in patients with (advanced stage) NSCLC. Table 2 summarizes the specific details on the recommendations of each participating country regarding biomarker testing as reported in the national guidelines of these countries. The interpretation of what constitutes valid, applicable national guidelines is not always clear-cut. For example, England does not have tumor-specific, detailed guidelines on molecular testing requirements, and Portugal has limited national guidelines supplemented by national expert consensus recommendations.^{57–59} National guidelines largely align with the biomarkers listed in the ESMO guidelines including *EGFR*, *ALK*, and *ROS1*, which were among the first biomarkers for which targeted therapies were available. Testing for these biomarkers is recommended in all participating countries.^{57–68} For the other biomarkers, however, the representation of testing recommendations in national guidelines varies, and the introduction of these biomarkers into the guidelines

has occurred over a broader range of time compared to *EGFR*, *ALK*, and *ROS1*. As other biomarkers each belong to more recently EMA-approved targeted therapies (see Fig. 1), a plausible explanation is that these differences are caused by delays in the updating of national guidelines.

Some countries have included *NRG1* testing as either mandatory (the Netherlands) or as recommendation (Austria, France) in their national guidelines. This inclusion is due to the relatively poor prognosis of patients with *NRG1* fusion-positive NSCLC, the lack of response to treatment with either chemotherapy or immunotherapy, and the potential inclusion of these patients in ongoing clinical trials (NCT02912949).⁶⁹

Predictive biomarker testing rates

Real-world data of predictive biomarker testing rates in patients with NSCLC is limited, and the available information is often outdated or grouped with older data (Table 3).^{7,44,70,72,76} However, recent studies from England, Spain, the Netherlands, and Norway have demonstrated high testing rates for *EGFR* and *ALK*.^{71,74,75,77} It should be noted that testing rates of

Biomarker	Austria	Belgium	Czech Republic	England ^a	France	Germany	Netherlands	Portugal	Slovenia	Spain	Sweden
<i>EGFR</i>	● 2013	● 2010	● 2012	● n/a	● 2008	● 2012	● 2011	● 2013	● 2010	● 2012	● 2009
<i>ALK</i>	● 2013	● 2013	● 2013	● n/a	● 2012	● 2012	● 2015	● 2013	● 2013	● 2012	● 2013
<i>ROS1</i>	● 2013	● 2017	● 2016	● n/a	● 2012	● 2015	● 2015	●* 2022	● 2015	● 2020	● 2014
<i>BRAF V600</i>	● 2020	● 2017	●*	● n/a	● 2012	● 2016	● 2015	●* 2022	● 2018	● 2020	● 2018
<i>RET</i>	● 2020	● 2022	●*	● n/a	● 2018	● 2016	● 2015	●* 2022	● 2022	● 2023	● 2023
<i>MET</i> exon 14	● 2020	● 2020	●*	● n/a	● 2012	● 2018	● 2020	●* 2022	● 2022	● 2023	● 2023
<i>MET</i> amp	● 2020	● 2023	●	● n/a	●* 2023	●* 2018	●	●* 2022	●	● 2023	●* 2023
<i>KRAS G12C</i>	● 2020	● 2021	●*	● n/a	● 2008	● 2021	● 2015	●* 2022	● 2022	● 2023	● 2022
<i>NTRK</i>	● 2020	● 2021	●*	● n/a	● 2018	● 2016	● 2020	●* 2022	● 2020	● 2023	● 2022
<i>HER2/ERBB2</i>	● 2020	● 2016	●*	●	● 2022	● 2016	● 2015	●* 2022	● 2022	● 2023	●* 2018
<i>NRG1</i>	● 2020	●	●	●	● 2022	●* 2019	● 2020	●	●	●	●* 2019
<i>PD-L1</i>	● 2020	● 2018	● 2016	● n/a	● 2015	● 2016	● 2020	●	● 2017	● 2020	● 2015

●, required by current national guidelines; ●*, recommended by current national guidelines; ●*, not specified in current national guidelines but recommended by national expert consensus; ●, testing not recommended or biomarker not present in current national guidelines.

^aGuidelines state that presence of *MET* amplification can/is recommended to be tested after progression on EGFR TKI. ^bBased on national genomic test directory of the NHS England are used, due to lack of national, tumor-specific guidelines on molecular testing requirements.

^cGuidelines state that *MET* amplification and *NRG1* fusion testing can be performed, but no recommendation regarding their application. n/a, the year of introduction is not available. ^dGuidelines previously (2016) recommended *MET* amplification testing only in the setting of EGFR TKI-resistance.

Table 2: Biomarker recommendations and year of introduction in national guidelines of European countries for molecular testing of patients with metastatic NSCLC.

predictive biomarkers introduced after *EGFR* and *ALK* tend to be lower, but should be interpreted with caution, due to the often limited reporting of these biomarkers and potential variations in testing methodologies over time. Observational studies have also highlighted substantial variation in molecular testing practices for NSCLC between institutions within a country, both in European countries and the United States.^{78–80} This suggests that access to predictive biomarker testing may be influenced by factors such as the location of the hospital where patients receive their diagnosis and treatment, accessibility of technology for comprehensive biomarker testing, reimbursement issues and/or lack of knowledge. In some countries, efforts have been made to actively improve access and overall functioning of molecular testing for NSCLC patients through dedicated projects and policies.^{81–84} With regard to England, it should be noted that a comprehensive overview of national predictive biomarker testing rates is currently being established by the National Disease Registration Service. In the Netherlands, molecular testing results are registered nationwide in the Dutch Pathology Registry (Palga), which enables researchers to examine these reports of patients diagnosed with NSCLC within a specified time period. Using this registry, reliable national testing rates of *EGFR*, *ALK*, *ROS*, *MET*, *RET*, *BRAF*, and *ERBB2* in 2017, and *KRAS* in 2013–2017 have previously been determined (see Table 3).^{44,74} Importantly, for other countries, national testing rates are not yet available and (publicly) available testing rates are limited to regional or multi-institutional patient populations, thereby potentially limiting their representativeness for the entire country.

Interpretation of molecular test results

Among the participating centers, there is a significant overlap in the elements provided in molecular pathology reports for NSCLC. These include information on the genes tested, the specific variants detected, variant allele frequency, and tumor cell percentage (Table 4). However, differences arise in the translation of test outcome to report conclusion, and the interpretation of reports by healthcare professionals for patient care. While most participating centers provide general treatment advice based on test results, only a few centers offer mutation-specific recommendations. International guidelines on reporting molecular results recommend a general treatment advice in cases with pathogenic, actionable molecular aberrations.⁸⁵ However, in some countries, national guidelines state that such treatment recommendations are not allowed to be stated in the pathology report.

We also inquired among the participating centers what methodology was used for discussing and advising on complex molecular results of patients with metastatic NSCLC (Table 5). The establishment of dedicated Molecular Tumor Boards (MTBs) for the interpretation of complex molecular testing results is common among the centers. However, only a few countries, including Germany and the Netherlands, have national guidelines in place that outline the composition and functioning of these treatment advisory boards.^{65,86,87} The variation in the translation and interpretation of complex or rare molecular testing results highlights the need for both to use and report following existing guidelines as well as further standardized guidelines and recommendations to ensure a consistent and accurate understanding of

	Austria ⁷⁰	Belgium ^a	Czech Republic ^c	England ⁷¹	France ⁷²	Germany ⁷³	Netherlands ^{44,74}	Portugal	Slovenia ⁶¹	Spain ⁷⁵	Sweden ^d
EGFR	77.1%	51%	79.3%	92%	53%	72.5%	~60–82% ^b	No data	91%	91.4%	No data (authors estimate: ~90%)
Year(s) of data	(2013–2015)	(2011)	(2011)	(2017)	(2015–2018)	(2015–2019)	(2017)		(Single center, 2020)	(2018–2019)	
ALK	62.5%	No data	82.7%	80%	46%	74.5%	~35%–55% ^{b,e}	No data	87%	80.1%	No data (authors estimate: ~90%)
Year(s) of data	(2013–2015)		(2013)	(2017)	(2015–2018)	(2015–2019)	(2017)		(Single center, 2018)	(2018–2019)	
ROS1	No data	No data	82.7%	No data	34%	66.1%	~28–38% ^{b,f}	No data	86%	56.2%	No data (authors estimate: ~90%)
Year(s) of data			(2015)		(Mutation and/or fusion, 2015–2018)	(2015–2019)	(2017, total)		(Single center, 2018)	(2018–2019)	
BRAF	No data	No data	No data	No data	38%	53.0%	~60–78% ^b	No data	No data	No data	No data (authors estimate: ~70%)
Year(s) of data					(2015–2018)	(2015–2019)	(2017)				
RET	No data	No data	No data	No data	No data	26.9%	~18–19% ^{b,g}	No data	No data	No data	No data (authors estimate: ~70%)
Year(s) of data						(2015–2019)	(2017, in total)				
MET mut.	No data	No data	No data	No data	12%	35.4%	~55–68% ^b	No data	No data	No data	No data (authors estimate: ~70%)
Year(s) of data					(Mutation and/or fusion, 2015–2018)	(2015–2019)	(2017)				
MET amp.	No data	No data	No data	No data	See 'MET mut.'	No data	No data	No data	No data	No data	No data
Year(s) of data											
KRAS	No data	No data	No data	No data	45%	44.9%	82.0%	No data	No data	No data	No data (authors estimate: ~70%)
Year(s) of data					(2015–2018)	(2015–2019)	(2017)				
NTRK	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data (authors estimate: ~50%)
Year(s) of data											
HER2/ERBB2	No data	No data	No data	No data	30%	15.2%	~55–76% ^b	No data	No data	No data	No data (authors estimate: ~50%)
Year(s) of data					(2015–2018)	(2015–2019)	(2017)				
NRG1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data (authors estimate: ~20%)
Year(s) of data											
PD-L1	No data	No data	89.5%	87%	17%	66.2%	No data	No data	74–91%	58.1%	No data (authors estimate: ~90%)
Year(s) of data			(2016)	(2017)	(2015–2018)	(2015–2019)			(Single center, 2018)	(2019–2020)	

^aRecent data not publicly available, but could be requested at RIZIV and Belgian Cancer registration. ^bHigher rate is applicable to adenocarcinoma, lower rate is applicable to NSCLC-NOS. ^cNational testing rates from national cancer registry, calculated from cases eligible for molecular testing (e.g., molecular testing for NSCLC NOS and adenocarcinoma, PD-L1 for all NSCLC). ^dNo public data available on testing rates, displayed numbers are the Swedish authors' estimations based on their expected national adherence to guidelines, personal communication of authors, degree of sequential testing. ^e~70–80% if EGFR/KRAS/BRAF/ERBB2/MET wildtype. ^f~60–70% if EGFR/KRAS/BRAF/ERBB2/MET wildtype. ^g~42–44% if EGFR/KRAS/BRAF/ERBB2/MET wildtype.

Table 3: Published molecular testing rates for patients with metastatic NSCLC across participating European countries.

Topic	Austria [§]	Belgium	Czech Republic	England	France	Germany	Netherlands	Portugal	Slovenia	Spain	Sweden
Affected gene (e.g., <i>EGFR</i>)	●	●	●	●	●	●	●	●	●	●	●
Mutation group (e.g., exon 19 deletion)	●	●	●	●	●	●	●	●	●	●	●
Specific mutation (e.g. E746_A750del)	●	●	●	●	●	●	●	●	●	●	●
Variant allele frequency of mutation(s)	●	●	●	●	●	●	●	●	●	●	●
Tumor cell percentage	●	●	●	●	●	●	●	●	●	●	●
Generic treatment advice (related to affected gene or type of mutation)	●	●	●	●	●	●	●	●	●	●	●
Specific treatment advice (tailored to mutation type or combination of multiple mutations)	●	●*	●	●	●	●**	●***	●	●	●	●

●, applicable at participating center; ●, not applicable at participating center, [§]Pathogenicity of mutation is also provided, and it is only whether or not there currently is therapeutic relevance. *Treatment advice is most often generic, but specific treatment advice may be given in certain circumstances (e.g., in the context of resistance). **Other information includes short description of the method including thresholds, regions that dropped out of analysis, coverage at mutation localization, biological interpretation of variants (e.g. activating, loss of function), information of clinical studies, if available. ***Reports state the possibility to discuss molecular testing results in molecular tumor board to receive specific treatment advice.

Table 4: Reporting of molecular testing results for patients with metastatic NSCLC across participating centers (of clinical scientist in molecular pathology/(molecular) pathologist).

complex/rare molecular testing results among health-care professionals.⁸⁵ These guidelines should help to guide appropriate treatment decision-making and patient care optimization in the context of NSCLC.

Barriers for off-label use, compassionate use programs and trial participation

As stated in Section [Availability of targeted therapies](#), the availability of targeted therapies for patients with

Statement	Austria	Belgium	Czech Republic	England	France	Germany	Netherlands	Portugal	Slovenia	Spain	Sweden
Specific treatment advice is given in the report	●	●	●	●	●	●	●	●	●	●	●***
There are standard meetings for discussion of complex molecular testing results	●	●	●	●	●	●	●	●	●	●	●
Cases are discussed in regular (multidisciplinary) meetings for oncological care	●	●	●	●	●	●	●	●	●	●	●
Cases can be discussed in a dedicated MTB	●	●	●	●	●	●	●	●	●	●	●
There are national guidelines regarding the compositions of MTBs	●	●	●	●	●	●*	●**	●	●	●	●

●, applicable at participating center; ●, not applicable at participating center. *Experts that have to participate include medical oncologist(s), pulmonary or thoracic oncologist(s), molecular biologist(s) in pathology/clinical scientist(s) in molecular pathology, and clinical geneticist(s). **For all pulmonary oncology, participants include pulmonary oncologist(s), pulmonary pathologist(s), clinical scientist(s) in molecular pathology, and upon indication participants also include clinical geneticist(s), structural biologist, medical oncologist(s), bioinformatician(s), and pharmacist(s). ***No specific recommendations, but links to case reports and exploratory studies.

Table 5: Methodology of discussing complex molecular results of patients with metastatic NSCLC across participating centers (of clinical scientist in molecular pathology/(molecular) pathologist).

Biomarker	Test type	Austria	Belgium	Czech Republic	England	France	Germany	Netherlands	Portugal	Slovenia	Spain	Sweden
<i>EGFR</i> mutations	NGS	●	●	●	●	●	●	●	●	●	●	●
	Sanger	●	●	●	●	●	●	●	●	●	●	●
	HRM	●	●	●	●	●	●	●	●	●	●	●
	Cobas	●	●	●	●	●	●	●	●	●	●	●
	ddPCR	●	●	●	●	●	●	●*	●	●	●	●
	Idylla	●	●	●	●	●	●	●*	●	●	●	●
<i>KRAS</i> G12C mutations	Other	●	●	AmoyDx	●	●	●	●	●	●	●	●
	NGS	●	●	●	●	●	●	●	●	●	●	●
	Sanger	●	●	●	●	●	●	●	●	●	●	●
	HRM	●	●	●	●	●	●	●	●	●	●	●
	Cobas	●	●	●	●	●	●	●	●	●	●	●
	ddPCR	●	●	●	●	●	●	●*	●	●	●	●
<i>BRAF</i> V600 mutations	Idylla	●	●	●	●	●	●	●*	●	●	●	●
	NGS-DNA	●	●	●	●	●	●	●	●	●	●	●
	Sanger	●	●	●	●	●	●	●	●	●	●	●
	HRM	●	●	●	●	●	●	●	●	●	●	●
	Cobas	●	●	●	●	●	●	●	●	●	●	●
	RNA seq.	●	●	●	●	●	●	●	●	●	●	●
<i>MET</i> exon 14 skipping mutations	NGS-RNA	●	●	●	●	●	●	●	●	●	●	●
	ddPCR	●	●	●	●	●	●	●*	●	●	●	●
	Idylla	●	●	●	●	●	●	●*	●	●	●	●
	IHC (V600E)	●	●	●	●	●	●	●	●	●	●	●
	NGS-DNA	●	●	●	●	●	●	●	●	●	●	●
	Sanger	●	●	●	●	●	●	●	●	●	●	●
<i>MET</i> amp.	RNA seq.	●	●	●	●	●	●	●	●	●	●	●
	NGS-RNA	●	●	●	●	●	●	●	●	●	●	●
	ddPCR	●	●	●	●	●	●	●	●	●	●	●
<i>ERBB2</i> mutations	Idylla	●	●	●	●	●	●	●	●	●	●	●
	FISH	●	●*	●	●	●	●	●*	●	●	●	●
	NGS-DNA	●	●	●	●	●	●	●	●	●	●	●
<i>ALK</i> fusions	NGS	●	●	●	●	●	●	●	●	●	●	●
	Sanger	●	●	●	●	●	●	●	●	●	●	●
	HRM	●	●	●	●	●	●	●	●	●	●	●
	Cobas	●	●	●	●	●	●	●	●	●	●	●
<i>ROS1</i> fusions	IHC	●	●	●	●	●	●	●	●	●	●	●
	RNA seq.	●	●	●	●	●	●	●	●	●	●	●
	FISH	●	●	●	●	●	●	●*	●	●	●	●
	NGS-RNA	●	●	●	●	●	●	●	●	●	●	●
<i>RET</i> fusions	IHC	●	●	●	●	●	●	●	●	●	●	●
	RNA seq.	●	●	●	●	●	●	●	●	●	●	●
	FISH	●	●	●	●	●	●	●*	●	●	●	●
	NGS-RNA	●	●	●	●	●	●	●	●	●	●	●
	Other	●	●	●	●	●	●	●	●	●	●	●
	Idylla	●	●	●	●	●	●	●	●	●	●	●
<i>NTRK</i> fusions	Other	●	●	●	AmoyDx	●	●	●	●	●	●	●
	IHC	●	●	●	●	●	●	●*	●	●	●	●
	RNA seq.	●	●	●	●	●	●	●	●	●	●	●
	FISH	●	●	●	●	●	●	●	●	●	●	●
	NGS-RNA	●	●	●	●	●	●	●	●	●	●	●
<i>NRG1</i> fusions	Other	●	●	●	●	●	●	●	●	●	●	●
	IHC	●	●	●	●	●	●	●*	●	●	●	●
In what order are above biomarkers routinely tested?	NGS-RNA	●	●	●	●	●	●	●	●	●	●	●
	Sim.	Sim.	Sim.	Sim.	Sim.	Sim.	Seq.**	Sim.	Sim.	Sim.	Sim.	Sim.

●, performed at participating center; ●, not performed at participating center. *Tests only performed in specific circumstances (e.g., confirmation of difficult NGS data, insufficient quality for NGS, or amount of tissue/DNA/tumor cell percentage is too low for routine NGS and re-biopsy is not possible). **PD-L1 IHC, DNA-NGS, and Fast track using the PGX system of Diatech is started simultaneously; RNA-based fusion analysis is started in wildtype samples. Sim., simultaneously; seq. sequentially.

Table 6: Available biomarker test types for patients with metastatic NSCLC in participating centers (of clinical scientist in molecular pathology/(molecular) pathologist).

NSCLC varies among European countries. Some targeted therapies have not (yet) received approval from the EMA, or they may not be registered or covered by national health insurance systems. In cases where targeted therapy is not standardly available, there are alternative options for patients to access these treatments. These options include participation in clinical trials, off-label targeted therapy use, and compassionate use or patient-named programs. Off-label use refers to the use of a drug for a purpose not (yet) specifically approved by regulatory authorities. Compassionate use programs allow patients with serious or life-threatening conditions to access experimental therapies outside of clinical trials.

Though these alternative options for targeted drug access are valuable for patients who would otherwise not receive (potentially) effective targeted therapy treatment, various authors of this review have encountered issues when attempting to utilize these opportunities for their patients. These issues include the lack of up-to-date and user-friendly databases for clinical trials and compassionate use programs, limited access to ongoing (international) clinical studies, challenges related to ineligibility criteria such as brain metastases, and compulsory individual negotiations with health insurance providers for off-label use, which may vary depending of the treating center. In some cases, university hospitals may have better chances of obtaining approval for off-label use due to their expertise and resources. Overall, navigating access to targeted therapies outside of standard availability can be complex and may require close collaboration between healthcare providers, patients, and regulatory bodies to ensure the best possible available treatment options for individual patients. In addition, we should aim for equal access to all presently available innovative systemic treatments (i.e., targeted therapies, immune checkpoints inhibitors, antibody-drug conjugates) for all patients with metastatic NSCLC.

Conclusions

The availability of targeted therapies for advanced stage NSCLC varies across European countries, with high availability of EGFR- (except amivantamab), ALK-, ROS1-, and BRAF V600-targeted therapies. However, there is considerable variability in the availability of other targeted therapies, and not all national guidelines align with European guidelines or reflect the availability of specific targeted therapies in each country. Recent real-world data studies in Norway, Spain, the Netherlands, and England have reported encouraging *EGFR* testing rates, though recent data are not publicly available for other participating countries. Data on testing rates for other predictive markers are often either outdated or not widely reported. Therefore, it remains uncertain what proportion of European patients with advanced stage NSCLC receive adequate molecular

Search strategy and selection criteria

References for this Review were identified through EMA European public assessments reports (EPAR) of EMA-authorized targeted therapies for non-small cell lung cancer (see [Supplementary File 2](#) for references and last date of access), subsequent scrutinization of references of included articles and through searches of the authors' own files. References of national guidelines and (national) biomarker testing rates were provided by participants from each country, if available. There was no selection based on language or date of publication.

testing. All centers contributing to this review have the capacity to perform DNA- and RNA-based NGS for recommended biomarkers ([Table 6](#)), and nearly all have established MTBs to discuss complex/rare molecular testing results. Molecular test reports of participating centers were largely overlapping in their content, with the exception of the comprehensiveness of treatment recommendations. While providing an overview of the characteristics of this select group of high-expertise, often university-affiliated centers, the overall real-world implementation of (large-panel) NGS and use of MTB may be suboptimal in other laboratories performing molecular testing within these European countries, as indicated in recent literature.^{88,89} Importantly, there is still a lack of international guidelines that define the role, criteria for patient enrollment, and composition of MTBs. Developing such consensus guidelines will provide a framework for interpretation of rare or complex mutations, guide appropriate treatment-decision making, and ensure quality control within MTBs. A future perspective for molecular testing in advanced stage NSCLC is described in the separate viewpoint paper.¹¹

Contributors

VdJ contributed to the conceptualization, data curation, methodology, visualization, writing—original draft, writing—review & editing; WT contributed to writing—original draft, writing—review & editing; AB, JB, LB, RB, GF, LH, MH, PH, AJ, MJ, Lvk, IK, FLR, ML, JCM, KM, LPA, AR, PT, and JW contributed to the investigation/data acquisition, writing—review & editing; ES and AvdW contributed to the conceptualization, methodology, supervision, writing—original draft, writing—review & editing.

Editor note

The Lancet Group takes a neutral position with respect to territorial claims in published maps and institutional affiliations.

Declaration of interests

WT has received consulting fees from Merck Sharp and Dohme, Bristol-Myers Squibb, and Altana (fees to institution), is board member of Dutch Society of Pathology and member of Council for Research and Innovation of the Federation of Medical Specialists (FMS); AB has received consulting fees from Sanofi (OncoCollective advisory board), payments or honoraria from Roche (oral presentation), support for attending ASCO 2023 from Pfizer; JB has received research grants from Amgen and Bristol-Myers Squibb, lecture honoraria from AstraZeneca, Merck Sharp and Dohme, Roche, Pfizer, Bristol-Myers Squibb, Boehringer Ingelheim, Novartis, GSK, Eli Lilly, Amgen, and Sanofi, and support for attending meetings and/or travel from Amgen; LB has received grants or contract from Takeda, Roche, AstraZeneca, and Bristol-Myers Squibb, payments or honoraria from Invitae, Eli Lilly,

AstraZeneca, Roche, Merck Sharp and Dohme, Merck, Bristol-Myers Squibb, Pfizer, Novartis, Takeda, and Janssen, has participated in data safety monitoring boards or advisory boards for Invitae, Eli Lilly, AstraZeneca, Roche, Merck Sharp and Dohme, Merck, Bristol-Myers Squibb, Pfizer, Novartis, Takeda, and Janssen, is Int. Secretary of the Austrian Society of Pathology, PPS Membership and Awards Committee, and is Member of the Mesothelioma Committee of IASLC; RB has received payments or honoraria for lectures and advisory boards for Abbvie, Amgen, AstraZeneca, Bayer, Bristol-Myers Squibb, Boehringer-Ingelheim, Illumina, Janssen, Lilly, Merck-Serono, Merck Sharp and Dohme, Novartis, Qiagen, Pfizer, Roche, and Targos MP Inc., has received payments for expert testimony from Merck Sharp and Dohme, is co-founder and co-owner of Gnothis Inc. (SE), Timer Therapeutics Inc. (GE); GF has received fees for advisory boards participation, meetings, or as invited speaker, from AstraZeneca, Boehringer Ingelheim, Bristol Myers Squibb, Lilly, Janssen, Merck Sharp and Dohme, Pfizer, Roche Farmacêutica Química Lda, and Takeda; MH has received payments or honoraria from Merck Sharp and Dohme, Roche, Lilly, AstraZeneca, and Takeda; PH has received consulting fees from AstraZeneca, Roche, Abbvie, Qiagen, and Pfizer, has received payments or honoraria from Thermofisher Scientist, Novartis, Janssen, AstraZeneca, and Biocartis, has received support for attending meetings and/or travel from Thermofisher Scientist, Novartis, Janssen, AstraZeneca, and Biocartis, has participated on a Data Safety Monitoring Board or Advisory Board for Thermofisher Scientist, Novartis, Janssen, AstraZeneca, Biocartis, Bristol-Myers Squibb, Sanofi, Roche, and Abbvie; LvK has received institutional grants or contract from Amgen, AstraZeneca, Bayer, Janssen-Cilag, Merck, Roche, and Servier, has received institutional payments or honoraria from Amgen, AstraZeneca, Bayer, Bristol-Myers Squibb, Eli Lilly, Novartis, Pfizer, and Roche, has received institutional support for attending meeting and/or travel from Roche, has participated on a Data Safety Monitoring Board or Advisory Board from Janssen-Cilag, Merck, and Roche, has a leadership or fiduciary role in the Commission Personalized Medicine—Belgium (unpaid), has stock or stock options in Cyclomics (institutional/personal); JCM has received payments or honoraria from Lilly Portugal, Novartis Portugal, Roche Portugal, Fresco Produções, Janssen-Cilag Portugal, Pierre Fabre Portugal, and Servier Portugal; has participated on a Data Safety Monitoring Board or Advisory Board of Roche Portugal; KM has received payments or honoraria from Takeda, Janssen, Pfizer, Merck Sharp and Dohme, Bristol-Myers Squibb, Roche, Amgen, Novartis, and Eli Lilly, has participated on a Data Safety Monitoring Board or Advisory Board of BMS, Takeda, AstraZeneca, Amgen, Roche, Boehringer Ingelheim, and Janssen; SP has received consulting fees (personal fees) from Amgen, AstraZeneca, Bayer, Blueprint, Bristol-Myers Squibb, Boehringer Ingelheim, Daiichi Sankyo, GSK, Guardant Health, Incyte, Janssen, Lilly, Merck Serono, Merck Sharp and Dohme, Novartis, Roche, Takeda, Pfizer, Seattle Genetics, Turning Point Therapeutics, and EQRx, has received payments or honoraria (personal fees) from AstraZeneca, Bayer, Guardant Health, Janssen, Merck Serono, Roche, and Takeda, has received payments for expert testimony (personal fees) from Roche, and Merck Serono, has received reimbursement of travel expenses from Janssen, and Roche, has participated on a Data Safety Monitoring Board or Advisory Board as per consulting fees, has a leadership role or fiduciary role (all unpaid) in the British Thoracic Oncology Group, ALK Positive UK, Lung Cancer Europe, Ruth Strauss Foundation, Mesothelioma Applied Research Foundation, and ETOP-IBCSG Partners Foundation Board; AR has received consulting fees for participation on advisory boards of AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Amgen, Merck Sharp and Dohme, Gilead, and Sanofi, has received honoraria for lectures and presentations from AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Amgen, Merck Sharp and Dohme, Gilead, Roche, Merck-Serono, has received support for attending meetings and travel from Sanofi, Gilead, Roche, and Bristol-Myers Squibb; JW has received payments or honoraria, has received support for attending meetings and travel, and has participated in a Data Safety Monitoring Board or Advisory Board, from Amgen, AstraZeneca, Bayer, Blueprint, BMS, Boehringer-Ingelheim, Chugai, Daiichi Sankyo, Janssen, Lilly, Loxo, Merck, Mirati, MSD, Novartis, Nuvalent,

Pfizer, Pierre-Fabre, Roche, Seattle Genetics, Takeda, and Turning Point; ES has received unrestricted grants (all paid to UMCG institution) from Abbott, Biocartis, AstraZeneca, Invitae/Archer, Bayer, Bio-Rad, Roche, Agena Bioscience, CC Diagnostics, MSD/MERCK, and Boehringer Ingelheim, has received consulting fees (all paid to UMCG institution) from MSD/Merck, AstraZeneca, Roche, Novartis, Bayer, BMS, Lilly, Amgen, Illumina, Agena Bioscience, CC Diagnostics, Janssen Cilag (Johnson & Johnson), Astellas Pharma, GSK, Sinnovisionlab, and Sysmex, has received payments or honoraria (all paid to UMCG institution) from Bio-Rad, Seracare, Roche, Biocartis, Lilly, Agena Bioscience, and Illumina, has received support for attending meetings and/or travel from BioRad, Biocartis, Agena Sciences, and Illumina, is a board member for the Dutch Society of Pathology (unpaid), European Society of Pathology (unpaid), European Liquid Biopsy Society (unpaid), is a secretary/member of the advisory committee for assessment of molecular diagnostics (cieBOD) (honoraria paid to UMCG institution), is committee member of national guideline advisory (honoraria paid to UMCG institution); AvdW, has received grants or contracts from AstraZeneca, Boehringer-Ingelheim, Pfizer, Roche, and Takeda, has received consulting fees from AstraZeneca, Janssen, Lilly, Roche, and Takeda, has received payments or honoraria from AstraZeneca, BMS, Lilly, Pfizer, and Roche, has a leadership or fiduciary role in the oncology section NVALT, guideline committee NSCLC and CUP, dure genesmiddelen committee NVALT and FMS. All other authors declare no competing interests.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.lanepe.2024.100838>.

References

- 1 Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2021;71(3):209–249.
- 2 Malvezzi M, Santucci C, Boffetta P, et al. European cancer mortality predictions for the year 2023 with focus on lung cancer. *Ann Oncol*. 2023;34(4):410–419.
- 3 Gregor A, Inage T, Hwangbo B, Yasufuku K. Lung cancer staging: state of the art in the era of ablative therapies and surgical segmentectomy. *Respirology*. 2020;25(9):924–932.
- 4 Howlader N, Forjaz G, Mooradian MJ, et al. The effect of advances in lung-cancer treatment on population mortality. *N Engl J Med*. 2020;383(7):640–649.
- 5 Hendriks LE, Kerr KM, Menis J, et al. Oncogene-addicted metastatic non-small-cell lung cancer: ESMO clinical practice guideline for diagnosis, treatment and follow-up. *Ann Oncol*. 2023;34(4):339–357.
- 6 Thunnissen E, Weynand B, Udovicic-Gagula D, et al. Lung cancer biomarker testing: perspective from Europe. *Transl Lung Cancer Res*. 2020;9(3):887–897.
- 7 Ryska A, Berzinec P, Brcic L, et al. NSCLC molecular testing in Central and Eastern European countries. *BMC Cancer*. 2018;18(1):269.
- 8 Kerr KM, Bibeau F, Thunnissen E, et al. The evolving landscape of biomarker testing for non-small cell lung cancer in Europe. *Lung Cancer*. 2021;154:161–175.
- 9 Normanno N, Apostolidis K, Wolf A, et al. Access and quality of biomarker testing for precision oncology in Europe. *Eur J Cancer*. 2022;176:70–77.
- 10 Bayle A, Bonastre J, Chaltiel D, et al. ESMO study on the availability and accessibility of biomolecular technologies in oncology in Europe. *Ann Oncol*. 2023;34(10):934–945.
- 11 de Jager VD, Timens W, Bayle A, et al. Future perspective for the application of predictive biomarker testing in advanced stage non-small cell lung cancer. *Lancet Reg Health Eur*. 2024;38:100839. <https://doi.org/10.1016/j.lanepe.2024.100839>.
- 12 Houda I, Dickhoff C, Uyl-de Groot CA, et al. New systemic treatment paradigms in resectable non-small cell lung cancer and variations in patient access across Europe. *Lancet Reg Health Eur*. 2024;38:100840. <https://doi.org/10.1016/j.lanepe.2024.100840>.
- 13 Adib E, Nassar AH, Abou Alaiwi S, et al. Variation in targetable genomic alterations in non-small cell lung cancer by genetic

- ancestry, sex, smoking history, and histology. *Genome Med.* 2022;14(1):39.
- 14 Koopman B, Cajiiao Garcia BN, Kuijpers C, et al. A nationwide study on the impact of routine testing for EGFR mutations in advanced NSCLC reveals distinct survival patterns based on EGFR mutation subclasses. *Cancers.* 2021;13(14):3641.
 - 15 Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med.* 2018;378(2):113–125.
 - 16 Yang JC, Sequist LV, Geater SL, et al. Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6. *Lancet Oncol.* 2015;16(7):830–838.
 - 17 Elamin YY, Robichaux JP, Carter BW, et al. Pozotinib for EGFR exon 20-mutant NSCLC: clinical efficacy, resistance mechanisms, and impact of insertion location on drug sensitivity. *Cancer Cell.* 2022;40(7):754–767.e6.
 - 18 Zhou C, Ramalingam SS, Kim TM, et al. Treatment outcomes and safety of mobocertinib in platinum-pretreated patients with EGFR Exon 20 insertion-positive metastatic non-small cell lung cancer: a phase 1/2 open-label nonrandomized clinical trial. *JAMA Oncol.* 2021;7(12):e214761.
 - 19 Park K, Haura EB, Leighl NB, et al. Amivantamab in EGFR exon 20 insertion-mutated non-small-cell lung cancer progressing on platinum chemotherapy: initial results from the CHRYSALIS phase I study. *J Clin Oncol.* 2021;39(30):3391–3402.
 - 20 Duke ES, Stapleford L, Drezner N, et al. FDA approval summary: mobocertinib for metastatic non-small cell lung cancer with EGFR exon 20 insertion mutations. *Clin Cancer Res.* 2023;29(3):508–512.
 - 21 Chon K, Larkins E, Chatterjee S, et al. FDA approval summary: amivantamab for the treatment of patients with non-small cell lung cancer with EGFR exon 20 insertion mutations. *Clin Cancer Res.* 2023;29(17):3262–3266.
 - 22 Wang C, Zhao K, Hu S, Dong W, Gong Y, Xie C. Clinical outcomes of afatinib versus osimertinib in patients with non-small cell lung cancer with uncommon EGFR mutations: a pooled analysis. *Oncologist.* 2023;28(6):e397–e405.
 - 23 Robichaux JP, Le X, Vijayan RSK, et al. Structure-based classification predicts drug response in EGFR-mutant NSCLC. *Nature.* 2021;597(7878):732–737.
 - 24 Wu YL, Tsuboi M, He J, et al. Osimertinib in resected EGFR-mutated non-small-cell lung cancer. *N Engl J Med.* 2020;383(18):1711–1723.
 - 25 Houda I, Dickhoff C, Uyl-de Groot CA, et al. Challenges and controversies in resectable non-small cell lung cancer: a clinician's perspective. *Lancet Reg Health Eur.* 2024;38:100841. <https://doi.org/10.1016/j.lanepe.2024.100841>.
 - 26 Koopman B, Groen HJM, Schuurin E, et al. Actionability of on-target ALK resistance mutations in patients with non-small cell lung cancer: local experience and review of the literature. *Clin Lung Cancer.* 2022;23(2):e104–e115.
 - 27 Ariyasu R, Kakuto S, Miyadera K, et al. Real-world outcome analysis of patients with stage IV NSCLC treated with tyrosine kinase and immune checkpoint inhibitors. *JTO Clin Res Rep.* 2023;4(6):100524.
 - 28 Ou SI, Nagasaka M, Brazel D, Hou Y, Zhu VW. Will the clinical development of 4th-generation "double mutant active" ALK TKIs (TPX-0131 and NVL-655) change the future treatment paradigm of ALK+NSCLC? *Transl Oncol.* 2021;14(11):101191.
 - 29 Shaw AT, Ou SH, Bang YJ, et al. Crizotinib in ROS1-rearranged non-small-cell lung cancer. *N Engl J Med.* 2014;371(21):1963–1971.
 - 30 Drilon A, Siena S, Dziadziuszko R, et al. Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three phase 1-2 trials. *Lancet Oncol.* 2020;21(2):261–270.
 - 31 Ou SI, Zhu VW. CNS metastasis in ROS1+ NSCLC: an urgent call to action, to understand, and to overcome. *Lung Cancer.* 2019;130:201–207.
 - 32 Dankner M, Rose AAN, Rajkumar S, Siegel PM, Watson IR. Classifying BRAF alterations in cancer: new rational therapeutic strategies for actionable mutations. *Oncogene.* 2018;37(24):3183–3199.
 - 33 Lim GHT, Balbi KJ, Poskitt B, Bennett P, Moore DA. Prevalence and breakdown of non-small cell lung cancer BRAF driver mutations in a large UK cohort. *Lung Cancer.* 2022;173:71–74.
 - 34 Dankner M, Wang Y, Fazelzad R, et al. Clinical activity of mitogen-activated protein kinase-targeted therapies in patients with non-V600 BRAF-mutant tumors. *JCO Precis Oncol.* 2022;6:e2200107.
 - 35 Forsythe A, Zhang W, Phillip Strauss U, Fellous M, Korei M, Keating K. A systematic review and meta-analysis of neurotrophic tyrosine receptor kinase gene fusion frequencies in solid tumors. *Ther Adv Med Oncol.* 2020;12:1758835920975613.
 - 36 Hong DS, DuBois SG, Kummer S, et al. Larotrectinib in patients with TRK fusion-positive solid tumours: a pooled analysis of three phase 1/2 clinical trials. *Lancet Oncol.* 2020;21(4):531–540.
 - 37 Drilon A, Siena S, Ou SI, et al. Safety and antitumor activity of the multitargeted pan-TRK, ROS1, and ALK inhibitor entrectinib: combined results from two phase I trials (ALKA-372-001 and STARTRK-1). *Cancer Discov.* 2017;7(4):400–409.
 - 38 Drilon A, Tan DSW, Lassen UN, et al. Efficacy and safety of larotrectinib in patients with tropomyosin receptor kinase fusion-positive lung cancers. *JCO Precis Oncol.* 2022;6:e2100418.
 - 39 Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol.* 2020;21(2):271–282.
 - 40 Drilon A, Oxnard GR, Tan DSW, et al. Efficacy of selpercatinib in RET fusion-positive non-small-cell lung cancer. *N Engl J Med.* 2020;383(9):813–824.
 - 41 Gainor JF, Curigliano G, Kim DW, et al. Pralsetinib for RET fusion-positive non-small-cell lung cancer (ARROW): a multi-cohort, open-label, phase 1/2 study. *Lancet Oncol.* 2021;22(7):959–969.
 - 42 Hobbs GA, Der CJ, Rossman KL. RAS isoforms and mutations in cancer at a glance. *J Cell Sci.* 2016;129(7):1287–1292.
 - 43 Terrenato I, Ercolani C, Di Benedetto A, et al. A real-world systematic analysis of driver mutations' prevalence in early- and advanced-stage NSCLC: implications for targeted therapies in the adjuvant setting. *Cancers.* 2022;14(12):2971.
 - 44 Garcia BNC, van Kempen LC, Kuijpers C, Schuurin E, Willems SM, van der Wekken AJ. Prevalence of KRAS p.(G12C) in stage IV NSCLC patients in the Netherlands; a nation-wide retrospective cohort study. *Lung Cancer.* 2022;167:1–7.
 - 45 Isaksson J, Berglund A, Louie K, et al. KRAS G12C mutant non-small cell lung cancer linked to female sex and high risk of CNS metastasis: population-based demographics and survival data from the National Swedish Lung Cancer Registry. *Clin Lung Cancer.* 2023;24(6):507–518.
 - 46 Cox AD, Fesik SW, Kimmelman AC, Luo J, Der CJ. Drugging the undruggable RAS: mission possible? *Nat Rev Drug Discov.* 2014;13(11):828–851.
 - 47 Ostrem JM, Peters U, Sos ML, Wells JA, Shokat KM. K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions. *Nature.* 2013;503(7477):548–551.
 - 48 Lanman BA, Allen JR, Allen JG, et al. Discovery of a covalent inhibitor of KRAS(G12C) (AMG 510) for the treatment of solid tumors. *J Med Chem.* 2020;63(1):52–65.
 - 49 Skoulidis F, Li BT, Dy GK, et al. Sotorasib for lung cancers with KRAS p.G12C mutation. *N Engl J Med.* 2021;384(25):2371–2381.
 - 50 de Langen AJ, Johnson ML, Mazieres J, et al. Sotorasib versus docetaxel for previously treated non-small-cell lung cancer with KRAS(G12C) mutation: a randomised, open-label, phase 3 trial. *Lancet.* 2023;401(10378):733–746.
 - 51 Fujino T, Suda K, Mitsudomi T. Lung cancer with MET exon 14 skipping mutation: genetic feature, current treatments, and future challenges. *Lung Cancer.* 2021;12:35–50.
 - 52 Awad MM, Oxnard GR, Jackman DM, et al. MET exon 14 mutations in non-small-cell lung cancer are associated with advanced age and stage-dependent MET genomic amplification and c-met overexpression. *J Clin Oncol.* 2016;34(7):721–730.
 - 53 Remon J, Hendriks LEL, Mountzios G, et al. MET alterations in NSCLC-current perspectives and future challenges. *J Thorac Oncol.* 2023;18(4):419–435.
 - 54 Wolf J, Seto T, Han JY, et al. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med.* 2020;383(10):944–957.
 - 55 Paik PK, Filip E, Veillon R, et al. Tepotinib in non-small-cell lung cancer with MET exon 14 skipping mutations. *N Engl J Med.* 2020;383(10):931–943.
 - 56 Planchard D, Popat S, Kerr K, et al. Metastatic non-small cell lung cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2018;29(Suppl 4):iv192–iv237.
 - 57 NHS England. National genomic test directory (version June 2023). <https://www.england.nhs.uk/publication/national-genomic-test-directories/>. Accessed July 13, 2023.

- 58 Lung cancer: diagnosis and management (NICE guideline [NG122]); 2023. <https://www.nice.org.uk/guidance/ng122>. Accessed July 2, 2023.
- 59 Teixeira ME, Oliveira J, Borralho P, et al. Portuguese consensus recommendations for next-generation sequencing of lung cancer, rare tumors, and cancers of unknown primary origin in clinical practice. *Acta Med Port.* 2022;35(9):677–690.
- 60 Popper HH, Gruber-Mosenbacher U, Pall G, et al. The 2020 update of the recommendations of the Austrian working group on lung pathology and oncology for the diagnostic workup of non-small cell lung cancer with focus on predictive biomarkers. *Memo.* 2020;13(1):11–26.
- 61 Janžič U. POROČILO BOLNIŠNIČNEGA REGISTRA TUMORJEV PRSNEGA KOŠA KLINIKE GOLNIK ZA LETA 2010 – 2020; 2022. <https://www.klinika-golnik.si/dejavnost-klinike/onkoloska-dejavnost/klinicni-register-bolnikov-z-rakom-pljuc>. Accessed July 2, 2023.
- 62 Oncologie: Terugbetaling van de moleculair biologische testen met “next generation sequencing” (NGS). <https://www.riziv.fgov.be/nl/professionals/verzorgingsinstellingen/laboratoria/Paginas/oncologie-terugbetalings-moleculair-biologische-ngs.aspx>. Accessed October 5, 2023.
- 63 Griesinger F, Absenger G, Bleckmann A, et al. Lungenkarzinom, nicht-kleinzellig (NSCLC); 2022. <https://www.onkopedia.com/de/onkopedia/guidelines/lungenkarzinom-nicht-kleinzellig-nsclc/@@guideline/html/index.html>. Accessed July 2, 2023.
- 64 Patients atteints d'un cancer bronchique non à petites cellules: indications des tests moléculaires en vue de la prescription de traitements de précision; 2023. <https://www.e-cancer.fr/Expertises-et-publications/Catalogue-des-publications/Patients-atteints-d-un-cancer-bronchique-non-a-petites-cellules-indications-des-tests-moleculaires-en-vue-de-la-prescription-de-traitements-de-precision>. Accessed August 14, 2023.
- 65 Federatie Medisch Specialisten (FMS) - Richtlijn Niet-kleincellig longcarcinoom; 2020. https://richtlijndatabase.nl/richtlijn/niet_kleincellig_longcarcinoom/startpagina_-_niet-kleincellig_longcarcinoom.html. Accessed July 2, 2023.
- 66 Lungcancer - Nationellt vårdprogram (v7.0); 2023. <https://kunska-psbanken.cancercentrum.se/globalassets/cancerdiagnoser/lungaoch-lungosack/vardprogram/nationellt-vardprogram-lungcancer.pdf>. Accessed July 3, 2023.
- 67 Isla D, Lozano MD, Paz-Ares L, et al. [New update to the guidelines on testing predictive biomarkers in non-small-cell lung cancer: a National Consensus of the Spanish Society of Pathology and the Spanish Society of Medical Oncology]. *Rev Esp Patol.* 2023;56(2):97–112.
- 68 Direção-Geral da Saúde (DGS) - Diagnóstico e Tratamento do Carcinoma de Não Pequenas Células do Pulmão; 2013. <https://normas.dgs.min-saude.pt/wp-content/uploads/2019/09/diagnostico-e-tratamento-do-carcinoma-de-nao-pequenas-celulas-do-pulmao.pdf>. Accessed July 5, 2023.
- 69 Drilon A, Duruisseaux M, Han JY, et al. Clinicopathologic features and response to therapy of NRG1 fusion-driven lung cancers: the eNRGy1 global multicenter registry. *J Clin Oncol.* 2021;39(25):2791–2802.
- 70 Burghuber OC, Kirchbacher K, Mohn-Staudner A, et al. Results of the Austrian national lung cancer audit. *Clin Med Insights Oncol.* 2020;14:1179554920950548.
- 71 Adizie JB, Tweedie J, Khakwani A, et al. Biomarker testing for people with advanced lung cancer in England. *JTO Clin Res Rep.* 2021;2(6):100176.
- 72 Lamy T, Cabarro B, Planchard D, et al. Biomarker testing in older patients treated for an advanced or metastatic non-squamous non-small-cell lung cancer: the French ESME real-life multicenter cohort experience. *Cancers.* 2021;14(1):92.
- 73 Griesinger F, Eberhardt W, Nusch A, et al. Biomarker testing in non-small cell lung cancer in routine care: analysis of the first 3,717 patients in the German prospective, observational, nation-wide CRISP registry (AIO-TRK-0315). *Lung Cancer.* 2021;152:174–184.
- 74 Steeghs EMP, Groen HJM, Schuur E, et al. Mutation-tailored treatment selection in non-small cell lung cancer patients in daily clinical practice. *Lung Cancer.* 2022;167:87–97.
- 75 Salas C, Martín-López J, Martínez-Pozo A, et al. Real-world biomarker testing rate and positivity rate in NSCLC in Spain: prospective central lung cancer biomarker testing registry (Lung-Path) from the Spanish Society of Pathology (SEAP). *J Clin Pathol.* 2022;75(3):193–200.
- 76 Kuipers C, van den Heuvel MM, Overbeek LIH, et al. [National variation in molecular diagnostics in metastatic lung cancer]. *Ned Tijdschr Geneesk.* 2018;162:D1607.
- 77 Eide IJZ, Nilssen Y, Stensland EM, Brustugun OT. Real-world data on EGFR and ALK testing and TKI usage in Norway-a nation-wide population study. *Cancers.* 2023;15(5):1505.
- 78 Baijal S, Crosbie P, Fenemore J, Desai K. Variations in genomic testing in non-small cell lung carcinoma: a healthcare professional survey of current practices in the UK. *Oncologist.* 2023;28(8):e699–e702.
- 79 Fox AH, Jett JR, Roy UB, et al. Knowledge and practice patterns among pulmonologists for molecular biomarker testing in advanced non-small cell lung cancer. *Chest.* 2021;160(6):2293–2303.
- 80 Roberts TJ, Kehl KL, Brooks GA, et al. Practice-level variation in molecular testing and use of targeted therapy for patients with non-small cell lung cancer and colorectal cancer. *JAMA Netw Open.* 2023;6(4):e2310809.
- 81 Cox S, Powell C, Morgan S. Implementing genomic testing for lung cancer into routine clinical practice - the Welsh experience. *Clin Oncol.* 2022;34(11):716–723.
- 82 de Jager VD, Cajiao Garcia BN, ter Elst A, et al. Optimalisatie van de zorgketen moleculaire diagnostiek bij niet-kleincellig longcarcinoom in Noord-Nederland. *Ned Tijdschr Oncol.* 2023;20:70–76.
- 83 Ossowski S, Neeman E, Borden C, et al. Improving time to molecular testing results in patients with newly diagnosed, metastatic non-small-cell lung cancer. *JCO Oncol Pract.* 2022;18(11):e1874–e1884.
- 84 Navani N, Butler R, Ibrahim S, et al. Optimising tissue acquisition and the molecular testing pathway for patients with non-small cell lung cancer: a UK expert consensus statement. *Lung Cancer.* 2022;172:142–153.
- 85 Deans ZC, Costa JL, Cree I, et al. Integration of next-generation sequencing in clinical diagnostic molecular pathology laboratories for analysis of solid tumours; an expert opinion on behalf of IQN Path ASBL. *Virchows Arch.* 2017;470(1):5–20.
- 86 Koopman B, van der Wekken AJ, Ter Elst A, et al. Relevance and effectiveness of molecular tumor board recommendations for patients with non-small-cell lung cancer with rare or complex mutational profiles. *JCO Precis Oncol.* 2020;4:393–410.
- 87 Koopman B, Groen HJM, Ligtenberg MJL, et al. Multicenter comparison of molecular tumor boards in the Netherlands: definition, composition, methods, and targeted therapy recommendations. *Oncologist.* 2021;26(8):e1347–e1358.
- 88 Stenzinger A, Moltzen EK, Winkler E, et al. Implementation of precision medicine in healthcare-a European perspective. *J Intern Med.* 2023;294(4):437–454.
- 89 Hofman P, Calabrese F, Kern I, et al. Real-world EGFR testing practices for non-small-cell lung cancer by thoracic pathology laboratories across Europe. *ESMO Open.* 2023;8(5):101628.