



## Short Communication

### Emergence of ST654 *Pseudomonas aeruginosa* co-harboring *bla*<sub>NDM-1</sub> and *bla*<sub>GES-5</sub> in novel class I integron In1884 from Bulgaria



## Keywords:

*Pseudomonas aeruginosa*  
Carbapenemase  
Carbapenem resistance  
New Delhi metallo- $\beta$ -lactamase  
Guiana extended-spectrum  $\beta$ -lactamase

Multidrug-resistant (MDR) *Pseudomonas aeruginosa* is a common cause of hospital-acquired infections. Carbapenem resistance, in particular, represents a substantial problem in terms of treatment of infections due to this pathogen and leads to increased mortality, longer duration of hospital stay and increased healthcare costs [1]. The class B New Delhi metallo- $\beta$ -lactamase 1 (NDM-1) carbapenemase has spread to and has been described in many bacterial species, including *P. aeruginosa* [2]. Another carbapenem-hydrolysing enzyme, the Guiana extended-spectrum  $\beta$ -lactamase-5 (GES-5), initially found in *Escherichia coli* in Greece and belonging to class A  $\beta$ -lactamase family, has also been detected in *P. aeruginosa* in Brazil, China, Spain and South Africa [3].

Here, we report the first detection of NDM-1-producing *P. aeruginosa* isolates in Bulgaria and the chromosomal co-harboring of *bla*<sub>NDM-1</sub> and *bla*<sub>GES-5</sub> genes in these MDR *P. aeruginosa* in novel class I integron In1884.

Five carbapenem-resistant (CR) *P. aeruginosa* strains were isolated from clinical samples of patients in two Bulgarian hospitals. The first two isolates, named Pae1250 and Pae1251, were recovered in September 2017 from urine samples ( $>10^5$  CFU/mL) of two different patients hospitalised in Alexandrovska University Hospital in Sofia. In August 2018, we detected three other CR *P. aeruginosa* isolates (Pae1252, Pae1255 and Pae1257) from tracheobronchial aspirates of three patients hospitalised in 'St. Ivan Rilski' University Hospital in Sofia. A CR *Klebsiella pneumoniae* (Kpn1256) was also isolated from the patient from whom CR *P. aeruginosa* Pae1255 was collected.

Antibiotic susceptibility testing was performed using Etest (bioMérieux, la Balme-les-Grottes, France). Colistin susceptibility was tested by broth microdilution method using MICRONAUT plate (MERLIN Diagnostika GmbH, Bornheim, Germany) and interpreted according to European Committee on Antimicrobial Susceptibility Testing (EUCAST) guidelines v9.0 ([http://www.eucast.org/clinical\\_breakpoints/](http://www.eucast.org/clinical_breakpoints/)). All six strains were resistant to carbapenems and had minimum inhibitory concentration (MIC) values of imipenem and meropenem  $>32$  mg/L. They were also resistant to ceftazidime, cefepime, piperacillin/tazobactam, ceftolozane/tazobactam, ceftazidime/avibactam, ciprofloxacin,


levofloxacin, amikacin, tobramycin and gentamicin, but were susceptible to colistin (MIC range 1–2 mg/L). Genomic DNA from all strains (five *P. aeruginosa* and one *K. pneumoniae*) was extracted using the MasterPure™ DNA Purification Kit (Epicentre Technologies Inc.) and sequenced (via  $2 \times 250$  bp, MiSeq, Illumina, San Diego, USA), and long-read sequencing (MinION, Oxford Nanopore Technologies, Oxford, UK) of Pae1250 and Pae1255 was done according to the manufacturer's instructions. Data analysis was performed using an in-house tool (BacPipe) [4], and single-nucleotide polymorphism (SNP) calling and genetic context analysis was performed using CLC Genomics Workbench v9.5.1 (Qiagen, Hilden, Germany).

Utilising short-read sequencing and read mapping, the *bla*<sub>NDM-1</sub> gene in Kpn1256 was carried on an IncFII plasmid. *bla*<sub>NDM-1</sub> in Kpn1256 was also flanked upstream by the truncated ISAb125 (IS30 family) and bleomycin resistance gene downstream. In addition, the isolate also carried *bla*<sub>CTX-M-15</sub>, *bla*<sub>SHV-11</sub>, *bla*<sub>CMY-4</sub> and *bla*<sub>TEM-1B</sub> genes and belonged to ST11, a high-risk clone repeatedly reported in the Balkan region [5]. The close relatedness between the two Bulgarian NDM-1-producing isolates in 2016 and Kpn1256, recovered in 2018, potentially indicates a persistent circulation of this NDM-1-carrying ST11 clone.

The five *P. aeruginosa* were clonally related, all belonging to ST654, carrying similar accessory genomes with no plasmids and the same profile of resistance genes. The core genomes of these *P. aeruginosa* encompassed 11–19 SNP differences, indicating the close relatedness between these strains (GenBank accession no. BioProject ID: PRJNA628735). In addition to the *bla*<sub>NDM-1</sub> gene, isolates Pae1250 and 1251 also carried *bla*<sub>GES-1</sub>, while Pae1252, 1255 and 1257 harboured *bla*<sub>GES-5</sub>. GES-5 is a variant of GES-1 with only one amino acid difference (Gly165Ser, previously reported as Gly170Ser) and, unlike GES-1, possesses carbapenem-hydrolysing activity.

Genes *bla*<sub>GES-5</sub> and *bla*<sub>NDM-1</sub> were integrated into the chromosome of PA1255 in a ca. 42-kb region of divergence (125 054–167 819 bp) situated between universal stress protein (*Usp*) and nucleotidyltransferase. *Bla*<sub>GES-5</sub> harboured the novel class 1 integron In1884 with the 5'CS-*bla*<sub>GES-5</sub>/*aadB*-3'CS gene cassette array. However, the 3'CS was interrupted by a second integron 5'CS with *sul3* as first (not a gene cassette but a structure called In0 found in other class 1 integrons). It seems that a truncated *attI1* follows the last gene cassette of the In1884 integron. The composite genetic element was organised as follows: In1884 with a regular 3'CS on one side, then an In0 element with *attI1-sul3* fusion and *ISCR1* and *bla*<sub>NDM-1</sub> on the other side (Fig. 1).

This divergence region consisted of other genes conferring resistance towards aminoglycosides [*strA*, *strB*, *aph(3')*-*Via*, *aadB*], sulfonamide (*sul1*) and tetracycline (*tetA* and *tetR*). Although *P. aeruginosa* isolates are intrinsically resistant to sulfonamides and



**Fig. 1.** Composite genetic structure of novel integron In1884 harbouring 5'CS-*bla*<sub>GES-5</sub>/*aadB*-3'CS gene cassette array. The new integron numbering is assigned based on the INTEGRALL database (<http://integrall.bio.ua.pt/>).

tetracyclines, the presence of this composite and its potential mobilisation are of particular concern as its transfer to other microorganisms, such as Enterobacterales, could incur resistance to multiple antimicrobial agents simultaneously, including the new  $\beta$ -lactam  $\beta$ -lactamase inhibitor combinations. To the best of our knowledge, this is the first report of the co-existence of *bla*<sub>NDM-1</sub> and *bla*<sub>GES-5</sub> in the same genetic element harboured in the chromosome and the first report of NDM-1-producing *P. aeruginosa* from hospitals in Bulgaria and a novel class 1 integron In1884. Moreover, the simultaneous presence of NDM-1 *K. pneumoniae* and *P. aeruginosa* isolates in one of the patient's samples implied an initial hypothesis of a potential horizontal transfer of *bla*<sub>NDM-1</sub> between these strains. Based on the analysis of the adjacent genetic structures by long-read sequencing, this hypothesis was rejected with the conclusion that the co-existence of two CR *bla*<sub>NDM-1</sub>-harbouring microorganisms in this patient was coincidental. However, this might have happened in other unresearched MDR isolates.

## Funding

TK, BBX and the research leading to these results have received support from the Innovative Medicines Initiative Joint Undertaking (<https://www.imi.europa.eu/>) under grant agreement no [115523] COMBACTE-NET (Combatting Bacterial Resistance in Europe) and no [115620] COMBACTE-CARE, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007–2013) and EFPIA companies' kind contribution. MNN is supported by COMPARE (Collaborative Management Platform for detection and Analyses of (Re-)emerging and Foodborne Outbreaks in Europe) [H2020 Grant number: 643476].

## Competing interests

No conflict of interest to declare.

## Ethical approval

Not required.

## References

- [1] Zavascki AP, Barth AL, Gonçalves AL, Moro AL, Fernandes JF, Martins AF, et al. The influence of metallo-beta-lactamase production on mortality in nosocomial *Pseudomonas aeruginosa* infections. *J Antimicrob Chemother* 2006;58:387–92.
- [2] Jovcic B, Lepsanovic Z, Suljagic V, Rackov G, Begovic J, Topisirovic L, et al. Emergence of NDM-1 metallo-beta-lactamase in *Pseudomonas aeruginosa* clinical isolates from Serbia. *Antimicrob Agents Chemother* 2011;55:3929–31, doi:<http://dx.doi.org/10.1128/AAC.00226-11>.

- [3] Girlich D, Poirel L, Szczepanowski R, Schlüter A, Nordmann P. Carbapenem-hydrolyzing GES-5-encoding gene on different plasmid types recovered from a bacterial community in a sewage treatment plant. *Appl Environ Microbiol* 2012;78:1292–5, doi:<http://dx.doi.org/10.1128/AEM.06841-11>.
- [4] Xavier BB, Mysara M, Bolzan M, Ribeiro-Gonçalves B, Alako BT, Harrison P, et al. BacPipe: a rapid, user-friendly whole-genome sequencing pipeline for clinical diagnostic bacteriology. *iScience* 2020;23:100769, doi:<http://dx.doi.org/10.1016/j.isci.2019.100769>.
- [5] Politi L, Gartzonika K, Spanakis N, Zarkotou O, Poulou A, Skoura L, et al. Emergence of NDM-1-producing *Klebsiella pneumoniae* in Greece: evidence of a widespread clonal outbreak. *J Antimicrob Chemother* 2019;74:2197–202, doi:<http://dx.doi.org/10.1093/jac/dkz176>.

Tomislav Kostyanev<sup>a,b,\*</sup>  
M.N. Nguyen<sup>a,b</sup>  
R. Markovska<sup>c</sup>  
P. Stankova<sup>c</sup>  
B.B. Xavier<sup>a,b</sup>  
C. Lammens<sup>a,b</sup>  
Y. Marteva-Proevska<sup>d</sup>  
T. Velinov<sup>d</sup>  
R. Cantón<sup>e,f</sup>  
H. Goossens<sup>a,b</sup>  
S. Malhotra-Kumar<sup>a,b</sup>

<sup>a</sup>Department of Medical Microbiology, University of Antwerp, Antwerp, Belgium

<sup>b</sup>Vaccine & Infectious Disease Institute, University of Antwerp, Antwerp, Belgium

<sup>c</sup>Department of Medical Microbiology, Faculty of Medicine, Medical University of Sofia, Sofia, Bulgaria

<sup>d</sup>Central Laboratory of Microbiology, Alexandrovska University Hospital, Sofia, Bulgaria

<sup>e</sup>Servicio de Microbiología, Hospital Universitario Ramón y Cajal and Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), Madrid, Spain

<sup>f</sup>Red Española de Investigación en Patología Infecciosa (REIPI), Madrid, Spain

\* Corresponding author at: Laboratory of Medical Microbiology, University of Antwerp, Campus Drie Eiken, Universiteitsplein 1, S.626, 2610 Wilrijk, Antwerp, Belgium.

E-mail address: [tomislav.kostyanev@uantwerpen.be](mailto:tomislav.kostyanev@uantwerpen.be) (T. Kostyanev).

Received 11 February 2020

Available online 22 June 2020