



Exposure levels, determinants and risk assessment of organophosphate flame retardants and plasticizers in adolescents (14–15 years) from the Flemish Environment and Health Study

Michiel Bastiaensen^a, Celine Gys^a, Ann Colles^b, Veerle Verheyen^{b,c}, Gudrun Koppen^b, Eva Govarts^b, Liesbeth Bruckers^d, Bert Morrens^e, Ilse Loots^e, Annelies De Decker^f, Vera Nelen^f, Tim Nawrot^g, Stefaan De Henauw^h, Nik Van Larebekeⁱ, Greet Schoeters^{b,c}, Adrian Covaci^{a,*}

^a Toxicological Center, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

^b VITO – Health, Boeretang 200, 2400 Mol, Belgium

^c Department of Biomedical Sciences, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

^d BioStat, Data Science Institute, Hasselt University, Martelarenlaan 42, 3500 Hasselt, Belgium

^e Faculty of Social Sciences, University of Antwerp, Sint-Jacobstraat 2, 2000 Antwerp, Belgium

^f Provincial Institute of Hygiene, Kronenburgstraat 45, 2000 Antwerp, Belgium

^g Centre for Environmental Sciences, Hasselt University, Agoralaan, 3590 Diepenbeek, Belgium

^h Department of Public Health, Ghent University, C. Heymanslaan 10, 9000 Gent, Belgium

ⁱ Analytical, Environmental and Geo-Chemistry, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium

ARTICLE INFO

Handling Editor: Martí Nadal

Keywords:

Organophosphate flame retardants
Exposure assessment
Human biomonitoring
FLEHS IV
Belgium
Adolescents

ABSTRACT

The ubiquitous use of organophosphate flame retardants and plasticizers (PFRs) in a variety of consumer products has led to widespread human exposure. Since certain PFRs are developmental and carcinogenic toxicants, detailed exposure assessments are essential to investigate the risk associated with environmental exposure levels. However, such data are still lacking for European countries. In this study, concentrations of thirteen PFR metabolites were measured in urine samples from 600 adolescents from Flanders, Belgium. 1-Hydroxy-2-propyl bis(1-chloro-2-propyl) phosphate (BCIPHIPP), diphenyl phosphate (DPHP), bis(1,3-dichloro-isopropyl) phosphate (BDCIPP), 2-hydroxyethyl bis(2-butoxyethyl) phosphate (BBOEHEP), 2-ethylhexyl phenyl phosphate (EHPPH) and 2-ethyl-5-hydroxyhexyl diphenyl phosphate (5-HO-EHDPPH) were frequently detected (>83%) in all participants. Comparisons with study populations from outside the EU showed that urinary levels of DPHP, BDCIPP and BCIPHIPP were generally within the same range. Only exposure to 2-ethylhexyl diphenyl phosphate (EHDPPH) was presumably higher in Flemish adolescents. However, determinants analysis through multivariate regression analyses did not reveal significant predictors that may explain this finding. Significantly higher levels of BDCIPP were observed in participants with new decorations at home, while adolescents with highly educated parents had higher levels of BBOEHEP and BDCIPP. Furthermore, multiple PFR metabolite concentrations followed a seasonal pattern. Estimated daily intakes (EDIs) were calculated from the internal dose by including fractions of urinary excretion (F_{UE}) estimated in *in vitro* metabolism studies. EDIs ranged from 6.3 ng/kg bw/day for TBOEP to 567.7 ng/kg bw/day for EHDPPH, which were well below the available oral reference doses for all investigated PFRs. This suggests that the associated risk is low at present. This is the first report on internal exposure to seven commonly used PFRs in a European population.

1. Introduction

Flame retardants are substances added to textiles, furniture, plastics, electronic devices, upholstery and other consumer goods to reduce the

risk of fire spreading and to meet flammability standards (van der Veen and de Boer, 2012). Until recently, polybrominated diphenyl ethers (PBDEs) were the most popular flame retardants used worldwide. However, the production and use of PBDE mixtures such as Penta- and

* Corresponding author.

E-mail address: adrian.covaci@uantwerpen.be (A. Covaci).

<https://doi.org/10.1016/j.envint.2020.106368>

Received 28 September 2020; Received in revised form 22 December 2020; Accepted 23 December 2020

Available online 6 January 2021

0160-4120/© 2020 The Author(s).

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Deca-BDE is now prohibited due to their listing as persistent organic pollutants (UNEP, 2009, 2017). Some of these mixtures already contained organophosphate flame retardants (PFRs), a class of organic chemicals that has recently overtaken PBDEs as most frequently used flame retardants (van der Veen and de Boer, 2012). PFRs are tri-esters of phosphate with varying side-chains. As a result, they are applied in a wide variety of products such as furniture, textiles, decoration and building materials, paints, floor polish, resins and polyvinyl chloride (PVC) plastics for their fire safety and/or plasticizing properties (Wei et al., 2015). Chlorinated PFRs, such as tris(2-chloroethyl) phosphate (TCEP), tris(1-chloro-isopropyl) phosphate (TCIPP) and tris(1,3-dichloro-isopropyl) phosphate (TDCIPP) and non-halogenated alkyl- and aryl-phosphates, such as triphenyl phosphate (TPHP), tri-*n*-butyl phosphate (TNBP), tris(2-butoxyethyl) phosphate (TBOEP) and 2-ethylhexyl diphenyl phosphate (EHDPHP) have been found in high levels in indoor dust, air and consumer products because PFRs are additive rather than reactive chemicals (Stapleton et al., 2009; Cequier et al., 2014; Kim et al., 2019; Li et al., 2019c). Potential routes of human exposure to PFRs not only include the ingestion of dust, inhalation of air or dermal contact (Xu et al., 2016), but also the intake of contaminated food as they have also been found in several food categories as a result of contamination during industrial processing (e.g., plastic packaging) (Poma et al., 2018; Li et al., 2019a). Levels of PFRs in the indoor environment now exceed those of PBDEs due to a sharp increase in PFR usage since the beginning of this century (Dodson et al., 2012). Exact numbers on production volumes are scarce, but all the PFRs of interest to this work are currently registered as high production volume chemicals under the European REACH regulation (ECHA, 2020). The total consumption of flame retardants in Europe was 465,000 tons in 2006 (of which 20% were PFRs) and is likely several times higher at this moment. Moreover, the worldwide consumption of flame retardants is expected to grow and exceed 4 million tons by 2025 (Grand View Research 2017).

Although PFRs were introduced as suitable alternatives to PBDEs, evidence from *in vitro* and *in vivo* toxicity tests suggest that exposure to these chemicals can lead to adverse health effects for humans and biota due to PFRs' carcinogenic, neurotoxic and endocrine disrupting potential (van der Veen and de Boer, 2012; Wei et al., 2015). Epidemiological evidence is scarce, but exposure to PFRs has been associated with adverse reproductive outcomes (Carignan et al., 2017; Ingle et al., 2018), increased odds of papillary thyroid cancer (Hoffman et al., 2017b), allergic symptoms (Araki et al., 2018) and increased oxidative stress biomarkers (Ait Bamai et al., 2019). As such, reliable exposure and risk assessments of human populations have gained significant scientific interest in recent years. Exposure to environmental contaminants is commonly monitored through the measurement of biomarkers in urine or blood. *In vitro* and *in vivo* studies have shown that PFRs are rapidly metabolized to dialkyl and diaryl phosphates (DAPs) such as diphenyl phosphate (DHP), bis(butoxyethyl) phosphate (BBOEP), bis(1,3-dichloro-isopropyl)phosphate (BDCIPP), and to different hydroxylation products (HO-PFRs), like 1-hydroxy-2-propyl bis(1-chloro-2-propyl) phosphate (BCIPHIPP) and 2-ethyl-5-hydroxyhexyl diphenyl phosphate (5-HO-EHDPHP), and 2-hydroxyethyl bis(2-butoxyethyl) phosphate (BBOEHEP) (Van den Eede et al., 2013; Ballesteros-Gómez et al., 2015; Völkel et al., 2018). These urinary metabolites have been used as biomarkers of exposure in studies from the US (Hoffman et al., 2018; Ospina et al., 2018), Australia (He et al., 2018a, 2018b), China (Chen et al., 2018; Sun et al., 2018; Ding et al., 2019) and Japan (Bastiaensen et al., 2019; Bastiaensen et al., 2020a). Although exposure levels of PFRs in humans are increasing as shown for populations in the US and Japan (Hoffman et al., 2017a; Bastiaensen et al., 2020a), the number of biomonitoring studies investigating PFR exposure in Europe is still very limited (Fromme et al., 2014; Cequier et al., 2015). Comprehensive analyses of internal exposure are needed to determine the extent of human exposure and the associated risk, especially in industrialized areas such as Europe. In the present study, we report the urinary concentrations of thirteen metabolites of seven frequently used

PFRs in adolescents from Flanders, Belgium. We also investigated which factors could significantly impact PFR exposure and if current exposure levels are of concern through the calculation of estimated daily intakes (EDI) and comparison with available reference doses (RfD).

2. Materials and methods

2.1. Study population

The Flemish Environment and Health Study (FLEHS), established in 2002, generates information on the presence and distribution of biomarkers of environmental pollutants relevant for public health. The study was implemented, funded and steered by the Environment, Nature and Energy Department of the Flemish government and carried out by a research consortium, the Centre for Expertise on Environment and Health. In previous cycles (2002–2015), reference values for persistent organic pollutants (POPs), metals, pesticides, phthalates, bisphenol A and polycyclic aromatic hydrocarbons (PAHs), among others, have been established for newborns, adolescents and adults (Schoeters et al., 2012). In the fourth cycle (FLEHS IV, 2016–2020), 600 adolescents (14–15 years old) were recruited in two parts as a sample of the general population of Flanders (northern part of Belgium): 200 newborns of the first FLEHS campaign (2002–2006, now adolescents) were asked to participate in FLEHS IV, as well as 400 adolescents newly recruited through schools. One of the objectives of the current program was to investigate the internal exposure to emerging contaminants, such as PFRs, new/alternative plasticizers and perfluoroalkyl substances in relation to (eco-)behavior and potential health effects. Participants were recruited through 20 schools from all five provinces of Flanders between September 2017 and June 2018. Inclusion criteria were: participants had to reside in Flanders for at least 5 years and had to be able to fill in questionnaires in Dutch. Each participant provided urine, hair and blood samples; these were immediately processed during fieldwork at school. Urine samples were collected in clean polyethylene containers, aliquoted into glass vials and stored frozen (-20 °C) until analysis by the responsible laboratory. The body weight (bw) and height of the adolescents were measured by trained nurses during fieldwork. Participants (adolescents and/or parents) completed self-administered questionnaires on personal and lifestyle factors, such as education, income, smoking and eating habits, and on the use of certain materials or products in their home environment (digital or on paper, carried out at home). The characteristics of the study population are shown in Table SI-1. Adolescents and their parents provided a written informed consent. The study protocol was approved by the Ethical Committee of the Antwerp University Hospital (Belgian Registry Number: B300201732753). All data were pseudonymized.

2.2. Measurement of PFR biomarkers in urine

Target analytes in this study were 1-hydroxy-2-propyl bis(1-chloro-2-propyl) phosphate (BCIPHIPP), bis(1-chloro-2-propyl) phosphate (BCIPP), diphenyl phosphate (DHP), 4-hydroxyphenyl phenyl phosphate (4-HO-DHP), hydroxyphenyl diphenyl phosphate (HO-TPHP), dibutyl phosphate (DNBP), bis(1,3-dichloro-isopropyl) phosphate (BDCIPP), tris(2-chloroethyl) phosphate (TCEP), 2-hydroxyethyl bis(2-butoxyethyl) phosphate (BBOEHEP), bis(butoxyethyl) phosphate (BBOEP), (3-hydroxy)-2-butoxyethyl bis(2-butoxyethyl) phosphate (3-HO-TBOEP), 2-ethylhexyl phenyl phosphate (EHPHP) and 2-ethyl-5-hydroxyhexyl diphenyl phosphate (5-HO-EHDPHP).

The method used for the extraction and quantification of PFR metabolites in urine has been discussed in detail elsewhere (Bastiaensen et al., 2018). Briefly, urine samples (2 mL) were spiked with mass-labelled internal standards (5 ng), adjusted to pH 6 with phosphate buffer (1 M) and deconjugated with β -glucuronidase (2 mg/mL). Samples were extracted on Bond-Elut C18 cartridges (3 mL, 200 mg, Agilent, Santa Clara, USA), conditioned with 3 mL of methanol and 2 mL of

water. Target analytes were eluted with methanol (3 mL) and filtered in a micro-centrifuge (0.2 µm nylon, VWR, Leuven, Belgium). Final extracts (water:methanol 1:1) were injected on a Agilent 1290 Infinity liquid chromatography system coupled to a triple quadrupole mass spectrometer (ESI-6460, Agilent). Separation was obtained by a biphenyl column (2.1 mm × 100 mm, 2.6 µm; Phenomenex Kinetex; Torrance, USA) and mobile phases of water with 2% methanol (A) and methanol with 2% water (5 mM ammonium acetate as additive). Target analytes were identified through dynamic multiple reaction monitoring (dMRM) in positive and negative ionization.

Quantification of target analytes was performed with calibration curves in neat solvents ranging from 0.04 to 10 ng/mL, except for BCIPP and 4-HO-DPHP for which the calibration ranged from 0.2 to 50 ng/mL. Procedural blanks were analyzed to check for background contamination. DPHP, DNBP and TCEP were found at trace levels in procedural blanks (0.14, 0.08 and 0.03 ng/mL, respectively). These blank concentrations were subtracted from values in urine samples. Limits of quantification (LOQ) were determined as three times the standard deviation of procedural blank concentrations (for DPHP, DNBP and TCEP) or as the concentration corresponding to a signal-to-noise ratio of 10 in spiked urine samples. Successful participation in inter-laboratory comparison exercises (HBM4EU ICI/EQUAS and OSEQAS, 2018–2020) assured additional external quality control for several target analytes (DPHP, BDCIPP and BCIPP). Metabolites and corresponding parent chemicals targeted in this study are shown in Table SI-2. The performance of the analytical method is summarized in Tables SI-2 and 3. Specific gravity was determined by refractometry at Algemeen Medisch Labo (AML, Antwerp, Belgium).

2.3. Statistical analysis

Statistical analyses were carried out for compounds which had exposure values above LOQ in at least 60% of the samples. A truncated lognormal distribution was fitted to the observed values (above the LOQ). This resulted in an estimation of the mean and standard deviation for the distribution of all log-normal transformed values (above and below LOQ). Values below the LOQ were then randomly assigned an imputed value below the log of the LOQ, drawn from the estimated lognormal distribution and retransformed to an imputed value between 0 and the LOQ on the original scale. The description of this imputation protocol and software has been recently communicated in a report for HBM4EU project – Dec 2020 (Statistical analysis plan for the co-funded studies – Deliverable 10.10 – <https://www.hbm4eu.eu/deliverables/>).

Questionnaire data were used to identify significant determinants of exposure by a stepwise multiple linear regression model per compound using backward selection. Dependent variables were PFR metabolite levels normalized for specific gravity using the following formula: $\text{biomarker_SG} = \text{conc}_{\text{biomarker}} * (1.024-1)/(SG-1)$, and then transformed by the natural logarithm. Independent variables were introduced in the multiple model if the p-value was <0.2 in univariate regression model and if the direction of the association was consistent with mechanistic or epidemiological insights. Correlations among independent variables was also verified by variance inflation factors. Non-significant explanatory variables ($p < 0.05$) were removed one by one until only significant variables were retained. Season and socio-economic variables were only introduced in the final step as they could be proxies for other determinants. The available variables that were tested in univariate regression models are shown in Table SI-4. Specific gravity was also added as an independent variable in each model. The R-square of the model reflects the percentage of variation in PFR metabolite levels that could be explained by the remaining independent variables in the final model.

Estimated daily intakes (EDI in ng/kg bw/day) were calculated based on urinary concentrations of frequently detected metabolites according to the following equation (Fromme et al., 2014; Chen et al., 2018):

$$EDI = \left(\frac{c_{\text{meta}} \times V_{\text{urine}}}{F_{\text{UE}} \times bw} \right) \times \frac{MW_p}{MW_m}$$

where c_{meta} is the specific-gravity corrected metabolite concentration (in ng/mL SG); V_{urine} is the daily excreted volume of urine (estimated at 1200 mL/day for adolescents) (Valentin 2002); F_{UE} is the urinary excretion factor; bw is the body weight of the participant (in kg); and MW_p and MW_m are the molecular weight of the parent compound and its metabolite respectively (in g/mol, Table SI-2). Studies on the toxicokinetics of PFRs in humans are scarce. Therefore, results from *in vitro* human liver microsomes (HLM) and S9 fractions experiments (Van den Eede et al., 2013) were used to estimate the urinary excretion fraction (F_{UE}) as previously suggested by Zhang et al. (2018b). EDIs were calculated based on both scenarios (HLM and S9) to investigate the total daily exposure to PFRs (exact F_{UE} values shown in Table 4). The estimated daily intakes of targeted PFRs were compared to available oral reference doses (RfD). RfD values provide an estimation of the daily exposure for humans that is likely without any adverse effects during a lifetime. As such, RfDs are a valuable tool for risk assessment of human exposure to toxic chemicals (Poma et al., 2019).

3. Results and discussion

3.1. Concentrations of PFR metabolites

The distribution of targeted PFR metabolites in urine of Flemish adolescents is shown in Table 1. Six from the thirteen measured metabolites, originating from seven different parent compounds were detected in the majority of the samples: DPHP as a metabolite of TPHP (detection frequency 99%), BCIPHIPP as a metabolite of TCIPP (96%), BDCIPP as a metabolite of TDCIPP (86%), BBOEHEP as a metabolite of TBOEP (96%), and EHPHP and 5-HO-EHDPHP both metabolites of EHDPHP (99% and 98%, respectively). The high detection frequencies of these metabolites show that participants were widely exposed to all targeted PFRs except TCEP (33%) and TNBP (16%). Detection frequencies of other metabolites derived from the same parent compounds were much lower. EHPHP, DPHP and BCIPHIPP were detected at the highest uncorrected geometric mean concentrations of 3.56 ng/mL, 1.23 ng/mL and 0.79 ng/mL, respectively.

Until now, most studies have focused on diaryl and dialkyl metabolites (DAPs) such as DPHP, BDCIPP, DNBP and BBOEP as targets for human biomonitoring. HO-PFRs have only been measured by a limited number of studies although they are excellent targets for internal exposure assessment as shown by the high detection frequencies of BCIPHIPP, BBOEHEP and 5-HO-EHDPHP in this study. However, not all HO-PFR metabolites are found in detectable levels. For example, exposure to TPHP is commonly monitored by measuring DPHP, but correlations between dust and urine were not statistically significant in most studies (Cequier et al., 2015; Bastiaensen et al., 2019). TPHP is used as a plasticizer and lubricant, but also as a flame retardant in polyurethane foam in combination with chlorinated PFRs (van der Veen and de Boer, 2012). However, other PFRs, such as EHDPHP and resorcinol bis-diphenyl phosphate (RDP) also form DPHP as a metabolite in urine. Thus, DPHP concentrations should be interpreted as the exposure dose resulting from the sum of all PFRs with at least two phenyl substituents (Dodson et al., 2014; Ballesteros-Gómez et al., 2015). Hydroxylated TPHP metabolites (4-HO-DPHP and HO-TPHP) have been proposed as TPHP specific biomarkers, but until now these metabolites have only been reported in low detection frequencies (Bastiaensen et al., 2019; Ding et al., 2019; Li et al., 2019b; Bastiaensen et al., 2020a).

As mentioned before, comparable results from European studies are scarce, but representative data for adolescents, children, adults and the general population have recently been reported for other countries. Median concentrations of urinary PFR metabolites as reported by various studies are summarized in Table 2. Levels of DPHP in this study

Table 1

Concentrations of PFR metabolites in urine of Flemish adolescents (n = 582, in ng/mL). Geometric means only for compounds with DF > 60%. Imputed values for measurements values < LOQ.

Parent compound	PFR metabolites	LOQ (ng/mL)	DF (%)	Percentile					GM	95% CI	
				5th	25th	50th	75th	95th		Lower bound	Upper bound
TPHP	DPHP	0.10	99	0.23	0.70	1.22	2.01	4.43	1.23	1.14	1.34
	4-HO-DPHP	0.50	15				n.d.	1.16	n.a.		
	HO-TPHP	0.01	7				n.d.	0.01	n.a.		
TCEP	TCEP	0.04	33				0.05	0.15	n.a.		
TCIPP	BCIPHIPP	0.04	96	0.07	0.26	0.61	1.42	6.12	0.79	0.69	0.89
	BCIPP	1.00	9				n.d.	2.40	n.a.		
TNBP	DNBP	0.15	16				n.d.	0.36	n.a.		
TDCIPP	BDCIPP	0.05	83	n.d.	0.12	0.29	0.63	2.20	0.42	0.38	0.46
TBOEP	BBOEHEP	0.005	96	0.01	0.02	0.03	0.07	0.24	0.04	0.04	0.04
	BBOEP	0.05	13				n.d.	0.10	n.a.		
	3-HO-TBOEP	0.01	7				n.d.	0.01	n.a.		
EHDPHP	EHPHP	0.05	99	0.57	2.18	3.77	6.15	12.90	3.56	3.28	3.85
	5-HO-EHDPHP	0.01	98	0.02	0.04	0.08	0.15	0.34	0.09	0.08	0.09

n.d.: not detected; n.a.: not applicable; LOQ: limit of quantification; DF: detection frequency; GM: geometric mean.

were higher than levels observed in adolescents from China (Ding et al., 2019), but comparable to adolescents from the US (Ospina et al., 2018). BDCIPP and BCIPHIPP levels were lower compared to these study populations, but generally higher in comparison to levels observed in children and adults from Japan (Bastiaensen et al., 2019; Bastiaensen et al., 2020a), Norway (Cequier et al., 2015) and China (Chen et al., 2018; Zhang et al., 2018a, 2018b; Li et al., 2019b). However, higher urinary concentrations have also been reported for BDCIPP and DPHP in children from Australia (He et al., 2018b) and BDCIPP in children from the US (Butt et al., 2016). Differences in internal exposure may be explained by several factors. Firstly, younger children are relatively more frequent exposed to PFRs than adults (Van den Eede et al., 2015). As such, comparisons between different age categories should be performed with caution. Secondly, regulations and usage patterns of chemicals can differ considerably between countries. For example, TCEP use declined in Europe due to stricter legislation. TCEP was classified as a category 2 reproductive toxicant and a category 2 carcinogen (EU 2009)) which led to increased use of TCIPP and TDCIPP. Conversely, no regulations exist for TCEP in various Asian countries which in turn resulted in relatively higher human exposure to TCEP (Li et al., 2019c). Therefore, it is important to consider sampling year and location because PFR consumption patterns change over time.

According to the results of this study, Belgian adolescents were mainly exposed to EHDPHP, DPHP, TCIPP and TDCIPP. Concentrations of EHPHP and 5-HO-EHDPHP were particularly high compared to other studies from China and Japan (Hu et al., 2019; Li et al., 2019b; Bastiaensen et al., 2020a), but similar to results from Belgian adults (Bastiaensen et al., 2018). This indicates that these compounds are present in products, used or present in the life environment of both age groups. EHDPHP is applied as a flame retardant and plasticizer in polyvinyl chloride (PVC), but is also used in paints, rubber, adhesives and food packaging materials (Ballesteros-Gómez et al., 2015). Migration of EHDPHP from packaging to food is likely a major exposure source to this PFR. Recent studies have shown that PFRs, and especially EHDPHP, contaminate food during industrial processing and manipulation (e.g., packaging, drying, canning) (Poma et al., 2018). EHDPHP was the major contributor to the total dietary exposure of PFRs in food basket studies from Belgium (Poma et al., 2018), Sweden (Poma et al., 2017) and Norway (Xu et al., 2017). Dietary intake was responsible for over 95% of the total EHDPHP exposure, compared to limited exposure through dust or air (Xu et al., 2019). We therefore presume that the higher levels of EHPHP and 5-HO-EHDPHP in the urine of Belgian adolescents are associated with dietary exposure. However, we could not identify any significant dietary factors from questionnaires (discussed in Section

3.2). Moreover, future studies are required to confirm these findings since most other studies are not reporting these metabolites (Table 2). EHDPHP is also frequently detected in dust samples, but generally in lower concentrations compared to other PFRs (Li et al., 2019c). However, one study from the UK found that EHDPHP levels in classroom dust significantly exceeded those of all other microenvironments studied (living rooms, offices, cars) (Brommer and Harrad, 2015). It is therefore possible that exposure during school hours also contributed to increased EHDPHP exposure. Food intake is only a minor exposure route for other PFRs, such as TCIPP, TDCIPP, TPHP and TBOEP. Exposure to these PFRs mainly arises from dust ingestion and air inhalation (Xu et al., 2016). TCIPP and TBOEP have been reported in the highest concentrations in dust from houses and other indoor environments in different countries, followed by TDCIPP and TPHP (Li et al., 2019c). TCIPP and TBOEP were also the most abundant PFRs in house dust samples from Belgium in two different studies (Van den Eede et al., 2011; de la Torre et al., 2020). These PFRs are applied in many products present in the indoor environment: chlorinated PFRs, such as TCIPP and TDCIPP are used in polyurethane foam found the upholstery of furniture, TBOEP in floor polish and TPHP as flame retardant in polyurethane foam and as plasticizer in lacquers, hydraulic fluids, resins and PVC (van der Veen and de Boer, 2012). Significant correlations have been reported between TDCIPP, TCIPP and TBOEP in house dust and their metabolites in urine (BDCIPP, BCIPHIPP and BBOEHEP, respectively), which suggest that dust ingestion was an important exposure route to these three PFRs for the participants of this study (Cequier et al., 2015; Bastiaensen et al., 2019). However, levels in dust and urine are not always significantly correlated as shown in children from the US (Phillips et al., 2018).

3.2. Determinants of exposure

Participants provided detailed information about their home environment and lifestyle habits by answering questionnaires, which could be related to the measured internal exposure. Results of the multiple regression models are summarized in Table 3. Season was a significant determinant for all PFR metabolites. Other studies have reported that concentrations of BDCIPP, BCIPHIPP, DPHP and BBOEHEP were higher in summer (Hoffman et al., 2018; Bastiaensen et al., 2020a). In this study, levels of BDCIPP and DPHP were also higher in warmer months, but the trend was not consistent across all metabolites (e.g. EHPHP, BBOEHEP, 5-HO-EHDPHP). In univariate analysis, we found that BDCIPP, BCIPHIPP and DPHP levels increased with higher average daily outdoor temperature and higher average daily UV radiation (data not shown). Temperature may be an important predictor of exposure to

Table 2
Comparison of median concentrations of PFR metabolites (ng/mL) reported in various studies.

Reference	n	Country	Sampling years	Population	TPHP		TDCIPP	TCIPP		TNBP	TBOEP		EHDPHP		
					DPHP	4-HO-DPHP	BDCIPP	BCIPHP	BCIPP	DNBP	BBOEHEP	BBOEP	3-HO-TBOEP	EHPHP	5-HO-EHDPHP
<i>This study</i>	582	Belgium	2017–2018	Adolescents (14–15 y/o) ^a	1.37	n.d.	0.34	0.71	n.d.	n.d.	0.04	n.d.	n.d.	4.07	0.01
Ding et al. (2019)	306	China	2015	Adolescents (12–15 y/o) ^b	0.42	n.d.	6.17	1.47	0.18	2.54	n.d.	0.11	n.d.		
Ospina et al. (2018)	427	USA	2013–2014	Adolescents (12–19 y/o) ^b	1.44		1.43		0.16	0.21					
Ospina et al. (2018)	2666	USA	2013–2014	General population ^b	0.82		0.88		0.16	0.25					
Bastiaensen et al. (2020a)	400	Japan	2012–2017	Children (7 y/o) ^b	0.46	n.d.	0.13	0.38	n.d.	n.d.	0.11	0.11	n.d.	0.31	0.01
Bastiaensen et al. (2019)	128	Japan	2009–2010	Children (7–12 y/ ^b o)	0.32	0.22	0.07	0.18	n.d.	n.d.	0.36	0.2	n.d.	n.a.	0.04
Hoffman et al. (2018)	203	USA	2014–2016	Children (3–6 y/o) ^a	2.3		6	1.3	0.46						
Gibson et al. (2018)	90	USA	2015	Children (3–5y/o) ^b	3.2		2.6	0.6	0.9						
Butt et al. (2016)	40	USA	2015	Children (0–3 y/o) ^a	2.5		7.4	2.0	n.d.						
Cequier et al. (2015)	54	Norway	2012	Children (6–12 y/o) ^a	1		0.23			n.d.			n.d.		
Chen et al. (2018)	411	China	2015	Children (8–12 y/o) ^a	0.28		0.05		0.15	0.12			0.05		
Fromme et al. (2014)	312	Germany	2011–2012	Children (2–6 y/o) ^b	0.8					0.2			2		
He et al. (2018b)	400	Australia	2014–2015	Children (0–5y/o) ^c	25		2.6	0.43	0.85	0.18	0.075	0.32	0.029		
He et al. (2018a)	61	Australia	2015–2016	Children (0–2 y/o) ^b	1		3.3	0.93	0.68	0.15	0.08	0.1	0.13		
Zhang et al. (2018a)	227	China	2016	Children (0–5y/o) ^b	0.27		0.08		0.69	0.06			0.04		
Sun et al. (2018)	180	China	2016–2017	Adults and children ^a	0.07		n.d.		n.d.	0.00			0.10		
Hammel et al. (2016)	40	USA	2015	Adults ^b	1.16		2.06	1.12	n.d.						
Hu et al. (2019)	15	China	2018	Adults ^b	2.47		0.09	0.49	n.d.	0.16	n.d.	0.12	n.d.	n.d.	n.d.
Bastiaensen et al. (2018)	14	Belgium	2017	Adults ^b	1.09	n.d.	0.19	3.93	n.d.	0.26	0.02	n.d.	n.d.	3.94	0.23
Dodson et al. (2014)	16	USA	2011	Adults ^b	0.44		0.09		n.d.	0.11			n.d.		
Siddique et al. (2020)	120	Canada	2013–2015	Adult women ^b	13.8		1.2	n.d.	0.5		n.d.		n.d.		
Chen et al. (2019)	84	China	2013–2015	Adult women ^a	0.26		0.25		0.14	0.18			0.06		
Li et al. (2019b)	46	China	2017	General population ^b	0.33	n.d.	0.08	0.68	n.d.	n.d.	n.d.	n.d.		0.57	0.004
Zhang et al. (2018b)	323	China	2017	General population ^b	0.30		0.15		0.30	0.29		0.03			

^a Specific gravity corrected median values.

^b Uncorrected median values.

^c Pooled mean values.

Table 3
Multivariate analysis for the identification of determinants of exposure of PFR metabolites (ln-transformed and standardized for specific gravity).
 Only variables significant at $p < 0.05$ were retained in the model. Estimates are multiplicative factors.

Log adjusted PFR metabolite	Variable (n)	β (95%CI)	p-Value
<u>BDCIPP</u> $R^2 = 0.059$	Highest education level of the mother		0.033
	Primary education (42)	Reference	
	Secondary education (84)	1.328 (0.034–1.129)	0.243
	Tertiary education (337)	1.679 (1.065–2.646)	0.026
	New decoration in living room (curtains, carpet, wallpapers, etc)		
	No (465)	Reference	
	Yes (98)	1.590 (1.163–2.175)	0.004
	Season		0.001
	Winter (169)	Reference	
	Spring (191)	1.387 (1.029–1.868)	0.032
Summer (66)	2.291 (1.527–3.438)	<0.001	
Autumn (137)	1.319 (0.958–1.817)	0.089	
<u>DPHP</u> $R^2 = 0.092$	Highest education level of the father		0.021
	Primary education (60)	Reference	
	Secondary education (203)	1.198 (0.983–1.459)	0.073
	Tertiary education (231)	1.009 (0.829–1.228)	0.930
	Home with mechanical ventilation		
	No (403)	Reference	
	Yes (91)	0.848 (0.726–0.990)	0.037
	Season		<0.001
	Winter (145)	Reference	
	Spring (164)	1.384 (1.186–1.613)	<0.001
Summer (61)	1.383 (1.129–1.696)	0.002	
Autumn (124)	1.284 (1.090–1.513)	0.003	
BMI	<18.5 kg/m ² (139)	Reference	0.032
	18.5–25 kg/m ² (301)	0.884 (0.771–1.013)	0.077
	>25 kg/m ² (54)	0.757 (0.609–0.939)	0.012
<u>BCIPHIPP</u> $R^2 = 0.088$	Building type		
	(549) house (549)	Reference	
	(32) apartment (32)	1.679 (1.035–2.721)	0.036
	Degree of urbanisation		
	Rural (<600 inhab/km ²) (376)	Reference	
	Urban (>600 inhab/km ²) (205)	0.790 (0.626–0.996)	0.046
	Season		<0.001
	Winter (173)	Reference	
	Spring (201)	1.319 (0.999–1.742)	0.051
	Summer (66)	1.391 (0.947–2.042)	0.092
Autumn (141)	0.551 (0.408–0.744)	<0.001	
<u>5-HO-EHDPHP</u> $R^2 = 0.153$	Building type		
	House (509)	Reference	
	Apartment (25)	1.422 (1.023–1.974)	0.036
	Insulation/building status		0.05
Others (428)	Reference		
	0.768 (0.612–0.965)	0.023	

Table 3 (continued)

Log adjusted PFR metabolite	Variable (n)	β (95%CI)	p-Value
<u>EHPHP</u> $R^2 = 0.119$	Newly built after 2006, but no energy-neutral building (56)	Reference	
	Energy-neutral building (50)	1.082 (0.856–1.368)	0.511
	Moment of urine collection		<0.001
	<10 h (111)	Reference	
	10–12 h (214)	1.080 (0.899–1.298)	0.409
	>12 h (209)	1.493 (1.240–1.800)	<0.001
	Season		<0.001
	Winter (156)	Reference	
	Spring (185)	0.603 (0.507–0.717)	<0.001
	Summer (64)	0.677 (0.536–0.856)	0.001
Autumn (129)	0.957 (0.794–1.153)	0.641	
<u>BBOEHP</u> $R^2 = 0.052$	Season		<0.001
	Winter (173)	Reference	
	Spring (202)	0.807 (0.694–0.936)	0.004
	Summer (66)	0.839 (0.683–1.033)	0.071
	autumn (141)	1.251 (1.064–1.471)	0.008
	BMI		0.019
	<18.5 (159)	Reference	
	18.5–25 (359)	0.937 (0.818–1.073)	0.344
	>25 (64)	0.740 (0.600–0.914)	0.005
	Highest education level of the father		0.02
Primary education (65)	Reference		
Secondary education (228)	1.309 (0.983–1.742)	0.065	
Tertiary education (254)	1.484 (1.142–1.968)	0.006	
Season		0.004	
Winter (162)	Reference		
Spring (189)	0.785 (0.630–0.977)	0.031	
Summer (63)	1.114 (0.824–1.507)	0.481	
Autumn (133)	0.703 (0.554–0.891)	0.004	

PFRs as seasonality has also been observed in PFR concentrations in indoor dust and outdoor urban air (Cao et al., 2014; Wang et al., 2019a). Temperature and UV radiation can not only change the emission rates of PFRs from applications into the environment, but also the partitioning of PFRs between air and dust (Cao et al., 2014). As such, exposure through inhalation of contaminated air might be higher for more volatile PFRs such as TCIPP and TDCIPP. Indirectly, temperature could also influence the behavior of people with regard to their diet, the clothes they wear, ventilation habits or time spent indoor which, in turn, would result in more frequent exposure to PFRs through different pathways (Hoffman et al., 2018; Bastiaensen et al., 2020a).

BDCIPP and BBOEHP concentrations were higher in adolescents with mothers or fathers with tertiary education. The same trend has been reported for BBOEP, another TBOEP metabolite, in Japanese children (Bastiaensen et al., 2019) and for BCIPP but not BCIPHIPP in children from the US (Hoffman et al., 2018). Here, DPHP was also associated with education level of the parents, but the trend was different (higher levels for parents with secondary education). These results suggest that educational attainment is a determinant of PFR exposure, but this is likely a proxy for another underlying factor, which could be the use of specific products in these households (such as

Table 4

Estimated daily intakes (EDI in ng/kg bw/day) to organophosphate flame retardants and plasticizers calculated based on urinary metabolite concentrations.

EDIs in ng/kg bw/day	TPHP		TDCIPP		TCIPP		BBOEP		EHDPHP	
	Based on DPHP		Based on BDCIPP		Based on BCIPHIP		Based on BBOEHEP		Based on EHPHP + 5-HO-EHDPHP ^a	
	S9 ^b	HLM ^b	S9 ^b	HLM ^b	S9 ^b	HLM ^b	S9 ^b	HLM ^b	High F _{ue} ^c	Low F _{ue} ^c
F _{UE}	0.19	0.41	0.68	0.46	0.28	0.33	0.16	0.81	0.2	0.8
min	8.3	3.8	0.1	0.2	0.3	0.2	0.1	0.02	12.0	3.0
5th per	64.3	29.8	0.8	1.2	8.1	6.9	1.2	0.3	194.3	48.6
25th per	124.1	57.5	6.4	9.4	26.2	22.2	3.6	0.7	356.7	89.2
50th per	197.7	91.6	14.5	21.4	57.4	48.7	6.3	1.2	567.7	141.9
75th per	310.9	144.1	29.5	43.6	126.2	107.1	11.9	2.4	905.6	226.4
95th per	640.0	296.6	119.1	176.1	517.5	439.1	44.2	8.7	2004.2	501.1
max	8080.9	3744.8	1958.9	2895.7	11282.4	9573.0	246.3	48.6	3308.2	827.1
Medians										
Male	216.3	100.3	14.3	21.1	61.3	52.1	6.0	1.2	562.1	140.5
Female	181.4	84.1	14.7	21.8	51.3	43.5	6.4	1.3	570.8	142.7
BMI < 18.5	249.5	115.6	18.8	27.7	70.0	59.4	7.6	1.5	694.4	173.6
BMI 18.5–25	188.6	87.4	13.8	20.4	50.7	43.0	5.6	1.1	562.1	140.5
BMI > 25	121.6	56.4	11.0	16.3	48.0	40.7	5.4	1.1	323.3	80.8
RfD (ng/kg bw/day) ^d	2.0 × 10 ⁴		2.0 × 10 ⁴		1.0 × 10 ⁴		1.5 × 10 ⁴		1.5 × 10 ⁴	
% >RfD	0%		0%		0.17% (1/582)		0%		0%	
Ratio RfD/95th per	31	67	168	114	19	23	340	1720	7	30

HLM: human liver microsome fraction; S9: S9 fraction; bw: bodyweight; per: percentile.

^b F_{ue} estimations taken from Van den Eede et al. (2013).

^c No F_{ue} available in literature, therefore low and high F_{ue} scenario.

^d Oral reference doses (RfDs) taken from Poma et al. (2019).

furniture, floor covering or electronic devices). However, the reasons remain unknown as shown in the results of this determinant analysis.

Other variables included in the final models were more heterogeneous between compounds and are discussed individually below. Adolescents with higher BMI (>25 kg/m²) had significantly lower levels of DPHP and EHPHP. Wang et al. (2019b) reported that levels of DPHP, BDCIPP and BCIPP were higher in obese individuals, whereas BBOEP levels were lower in obese individuals than those with normal BMI. No differences in PFR metabolite levels were found between boys and girls. BDCIPP concentrations were significantly higher in adolescents living in homes with new decoration such as curtains, carpets and wall papers. TDCIPP is mainly applied in upholstered furniture, but it is possible that it is also used as a flame retardant in decorative materials (Wei et al., 2015). We also found that DPHP levels were lower for adolescents that live in homes with mechanical ventilation. This would suggest that ventilation decreases the concentration of TPHP in indoor dust or air. However, in a previous study by our research group, an opposite trend was observed consistent across multiple PFR metabolites (higher BCIPHIP, DPHP, BDCIPP, BBOEHEP and BBOEP levels with more frequent ventilation) (Bastiaensen et al., 2020a). In this study, ventilation was only significant for DPHP. All other metabolites were not associated with any of the ventilation variables, neither in univariate analysis (Table SI-4). Concentrations of 5-HO-EHDPHP were lower in adolescents living in a recently constructed home, which could suggest that newer insulation or building materials would not or contain less EHDPHP. Although EHDPHP is possibly applied in building materials as part of PVC polymers, no public data exist on the type of products in which this PFR is used (de la Torre et al., 2020). Finally, there were two significant determinants of TCIPP exposure for which no reasonable explanation could be found: building type (higher levels for adolescents living in apartments) and degree of urbanization (lower levels for adolescents living in urban areas).

Despite the extensive list of available questions and large sample size, the overall proportion of variance predicted by significant variables in multiple models was low (max R² = 0.153). Because various PFRs are often applied in similar applications (some even within the same product), we hypothesized that some associations would be similar for multiple metabolites. However, there were no consistent associations for characteristics of the indoor environment such as the use of building

materials, electronic devices or ventilation. Furthermore, none of the investigated food variables were significant determinants of exposure. Other studies have reported similar difficulties to relate food intake with urinary metabolite concentrations (Thomas et al., 2017a, 2017b; Bastiaensen et al., 2020a). One possible explanation for the lack of association with food is that the questions of this study focused on fresh foods while dietary exposure mainly originates from industrially processed food (such as grains, oils and dairy products) (Poma et al., 2018). Finally, the low predictive value of the constructed models may also be attributable to the short-term variability of PFR metabolite concentrations in urine. In this study, random spot samples were collected at school (9–16 h). Sampling time was a significant predictor for 5-HO-EHDPHP but not for other PFR biomarkers (Table 3). Spot urine sample analysis may not accurately represent exposure due to within-individual variation as a result of the short-biological half-life of PFRs (Wei et al., 2015). However, intraclass correlation coefficient analysis of PFR metabolites in spot urine samples has shown fair-to-good reproducibility of most PFR metabolites with ICC ranging from 0.35 for BBOEP to 0.68 for BDCIPP over 5 weeks (Wang et al., 2019b) and from 0.40 for 5-HO-EHDPHP to 0.60 for BDCIPP over 5 consecutive days, except for DPHP and EHPHP (ICC of 0.30 and 0.23, respectively) (Bastiaensen et al., 2020b).

3.3. Estimated daily exposure

A major advantage of human biomonitoring is that it provides an overall view of the internal dose and, thus, that it encompasses all exposure pathways. However, data on the bioavailability and toxicokinetics of PFRs are still very limited which limits the performance of a thorough risk assessment.

The daily intakes calculated here should therefore be considered as an estimation and the interpretation should be done cautiously. We opted to discuss only the EDI results based on the F_{UE}-S9-fraction as it contains both phase I and phase II metabolites (contrary to HLM that only contain phase I enzymes) (Richardson et al., 2016). This is of importance since some PFR metabolites are excreted as glucuronide conjugates (Van den Eede et al., 2013).

The highest median EDI was observed for EHDPHP_{highFue} (567.7 ng/kg bw/day). Median EDIs of other PFRs were a factor 10 (TPHP, TDCIPP,

TCIPP) or 100 lower (TBOEP). For TPHP, TDCIPP and TCIPP, the maximum EDIs calculated were several times higher than the 95th percentile, which indicates that some participants had extreme levels of these metabolites in their urine. As reported above, the total daily intake of TPHP and EHDHPH was related to BMI but not to sex (see descriptive results in Table 4). The calculated median EDI for TDCIPP was similar to those reported for adults and children from China and the US based on urinary metabolite levels (Chen et al., 2018; Zhang et al., 2018b; Li et al., 2019b; Wang et al., 2019b). EDIs for TPHP, TCIPP and TBOEP were more heterogeneous between the available studies, but generally within the same range. For EHDHPH however, the median EDI was higher than the one reported for Chinese adults (Li et al., 2019b).

EDIs of this study were much higher compared to results from external exposure assessments for Belgian adults (not available for adolescents). The mean daily intake for Belgian adults via food intake was estimated at 46.6 ng/kg bw/day for TPHP, 9.6 for TDCIPP, 18.5 for TCIPP and 14.9 for EHDHPH (Poma et al., 2018). The median intakes of PFRs by dust ingestion for Belgian adults were between a factor 10 and 100 lower compared to the estimate for food intake (0.1 ng/kg bw/day for TPHP, 0.3 for TDCIPP, 0.5 for TCIPP, 0.7 for TBOEP and 0.1 for EHDHPH) (Van den Eede et al., 2011; de la Torre et al., 2020). However, even when intake from dust and food are combined, the total intake accounted for a fraction of the median intake calculated based on urinary metabolite concentrations (24% for TPHP_{S9}, 69% for TDCIPP_{S9}, 33% for TCIPP_{S9} and 3% for EHDHPH_{highFue}). These results suggested that other exposure pathways such as air inhalation and dermal absorption could contribute considerably to PFR exposure and that some sources of exposure may still be unknown such as direct exposure to diesters (Tan et al., 2019).

Oral reference doses (RfD) for human exposure were used according to Poma et al. (2019) to assess the potential risk associated with the estimated intakes. EDIs of all participants, except one for TCIPP_{S9}, were well below the RfDs of the investigated PFRs (Table 4). More precisely, the 95th percentile EDI (high exposure scenario) was 7 (EHDHPH_{highFue}) to 1720 (TBOEP_{HLM}) times lower than the RfD. Overall, these results suggest that the PFR exposure of Belgian adolescents is well below the risk threshold. However, it is important to note that these RfD values were determined based on toxicological data and should be interpreted within the context of their limitations. When RfDs are revised according to new toxicological evidence they may become stricter. Furthermore, the assessment presented here does also not account for the potential additive effect on health of combined exposure to multiple PFRs and/or other toxic chemicals. Finally, due to the lack of *in vivo* toxicokinetic data of PFRs in humans, the calculated EDI may be over- or underestimated. The toxic characteristics of most PFRs are relatively well documented in *in vitro* and animal studies but reports on the health effects of background levels on humans are limited. Urinary concentrations of PFR metabolites within the same range as the levels observed in this study have been associated with allergic symptoms and oxidative stress biomarkers in Japanese children (Araki et al., 2018; Ait Bamai et al., 2019). EHDHPH concentrations were also significantly correlated with sphingolipid levels in human blood (Zhao et al., 2016). Although these association studies may not be representative to adolescents or other age groups, they do suggest a relationship between exposure and effect. Additional epidemiological research is needed to understand sources of exposure as well as the potential health impact of PFR exposure.

4. Conclusions

This is the first study to report reference values for 13 PFR urinary metabolites originating from 7 parent compounds, determined in European adolescents. Metabolites of EHDHPH, TCIPP, TDCIPP, TPHP and TBOEP were detectable in practically every participant suggesting widespread exposure to PFRs in Flanders, Belgium. Concentrations of DPHP, BDCIPP and BCIPHIPP were generally within the same range as

in other study populations from non-European countries, only EHDHPH exposure was presumably higher in this study. However, we could not identify any explanatory factors related to the ingestion of EHDHPH or other PFRs through food. Determinant analysis did show higher levels of BDCIPP with the introduction of new decoration, lower levels of 5-HO-EHDHPH in adolescents living in newly built homes and higher levels of BBOEHP and BDCIPP in adolescents with highly educated parents. Season was the only determinant significant for several PFR metabolites. The estimated daily intakes for Flemish adolescents were all well below the available reference doses indicating that the risk associated to the measured internal doses is low.

CRedit authorship contribution statement

Michiel Bastiaensen: Conceptualization, Methodology, Formal analysis, Investigation, Writing - original draft, Visualization. **Celine Gys:** Conceptualization, Investigation, Writing - review & editing, Visualization. **Ann Colles:** Conceptualization, Investigation, Writing - review & editing. **Veerle Verheyen:** Conceptualization, Writing - review & editing. **Gudrun Koppen:** Conceptualization, Writing - review & editing. **Eva Govarts:** Conceptualization, Investigation, Writing - review & editing. **Liesbeth Bruckers:** Conceptualization, Investigation, Writing - review & editing. **Bert Morrens:** Conceptualization, Writing - review & editing. **Ilse Loots:** Conceptualization, Writing - review & editing. **Annelies De Decker:** Conceptualization, Writing - review & editing. **Vera Nelen:** Conceptualization, Writing - review & editing. **Tim Nawrot:** Conceptualization, Writing - review & editing. **Stefaan De Henauw:** Conceptualization, Writing - review & editing. **Nik Van Larbeke:** Conceptualization, Writing - review & editing. **Greet Schoeters:** Conceptualization, Methodology, Resources, Writing - review & editing, Supervision, Project administration, Funding acquisition. **Adrian Covaci:** Conceptualization, Methodology, Resources, Writing - review & editing, Supervision, Project administration, Funding acquisition.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We thank the participating adolescents and their families. Without their effort this study would not have been possible. We also thank J. Bombeke for his assistance during the sample preparation. This paper is based on research conducted within the framework of the Flemish Center of Expertise on Environment and Health (FLEHS 2016-2020), funded by the Government of Flanders, Department of Environment & Spatial Development. The views expressed herein are those of the author (s) and are not necessarily endorsed by the government of Flanders. The Flemish Environment and Health Study was commissioned, financed and steered by the Ministry of the Flemish Community (Department of Environment) (grant ID: Steunpunt Milieu & Gezondheid), including the partial funding of the Ph.D. of Michiel Bastiaensen. Celine Gys acknowledges the funding of a PhD fellowship from Research Foundation Flanders (project GOE5216N).

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2020.106368>.

References

Ait Bamai, Y.u., Bastiaensen, M., Araki, A., Goudarzi, H., Konno, S., Ito, S., Miyashita, C., Yao, Y., Covaci, A., Kishi, R., 2019. Multiple exposures to organophosphate flame

- retardants alter urinary oxidative stress biomarkers among children: The Hokkaido Study. *Environ. Int.* 131, 105003.
- Araki, A., Michiel Bastiaensen, Y.u., Bamai, A., Van den Eede, N., Kawai, T., Tsuboi, T., Ketema, R.M., Covaci, A., Kishi, R., 2018. Associations between allergic symptoms and phosphate flame retardants in dust and their urinary metabolites among school children. *Environ. Int.* 119, 438–446.
- Ballesteros-Gómez, A., Erratico, C.A., Van den Eede, N., Ionas, A.C., Leonards, P.E.G., Covaci, A., 2015. In vitro metabolism of 2-ethylhexyldiphenyl phosphate (EHDPHP) by human liver microsomes. *Toxicol. Lett.* 232, 203–212.
- Bastiaensen, M., Bamai, Y.A., Araki, A., Goudarzi, H., Konno, S., Ito, S., Miyashita, C., Yao, Y., Kishi, R., Covaci, A., 2020a. Temporal trends and determinants of PFR exposure in the Hokkaido Study. *Int. J. Hyg. Environ. Health* 228, 113523.
- Bastiaensen, M., Bamai, Y.A., Araki, A., Van den Eede, N., Kawai, T., Tsuboi, T., Kishi, R., Covaci, A., 2019. Biomonitoring of organophosphate flame retardants and plasticizers in children: Associations with house dust and housing characteristics in Japan. *Environ. Res.* 172, 543–551.
- Bastiaensen, Michiel, Gys, Celine, Malarvannan, Govindan, Fotache, Mihai, Jasper Bombeke, Yu, Bamai, Ait, Araki, Atsuko, Covaci, Adrian, 2020b. Short-term temporal variability of urinary biomarkers of organophosphate flame retardants and plasticizers. *Environ. Int.* 146.
- Bastiaensen, M., Fuchao, X.u., Been, F., Van den Eede, N., Covaci, A., 2018. Simultaneous determination of 14 urinary biomarkers of exposure to organophosphate flame retardants and plasticizers by LC-MS/MS. *Anal. Bioanal. Chem.* 410, 7871–7880.
- Brommer, S., Harrad, S., 2015. Sources and human exposure implications of concentrations of organophosphate flame retardants in dust from UK cars, classrooms, living rooms, and offices. *Environ. Int.* 83, 202–207.
- Butt, C.M., Hoffman, K., Chen, A., Lorenzo, A., Congleton, J., Stapleton, H.M., 2016. Regional comparison of organophosphate flame retardant (PFR) urinary metabolites and tetrabromobenzoic acid (TBBA) in mother-toddler pairs from California and New Jersey. *Environ. Int.* 94, 627–634.
- Cao, Z., Fuchao, X.u., Covaci, A., Wu, Min, Yu, Gang, Wang, B., Deng, S., Huang, J., 2014. Differences in the seasonal variation of brominated and phosphorus flame retardants in office dust. *Environ. Int.* 65, 100–106.
- Carignan, Courtney C., Mínguez-Alarcón, Lidia, Butt, Craig M., Williams, Paige L., Meeker, John D., Stapleton, Heather M., Toth, Thomas L., Ford, Jennifer B., Hauser, Russ, EARTH Study Team, 2017. Urinary concentrations of organophosphate flame retardant metabolites and pregnancy outcomes among women undergoing in vitro fertilization. *Environ. Health Perspect.* 125, 087018.
- Cequier, Enrique, Ionas, Alin C., Covaci, Adrian, Marcé, Rosa Maria, Becher, Georg, Thomsen, Cathrine, 2014. Occurrence of a broad range of legacy and emerging flame retardants in indoor environments in Norway. *Environ. Sci. Technol.* 48, 6827–6835.
- Cequier, Enrique, Sakhi, Amrit Kaur, Marcé, Rosa Maria, Becher, Georg, Thomsen, Cathrine, 2015. Human exposure pathways to organophosphate triesters—a biomonitoring study of mother-child pairs. *Environ. Int.* 75, 159–165.
- Chen, Yi, Fang, Jianzhang, Ren, Lu, Fan, Ruirang, Zhang, Jianqing, Liu, Guihua, Zhou, Li, Chen, Dingyan, Yu, Yingxin, Lu, Shaoyou, 2018. Urinary metabolites of organophosphate esters in children in South China: Concentrations, profiles and estimated daily intake. *Environ. Pollut.* 235, 358–364.
- Chen, Yi, Jiang, Lei, Shaoyou, Lu, Kang, Li, Luo, Xianru, Liu, Guihua, Cui, Xinyi, Yu, Yingxin, 2019. Organophosphate ester and phthalate ester metabolites in urine from primiparas in Shenzhen, China: Implications for health risks. *Environ. Pollut.* 247, 944–952.
- de la Torre, Adrián, Navarro, Irene, Sanz, Paloma, de los Ángeles Martínez, María, 2020. Organophosphate compounds, polybrominated diphenyl ethers and novel brominated flame retardants in European indoor house dust: Use, evidence for replacements and assessment of human exposure. *J. Hazard. Mater.* 382, 121009.
- Ding, Jinjian, Deng, Tongqing, Ye, Xiaoqing, Covaci, Adrian, Liu, Jing, Yang, Fangxing, 2019. Urinary metabolites of organophosphate esters and implications for exposure pathways in adolescents from Eastern China. *Sci. Total Environ.* 695, 133894.
- Dodson, Robin E., Perovich, Laura J., Covaci, Adrian, Van den Eede, Nele, Ionas, Alin C., Dirtu, Alin C., Brody, Julia Green, Rudel, Ruthann A., 2012. After the PBDE phase-out: a broad suite of flame retardants in repeat house dust samples from California. *Environ. Sci. Technol.* 46, 13056–13066.
- Dodson, Robin E., Van den Eede, Nele, Covaci, Adrian, Perovich, Laura J., Brody, Julia Green, Rudel, Ruthann A., 2014. Urinary biomonitoring of phosphate flame retardants: levels in California adults and recommendations for future studies. *Environ. Sci. Technol.* 48, 13625–13633.
- ECHA, 2020. Information on Chemicals. <https://echa.europa.eu/nl/information-on-chemicals>.
- EU, 2009. European Union Risk Assessment Report: Tris(2-chloroethyl) Phosphate, TCEP. European Union.
- Fromme, H., Lahrz, T., Kraft, M., Fembacher, L., Mach, C., Dietrich, S., Burkardt, R., Völkel, W., Göen, T., 2014. Organophosphate flame retardants and plasticizers in the air and dust in German daycare centers and human biomonitoring in visiting children (LUPE 3). *Environ. Int.* 71, 158–163.
- Gibson, Elizabeth A., Stapleton, Heather M., Calero, Lehyla, Holmes, Darrell, Burke, Kimberly, Martinez, Rodney, Cortes, Boris, Nematollahi, Amy, Evans, David, Herbstman, Julie B., 2018. Flame retardant exposure assessment: findings from a behavioral intervention study. *J. Exposure Sci. Environ. Epidemiol.* 1.
- Grand View Research, 2017. Global flame retardant market projected to reach US\$11.96 billion by 2025. *Addit. Polym.* 2017, 10–11.
- Hammel, Stephanie C., Hoffman, Kate, Webster, Thomas F., Anderson, Kim A., Stapleton, Heather M., 2016. Measuring personal exposure to organophosphate flame retardants using silicone wristbands and hand wipes. *Environ. Sci. Technol.* 50, 4483–4491.
- He, Chang, English, Karin, Baduel, Christine, Thai, Phong, Jagals, Paul, Ware, Robert S., Li, Yan, Wang, Xianyu, Sly, Peter D., Mueller, Jochen F., 2018a. Concentrations of organophosphate flame retardants and plasticizers in urine from young children in Queensland, Australia and associations with environmental and behavioural factors. *Environ. Res.* 164, 262–270.
- He, Chang, Toms, Leisa-Maree L., Thai, Phong, Van den Eede, Nele, Wang, Xianyu, Li, Yan, Baduel, Christine, Harden, Fiona A., Heffernan, Amy L., Hobson, Peter, Covaci, Adrian, Mueller, Jochen F., 2018b. Urinary metabolites of organophosphate esters: Concentrations and age trends in Australian children. *Environ. Int.* 111, 124–130.
- Hoffman, Kate, Butt, Craig M., Webster, Thomas F., Preston, Emma V., Hammel, Stephanie C., Makey, Colleen, Lorenzo, Amelia M., Cooper, Ellen M., Carignan, Courtney, Meeker, John D., 2017a. Temporal trends in exposure to organophosphate flame retardants in the United States. *Environ. Sci. Technol. Lett.* 4 (3), 112–118.
- Hoffman, Kate, Hammel, Stephanie C., Phillips, Allison L., Lorenzo, Amelia M., Chen, Albert, Calafat, Antonia M., Ye, Xiaoyun, Webster, Thomas F., Stapleton, Heather M., 2018. Biomarkers of exposure to SVOCs in children and their demographic associations: The TESIE Study. *Environ. Int.* 119, 26–36.
- Hoffman, Kate, Lorenzo, Amelia, Butt, Craig M., Hammel, Stephanie C., Henderson, Brittany Bohinc, Roman, Sanziana A., Scheri, Randall P., Stapleton, Heather M., Sosa, Julie Ann, 2017b. Exposure to flame retardant chemicals and occurrence and severity of papillary thyroid cancer: A case-control study. *Environ. Int.* 107, 235–242.
- Hu, Liqin, Tao, Yun, Luo, Dan, Feng, Jingwen, Wang, Limei, Yu, Meng, Li, Yaping, Covaci, Adrian, Mei, Surong, 2019. Simultaneous biomonitoring of 15 organophosphate flame retardants metabolites in urine samples by solvent induced phase transition extraction coupled with ultra-performance liquid chromatography-tandem mass spectrometry. *Chemosphere* 233, 724–732.
- Ingle, Mary E., Mínguez-Alarcón, Lidia, Carignan, Courtney C., Butt, Craig M., Stapleton, Heather M., Williams, Paige L., Ford, Jennifer B., Hauser, Russ, Meeker, John D., EARTH Study Team, 2018. The association between urinary concentrations of phosphorous-containing flame retardant metabolites and semen parameters among men from a fertility clinic. *Int. J. Hyg. Environ. Health* 221, 809–815.
- Kim, Un-Jung, Yu, Wang, Li, Wenhui, Kannan, Kurunthachalam, 2019. Occurrence of and human exposure to organophosphate flame retardants/plasticizers in indoor air and dust from various microenvironments in the United States. *Environ. Int.* 125, 342–349.
- Li, Jianhua, Zhao, Luming, Letcher, Robert J., Zhang, Yayun, Jian, Kang, Zhang, Jinhua, Su, Guanyong, 2019a. A review on organophosphate Ester (OPE) flame retardants and plasticizers in foodstuffs: Levels, distribution, human dietary exposure, and future directions. *Environ. Int.* 127, 35–51.
- Li, Mengqi, Yao, Yiming, Wang, Yu, Bastiaensen, Michiel, Covaci, Adrian, Sun, Hongwen, 2019b. Organophosphate ester flame retardants and plasticizers in a Chinese population: Significance of hydroxylated metabolites and implication for human exposure. *Environ. Pollut.* 113633.
- Li, Wenhui, Wang, Yu, Asimakopoulos, Alexandros G., Covaci, Adrian, Gevao, Bondi, Johnson-Restrepo, Boris, Kumosani, Taha A., Malarvannan, Govindan, Moon, Hyo-Bang, Nakata, Haruhiko, 2019c. Organophosphate esters in indoor dust from 12 countries: Concentrations, composition profiles, and human exposure. *Environ. Int.* 133, 105178.
- Ospina, Maria, Jayatilaka, Nayana K., Wong, Lee-Yang, Restrepo, Paula, Calafat, Antonia M., 2018. Exposure to organophosphate flame retardant chemicals in the US general population: Data from the 2013–2014 National Health and Nutrition Examination Survey. *Environ. Int.* 110, 32–41.
- Phillips, Allison L., Hammel, Stephanie C., Hoffman, Kate, Lorenzo, Amelia M., Chen, Albert, Webster, Thomas F., Stapleton, Heather M., 2018. Children's residential exposure to organophosphate ester flame retardants and plasticizers: Investigating exposure pathways in the TESIE study. *Environ. Int.* 116, 176–185.
- Poma, Giulia, Glynn, Anders, Malarvannan, Govindan, Covaci, Adrian, Darnerud, Per Ola, 2017. Dietary intake of phosphorus flame retardants (PFRs) using Swedish food market basket estimations. *Food Chem. Toxicol.* 100, 1–7.
- Poma, Giulia, Sales, Carlos, Bruyland, Bram, Christia, Christina, Goscinny, Séverine, Van Loco, Joris, Covaci, Adrian, 2018. Occurrence of Organophosphorus Flame Retardants and Plasticizers (PFRs) in Belgian Foodstuffs and Estimation of the Dietary Exposure of the Adult Population. *Environ. Sci. Technol.* 52, 2331–2338.
- Poma, Giulia, Yin, Shanshan, Tang, Bin, Fujii, Yukiko, Cuykx, Matthias, Covaci, Adrian, 2019. Occurrence of selected organic contaminants in edible insects and assessment of their chemical safety. *Environ. Health Perspect.* 127, 127009.
- Richardson, Samantha, Bai, April, Kulkarni, Ashutosh, Moghaddam, Mehran, 2016. Efficiency in drug discovery: liver S9 fraction assay as a screen for metabolic stability. *Drug Metab. Lett.* 10, 83–90.
- Schoeters, Greet, Den Hond, Elly, Colles, Ann, Loots, Ilse, Morrens, Bert, Keune, Hans, Bruckers, Liesbeth, Nawrot, Tim, Sioen, Isabelle, De Coster, Sam, 2012. Concept of the Flemish human biomonitoring programme. *Int. J. Hyg. Environ. Health* 215, 102–108.
- Siddique, Shabana, Harris, Shelley A., Kosarac, Ivana, Latifovic, Lidija, Kubwabo, Cariton, 2020. Urinary metabolites of organophosphate esters in women and its relationship with serum lipids: An exploratory analysis. *Environ. Pollut.* 114110.
- Stapleton, Heather M., Klosterhaus, Susan, Eagle, Sarah, Fuh, Jennifer, Meeker, John D., Blum, Arlene, Webster, Thomas F., 2009. Detection of organophosphate flame retardants in furniture foam and US house dust. *Environ. Sci. Technol.* 43, 7490–7495.

- Sun, Yan, Gong, Xia, Lin, Wanlong, Liu, Ye, Wang, Yujie, Minghong, Wu., Kannan, Kurunthachalam, Ma, Jing, 2018. Metabolites of organophosphate ester flame retardants in urine from Shanghai, China. *Environ. Res.* 164, 507–515.
- Tan, Hongli, Yang, Liu, Yu, Yunjiang, Guan, Qingxia, Liu, Xiaotu, Li, Liangzhong, Da, Chen, 2019. Co-existence of organophosphate di- and tri-esters in house dust from South China and Midwestern United States: implications for human exposure. *Environ. Sci. Technol.* 53, 4784–4793.
- Thomas, M.B., Stapleton, H.M., Dills, R.L., Violette, H.D., Christakis, D.A., Sathyanarayana, S., 2017a. Demographic and dietary risk factors in relation to urinary metabolites of organophosphate flame retardants in toddlers. *Chemosphere* 185, 918–925.
- UNEP, 2009. Decision SC-4/18: Listing of tetrabromodiphenyl ether and pentabromodiphenyl ether. In: Edited by Stockholm Convention on Persistent Organic Pollutants. Geneva, Switzerland: United Nations Environment Programme.
- UNEP, 2017. Decision SC-8/10: Listing of decabromodiphenyl ether. In: Edited by Stockholm Convention on Persistent Organic Pollutants. Geneva, Switzerland: United Nations Environment Programme.
- Valentin, Jack, 2002. Basic anatomical and physiological data for use in radiological protection: reference values: ICRP Publication 89. *Ann. ICRP* 32, 1–277.
- den Eede, Van, Nele, Alin C., Dirtu, Hugo Neels, Covaci, Adrian, 2011. Analytical developments and preliminary assessment of human exposure to organophosphate flame retardants from indoor dust. *Environ. Int.* 37, 454–461.
- den Eede, Van, Nele, Amy L., Heffernan, Lesa L., Aylward, Peter Hobson, Neels, Hugo, Mueller, Jochen F, Covaci, Adrian, 2015. Age as a determinant of phosphate flame retardant exposure of the Australian population and identification of novel urinary PFR metabolites. *Environ. Int.* 74, 1–8.
- den Eede, Van, Nele, Walid Maho, Erratico, Claudio, Neels, Hugo, Covaci, Adrian, 2013. First insights in the metabolism of phosphate flame retardants and plasticizers using human liver fractions. *Toxicol. Lett.* 223, 9–15.
- van der Veen, Ike, de Boer, Jacob, 2012. Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 88, 1119–1153.
- Völkel, Wolfgang, Fuchs, Veronika, Wöckner, Mandy, Fromme, Hermann, 2018. Toxicokinetic of tris (2-butoxyethyl) phosphate (TBOEP) in humans following single oral administration. *Arch. Toxicol.* 92, 651–660.
- Wang, Dou, Pu, Wang, Wang, Yiwen, Zhang, Weiwei, Zhu, Chaofei, Sun, Huizhong, Matsiko, Julius, Zhu, Ying, Li, Yingming, Meng, Wenying, 2019a. Temporal variations of PM_{2.5}-bound organophosphate flame retardants in different microenvironments in Beijing, China, and implications for human exposure. *Sci. Total Environ.* 666, 226–234.
- Wang, Yu, Li, Wenhui, Martínez-Moral, María Pilar, Sun, Hongwen, Kannan, Kurunthachalam, 2019b. Metabolites of organophosphate esters in urine from the United States: Concentrations, temporal variability, and exposure assessment. *Environ. Int.* 122, 213–221.
- Wei, Gao-Ling, Li, Ding-Qiang, Zhuo, Mu-Ning, Liao, Yi-Shan, Xie, Zhen-Yue, Guo, Tai-Long, Li, Jun-Jie, Zhang, Si-Yi, Liang, Zhi-Quan, 2015. Organophosphorus flame retardants and plasticizers: sources, occurrence, toxicity and human exposure. *Environ. Pollut.* 196, 29–46.
- Xu, Fuchao, Eulaers, Igor, Alves, Andreia, Papadopoulou, Eleni, Padilla-Sanchez, Juan Antonio, Lai, Foon Yin, Haug, Line Småstuen, Voorspoels, Stefan, Neels, Hugo, Covaci, Adrian, 2019. Human exposure pathways to organophosphate flame retardants: Associations between human biomonitoring and external exposure. *Environ. Int.* 127, 462–472.
- Xu, Fuchao, Giovanoulis, Georgios, Van Waes, Sofie, Padilla-Sanchez, Juan Antonio, Papadopoulou, Eleni, Magnér, Jorgen, Haug, Line Småstuen, Neels, Hugo, Covaci, Adrian, 2016. Comprehensive Study of Human External Exposure to Organophosphate Flame Retardants via Air, Dust, and Hand Wipes: The Importance of Sampling and Assessment Strategy. *Environ. Sci. Technol.* 50, 7752–7760.
- Xu, Fuchao, Tay, Joo-Hui, Covaci, Adrian, Padilla-Sánchez, Juan Antonio, Papadopoulou, Eleni, Haug, Line Småstuen, Neels, Hugo, Sellström, Ulla, de Wit, Cynthia A, 2017. Assessment of dietary exposure to organohalogen contaminants, legacy and emerging flame retardants in a Norwegian cohort. *Environ. Int.* 102, 236–243.
- Zhang, Bo, Lu, Shaoyou, Huang, Mingzhi, Zhou, Meizhou, Zhou, Ziqing, Zheng, Hangcong, Jiang, Yongchen, Bai, Xueyuan, Zhang, Tao, 2018a. Urinary metabolites of organophosphate flame retardants in 0–5-year-old children: Potential exposure risk for inpatients and home-stay infants. *Environ. Pollut.* 243, 318–325.
- Zhang, Tao, Bai, Xue-yuan, Lu, Shao-you, Zhang, Bo, Xie, Lei, Zheng, Hang-cong, Jiang, Yong-chen, Zhou, Mei-zhou, Zhou, Zi-qing, Song, Shi-ming, 2018b. Urinary metabolites of organophosphate flame retardants in China: health risk from tris (2-chloroethyl) phosphate (TCEP) exposure. *Environ. Int.* 121, 1363–1371.
- Zhao, Fanrong, Wan, Yi, Zhao, Haoqi, Hu, Wenxin, Mu, Di, Webster, Thomas F, Hu, Jianying, 2016. Levels of blood organophosphorus flame retardants and association with changes in human sphingolipid homeostasis. *Environ. Sci. Technol.* 50, 8896–8903.